

(Piecewise linear & birational) involutions on Dyck paths

Howard Mathematics Colloquium

Sam Hopkins

based on joint work with Michael Joseph (Dalton State College)

Howard University

September 24, 2021

Section 1

Catalan numbers, Dyck paths, Naryana numbers, and the Lalanne–Kreweras involution



Montserrat Mountain, Catalonia, Spain

Catalan numbers

The **Catalan numbers** C_n are a famous sequence of numbers

1, 2, 5, 14, 42, 132, 429, 1430, ...,

which count numerous combinatorial collections including:


Catalan numbers

The **Catalan numbers** C_n are a famous sequence of numbers


$$1, 2, 5, 14, 42, 132, 429, 1430, \dots,$$

which count numerous combinatorial collections including:


triangulations
of an $n + 2$ -gon



binary trees
with n nodes



plane trees with
 $n + 1$ nodes



bracketings of
 $n + 1$ terms

$$\begin{aligned} &a(b(cd)) \quad a((bc)d) \\ &(ab)(cd) \quad (a(bc))d \\ &((ab)c)d \end{aligned}$$

Catalan numbers

The **Catalan numbers** C_n are a famous sequence of numbers


$$1, 2, 5, 14, 42, 132, 429, 1430, \dots,$$

which count numerous combinatorial collections including:


triangulations
of an $n + 2$ -gon



binary trees
with n nodes



plane trees with
 $n + 1$ nodes



bracketings of
 $n + 1$ terms

$$\begin{aligned} &a(b(cd)) \quad a((bc)d) \\ &(ab)(cd) \quad (a(bc))d \\ &((ab)c)d \end{aligned}$$

There is a well-known product formula for the Catalan numbers:

$$C_n = \frac{1}{n+1} \binom{2n}{n} = \frac{(2n)!}{(n+1)!n!}$$

History of Catalan numbers

The Catalan numbers are named after Belgian mathematician *Eugène Catalan* (1814 – 1894), who studied them in conjunction with bracketings.

History of Catalan numbers

The Catalan numbers are named after Belgian mathematician *Eugène Catalan* (1814 – 1894), who studied them in conjunction with bracketings.

But they were studied combinatorially much earlier by *Leonhard Euler* (1707 – 1783), who showed they count triangulations of convex polygons.

History of Catalan numbers

The Catalan numbers are named after Belgian mathematician *Eugène Catalan* (1814 – 1894), who studied them in conjunction with bracketings.

But they were studied combinatorially much earlier by *Leonhard Euler* (1707 – 1783), who showed they count triangulations of convex polygons.

In fact, even earlier, Mongolian mathematician/scientist *Minggatu* (c.1692 – c.1763) used Catalan numbers in certain trigonometric identities.

History of Catalan numbers

The Catalan numbers are named after Belgian mathematician *Eugène Catalan* (1814 – 1894), who studied them in conjunction with bracketings.

But they were studied combinatorially much earlier by *Leonhard Euler* (1707 – 1783), who showed they count triangulations of convex polygons.

In fact, even earlier, Mongolian mathematician/scientist *Minggatu* (c.1692 – c.1763) used Catalan numbers in certain trigonometric identities.



E. Catalan



L. Euler



Minggatu

History of Catalan numbers

The Catalan numbers are named after Belgian mathematician *Eugène Catalan* (1814 – 1894), who studied them in conjunction with bracketings.

But they were studied combinatorially much earlier by *Leonhard Euler* (1707 – 1783), who showed they count triangulations of convex polygons.

In fact, even earlier, Mongolian mathematician/scientist *Minggatu* (c.1692 – c.1763) used Catalan numbers in certain trigonometric identities.



E. Catalan



L. Euler




Minggatu

It's a good thing the C_n are not named after Euler, since there are already

- *Euler numbers & Eulerian numbers*, counting certain permutations;
- *Euler's number* $e \approx 2.71$ & the *Euler–Mascheroni constant* $\gamma \approx 0.57$.


Catalan numbers: the book

Richard Stanley has a whole book devoted to the Catalan numbers.



Catalan numbers: the book

Richard Stanley has a whole book devoted to the Catalan numbers.



In it, he gives an astounding 214 different interpretations of C_n .


Dyck paths

The interpretation of C_n I want to focus on is in terms of Dyck paths.

Dyck paths

The interpretation of C_n I want to focus on is in terms of Dyck paths.


A **Dyck path** of length $2n$ is a lattice path in \mathbb{Z}^2 from $(0, 0)$ to $(2n, 0)$ consisting of n up steps $U = (1, 1)$ and n down steps $D = (1, -1)$ that never goes below the x -axis:



Dyck paths

The interpretation of C_n I want to focus on is in terms of Dyck paths.

A **Dyck path** of length $2n$ is a lattice path in \mathbb{Z}^2 from $(0, 0)$ to $(2n, 0)$ consisting of n up steps $U = (1, 1)$ and n down steps $D = (1, -1)$ that never goes below the x -axis:




The number of Dyck paths of length $2n$ is C_n :



Dyck paths

The interpretation of C_n I want to focus on is in terms of Dyck paths.

A **Dyck path** of length $2n$ is a lattice path in \mathbb{Z}^2 from $(0, 0)$ to $(2n, 0)$ consisting of n up steps $U = (1, 1)$ and n down steps $D = (1, -1)$ that never goes below the x -axis:



The number of Dyck paths of length $2n$ is C_n :



They are named after German algebraist *Walther von Dyck* (1856 – 1934).

Peaks and valleys in Dyck paths

Dyck paths look like mountain ranges. So we use some topographic terminology when working with Dyck paths.

Peaks and valleys in Dyck paths


Dyck paths look like mountain ranges. So we use some topographic terminology when working with Dyck paths.

A **peak** in a Dyck path is an up step that is immediately followed by a down step; a **valley** is a down step immediately followed by an up step.

Peaks and valleys in Dyck paths

Dyck paths look like mountain ranges. So we use some topographic terminology when working with Dyck paths.

A **peak** in a Dyck path is an up step that is immediately followed by a down step; a **valley** is a down step immediately followed by an up step.




Here the peaks are marked by red circles and the valleys by green circles.

Peaks and valleys in Dyck paths

Dyck paths look like mountain ranges. So we use some topographic terminology when working with Dyck paths.

A **peak** in a Dyck path is an up step that is immediately followed by a down step; a **valley** is a down step immediately followed by an up step.



Here the peaks are marked by red circles and the valleys by green circles.
It's easy to see that a Dyck path which has k valleys has $k + 1$ peaks.

Narayana numbers

The **Narayana number** $N(n, k)$ is the number of Dyck paths of length $2n$ with exactly k valleys.

Narayana numbers

The **Narayana number** $N(n, k)$ is the number of Dyck paths of length $2n$ with exactly k valleys.

$n \setminus k$	0	1	2	3
1	1			
2	1	1		
3	1	3	1	
4	1	6	6	1

← array of $N(n, k)$

Narayana numbers

The **Narayana number** $N(n, k)$ is the number of Dyck paths of length $2n$ with exactly k valleys.

$n \setminus k$	0	1	2	3
1	1			
2	1	1		
3	1	3	1	
4	1	6	6	1

\leftarrow array of $N(n, k)$

Evidently, the Narayana numbers $N(n, k)$ refine the Catalan number C_n :

$$C_n = \sum_{k=0}^{n-1} N(n, k).$$

Narayana numbers

The **Narayana number** $N(n, k)$ is the number of Dyck paths of length $2n$ with exactly k valleys.

$n \setminus k$	0	1	2	3
1	1			
2	1	1		
3	1	3	1	
4	1	6	6	1

\leftarrow array of $N(n, k)$

Evidently, the Narayana numbers $N(n, k)$ refine the Catalan number C_n :

$$C_n = \sum_{k=0}^{n-1} N(n, k).$$

They are named after Canadian mathematician/statistician *Tadepalli Venkata Narayana* (1930 – 1987), who in 1959 showed that

$$N(n, k) = \frac{1}{n} \binom{n}{k} \binom{n}{k+1}.$$

Symmetry of Narayana numbers

From Narayana's formula, it follows immediately that

$$N(n, k) = N(n, n - 1 - k)$$


for all k . That is, the sequence of Narayana numbers is *symmetric*.

Symmetry of Narayana numbers

From Narayana's formula, it follows immediately that

$$N(n, k) = N(n, n - 1 - k)$$

for all k . That is, the sequence of Narayana numbers is *symmetric*.




Symmetry of Narayana numbers

From Narayana's formula, it follows immediately that

$$N(n, k) = N(n, n - 1 - k)$$

for all k . That is, the sequence of Narayana numbers is *symmetric*.




However, it is not combinatorially obvious why the number of Dyck paths with k valleys should be the same as the number with $n - 1 - k$ valleys.

The Lalanne–Kreweras involution

The **Lalanne–Kreweras involution** (after *G. Kreweras* and *J.-C. Lalanne*) is a map on Dyck paths which combinatorially demonstrates the symmetry of the Narayana numbers: $\#\text{valleys}(\Gamma) + \#\text{valleys}(\text{LK}(\Gamma)) = n - 1$.

The Lalanne–Kreweras involution


The **Lalanne–Kreweras involution** (after G. Kreweras and J.-C. Lalanne) is a map on Dyck paths which combinatorially demonstrates the symmetry of the Narayana numbers: $\#\text{valleys}(\Gamma) + \#\text{valleys}(\text{LK}(\Gamma)) = n - 1$.



As depicted above, to compute the LK involution of a Dyck path Γ , we draw dashed lines emanating from the middle of every double up step and every double down step of Γ , at -45° and 45° respectively; these dashed lines intersect at the valleys of (an upside copy of) the Dyck path $\text{LK}(\Gamma)$.

The Lalanne–Kreweras involution

The **Lalanne–Kreweras involution** (after G. Kreweras and J.-C. Lalanne) is a map on Dyck paths which combinatorially demonstrates the symmetry of the Narayana numbers: $\#\text{valleys}(\Gamma) + \#\text{valleys}(\text{LK}(\Gamma)) = n - 1$.



As depicted above, to compute the LK involution of a Dyck path Γ , we draw dashed lines emanating from the middle of every double up step and every double down step of Γ , at -45° and 45° respectively; these dashed lines intersect at the valleys of (an upside copy of) the Dyck path $\text{LK}(\Gamma)$. That LK is an involution means $\text{LK}^2(\Gamma) = \Gamma$ for all Dyck paths Γ .

Section 2

Posets

Posets

We will now reinterpret the LK involution using the theory of finite posets.

Posets

We will now reinterpret the LK involution using the theory of finite posets.

A (finite) **poset**, or *partially ordered set*, is a (finite) set P together with a relation \leq satisfying the usual axioms of a partial order:

- *transitivity* ($x \leq y, y \leq z \Rightarrow x \leq z$);
- *anti-symmetry* ($x \leq y, y \leq x \Rightarrow x = y$);
- *reflexivity* ($x \leq x$).


Posets

We will now reinterpret the LK involution using the theory of finite posets.

A (finite) **poset**, or *partially ordered set*, is a (finite) set P together with a relation \leq satisfying the usual axioms of a partial order:

- *transitivity* ($x \leq y, y \leq z \Rightarrow x \leq z$);
- *anti-symmetry* ($x \leq y, y \leq x \Rightarrow x = y$);
- *reflexivity* ($x \leq x$).

We represent posets via their **Hasse diagrams**:




Posets

We will now reinterpret the LK involution using the theory of finite posets.

A (finite) **poset**, or *partially ordered set*, is a (finite) set P together with a relation \leq satisfying the usual axioms of a partial order:

- *transitivity* ($x \leq y, y \leq z \Rightarrow x \leq z$);
- *anti-symmetry* ($x \leq y, y \leq x \Rightarrow x = y$);
- *reflexivity* ($x \leq x$).

We represent posets via their **Hasse diagrams**:



Here an edge from x (below) to y (above) represents the **cover relation** $x \lessdot y$ in P , which means $x \leq y$ and there is no $p \in P$ with $x \leq p \leq y$.

Chains and antichains

Two elements x, y in a poset P are **comparable** if either $x \leq y$ or $y \leq x$.

Chains and antichains

Two elements x, y in a poset P are **comparable** if either $x \leq y$ or $y \leq x$. A **chain** $C \subseteq P$ of P is a subset of pairwise comparable elements (i.e., a chain is a *totally ordered* subset $C = \{x_1 < x_2 < \cdots < x_k\}$).

Chains and antichains

Two elements x, y in a poset P are **comparable** if either $x \leq y$ or $y \leq x$. A **chain** $C \subseteq P$ of P is a subset of pairwise comparable elements (i.e., a chain is a *totally ordered* subset $C = \{x_1 < x_2 < \cdots < x_k\}$). A chain C is **maximal** if it is not strictly contained in another chain.

Chains and antichains

Two elements x, y in a poset P are **comparable** if either $x \leq y$ or $y \leq x$. A **chain** $C \subseteq P$ of P is a subset of pairwise comparable elements (i.e., a chain is a *totally ordered* subset $C = \{x_1 < x_2 < \cdots < x_k\}$). A chain C is **maximal** if it is not strictly contained in another chain.

Two elements $x, y \in P$ are **incomparable** if they are not comparable.

Chains and antichains

Two elements x, y in a poset P are **comparable** if either $x \leq y$ or $y \leq x$. A **chain** $C \subseteq P$ of P is a subset of pairwise comparable elements (i.e., a chain is a *totally ordered* subset $C = \{x_1 < x_2 < \cdots < x_k\}$). A chain C is **maximal** if it is not strictly contained in another chain.

Two elements $x, y \in P$ are **incomparable** if they are not comparable. An **antichain** $A \subseteq P$ of P is a subset of pairwise incomparable elements.

Chains and antichains


Two elements x, y in a poset P are **comparable** if either $x \leq y$ or $y \leq x$. A **chain** $C \subseteq P$ of P is a subset of pairwise comparable elements (i.e., a chain is a *totally ordered* subset $C = \{x_1 < x_2 < \cdots < x_k\}$). A chain C is **maximal** if it is not strictly contained in another chain.

Two elements $x, y \in P$ are **incomparable** if they are not comparable. An **antichain** $A \subseteq P$ of P is a subset of pairwise incomparable elements. We use $\mathcal{A}(P)$ to denote the set of antichains of P .

Chains and antichains

Two elements x, y in a poset P are **comparable** if either $x \leq y$ or $y \leq x$. A **chain** $C \subseteq P$ of P is a subset of pairwise comparable elements (i.e., a chain is a *totally ordered* subset $C = \{x_1 < x_2 < \dots < x_k\}$). A chain C is **maximal** if it is not strictly contained in another chain.

Two elements $x, y \in P$ are **incomparable** if they are not comparable. An **antichain** $A \subseteq P$ of P is a subset of pairwise incomparable elements. We use $\mathcal{A}(P)$ to denote the set of antichains of P .



Here the red elements form a maximal chain C , and the blue elements form an antichain $A \in \mathcal{A}(P)$.

The poset Δ^{n-1}

One particular family of posets Δ^{n-1} is relevant to the LK involution.

The poset Δ^{n-1}

One particular family of posets Δ^{n-1} is relevant to the LK involution.

Δ^{n-1} is the poset whose elements are **intervals** $[i, j] := \{i, i + 1, \dots, j\}$ with $1 \leq i \leq j \leq n - 1$, and with the partial order given by **inclusion**:

$$[i, j] \leq [i', j'] \iff [i, j] \subseteq [i', j'] \iff i \leq i' \leq j' \leq j$$

The poset Δ^{n-1}

One particular family of posets Δ^{n-1} is relevant to the LK involution.

Δ^{n-1} is the poset whose elements are **intervals** $[i, j] := \{i, i + 1, \dots, j\}$ with $1 \leq i \leq j \leq n - 1$, and with the partial order given by **inclusion**:

$$[i, j] \leq [i', j'] \iff [i, j] \subseteq [i', j'] \iff i \leq i' \leq j' \leq j$$

Δ^{n-1} has a “triangular” Hasse diagram:

The poset Δ^{n-1}

One particular family of posets Δ^{n-1} is relevant to the LK involution.

Δ^{n-1} is the poset whose elements are **intervals** $[i, j] := \{i, i + 1, \dots, j\}$ with $1 \leq i \leq j \leq n - 1$, and with the partial order given by **inclusion**:

$$[i, j] \leq [i', j'] \iff [i, j] \subseteq [i', j'] \iff i \leq i' \leq j' \leq j$$

Δ^{n-1} has a “triangular” Hasse diagram:

$[1, 1]$
○

Δ^1


The poset Δ^{n-1}

One particular family of posets Δ^{n-1} is relevant to the LK involution.

Δ^{n-1} is the poset whose elements are **intervals** $[i, j] := \{i, i + 1, \dots, j\}$ with $1 \leq i \leq j \leq n - 1$, and with the partial order given by **inclusion**:

$$[i, j] \leq [i', j'] \iff [i, j] \subseteq [i', j'] \iff i \leq i' \leq j' \leq j$$

Δ^{n-1} has a “triangular” Hasse diagram:




The poset Δ^{n-1}

One particular family of posets Δ^{n-1} is relevant to the LK involution.

Δ^{n-1} is the poset whose elements are **intervals** $[i, j] := \{i, i + 1, \dots, j\}$ with $1 \leq i \leq j \leq n - 1$, and with the partial order given by **inclusion**:

$$[i, j] \leq [i', j'] \iff [i, j] \subseteq [i', j'] \iff i \leq i' \leq j' \leq j$$

Δ^{n-1} has a “triangular” Hasse diagram:




The poset Δ^{n-1}

One particular family of posets Δ^{n-1} is relevant to the LK involution.

Δ^{n-1} is the poset whose elements are **intervals** $[i, j] := \{i, i + 1, \dots, j\}$ with $1 \leq i \leq j \leq n - 1$, and with the partial order given by **inclusion**:

$$[i, j] \leq [i', j'] \iff [i, j] \subseteq [i', j'] \iff i \leq i' \leq j' \leq j$$

Δ^{n-1} has a “triangular” Hasse diagram:




Dyck paths are antichains in Δ^{n-1}

There is a natural, pictoral bijection between the Dyck paths of length $2n$ and the antichains of Δ^{n-1} :


Dyck paths are antichains in Δ^{n-1}

There is a natural, pictorial bijection between the Dyck paths of length $2n$ and the antichains of Δ^{n-1} :



Dyck paths are antichains in Δ^{n-1}


There is a natural, pictorial bijection between the Dyck paths of length $2n$ and the antichains of Δ^{n-1} :



Observe how, under this bijection, the number of valleys of a Dyck path Γ becomes the number of elements of an antichain A .

Dyck paths are antichains in Δ^{n-1}

There is a natural, pictorial bijection between the Dyck paths of length $2n$ and the antichains of Δ^{n-1} :



Observe how, under this bijection, the number of valleys of a Dyck path Γ becomes the number of elements of an antichain A .

Via this bijection, we can view the LK involution as an involution on antichains $\text{LK}: \mathcal{A}(\Delta^{n-1}) \rightarrow \mathcal{A}(\Delta^{n-1})$ which satisfies

$$\#A + \#\text{LK}(A) = n - 1.$$

The LK involution on antichains

D. Panyushev gave a simple description of the LK involution on $\mathcal{A}(\Delta^{n-1})$:

The LK involution on antichains

D. Panyushev gave a simple description of the LK involution on $\mathcal{A}(\Delta^{n-1})$:

Theorem (Panyushev, 2004)

Let $A = \{[i_1, j_1], [i_2, j_2], \dots, [i_k, j_k]\} \in \mathcal{A}(\Delta^{n-1})$ with $i_1 < i_2 < \dots < i_k$.

Then $\text{LK}(A) = \{[i'_1, j'_1], [i'_2, j'_2], \dots, [i'_{n-1-k}, j'_{n-1-k}]\} \in \mathcal{A}(\Delta^{n-1})$, where

- $\{i'_1 < i'_2 < \dots < i'_{n-1-k}\} = \{1, 2, \dots, n-1\} \setminus \{j_1, j_2, \dots, j_k\}$;
- $\{j'_1 < j'_2 < \dots < j'_{n-1-k}\} = \{1, 2, \dots, n-1\} \setminus \{i_1, i_2, \dots, i_k\}$.

The LK involution on antichains

D. Panyushev gave a simple description of the LK involution on $\mathcal{A}(\Delta^{n-1})$:


Theorem (Panyushev, 2004)

Let $A = \{[i_1, j_1], [i_2, j_2], \dots, [i_k, j_k]\} \in \mathcal{A}(\Delta^{n-1})$ with $i_1 < i_2 < \dots < i_k$.
 Then $\text{LK}(A) = \{[i'_1, j'_1], [i'_2, j'_2], \dots, [i'_{n-1-k}, j'_{n-1-k}]\} \in \mathcal{A}(\Delta^{n-1})$, where

- $\{i'_1 < i'_2 < \dots < i'_{n-1-k}\} = \{1, 2, \dots, n-1\} \setminus \{j_1, j_2, \dots, j_k\}$;
- $\{j'_1 < j'_2 < \dots < j'_{n-1-k}\} = \{1, 2, \dots, n-1\} \setminus \{i_1, i_2, \dots, i_k\}$.


From Panyushev's description, it is immediate that this operation is an involution (i.e., $\text{LK}^2(A) = A$), and that $\#A + \#\text{LK}(A) = n - 1$.

The LK involution on antichains: example




$$A = \{[1, \underline{2}], [4, \underline{4}], [5, \underline{6}], [6, \underline{9}]\}$$

The LK involution on antichains: example



$$A = \{[1, \underline{2}], [4, \underline{4}], [5, \underline{6}], [6, \underline{9}]\}$$



$$\text{LK}(A) = \{[1, \underline{2}], [\underline{3}, \underline{3}], [\underline{5}, \underline{7}], [\underline{7}, \underline{8}], [\underline{8}, \underline{9}]\}$$

$$\begin{aligned} \{1, 3, 5, 7, 8\} &= \{1, \dots, 9\} \setminus \{\underline{2}, \underline{4}, \underline{6}, \underline{9}\} \\ \{\underline{2}, \underline{3}, \underline{7}, \underline{8}, \underline{9}\} &= \{1, \dots, 9\} \setminus \{1, 4, 5, 6\} \end{aligned}$$

Section 3

Toggling

Toggling for antichains

Our first new result gives another expression for the LK involution in terms of certain “local” involutions called **toggles**.

Toggling for antichains

Our first new result gives another expression for the LK involution in terms of certain “local” involutions called **toggles**.

Let P be a poset and $A \in \mathcal{A}(P)$ an antichain. Let $p \in P$ be any element. The **toggle of p in A** is the antichain $\tau_p(A) \in \mathcal{A}(P)$, where

$$\tau_p(A) := \begin{cases} A \setminus \{p\} & \text{if } p \in A; \\ A \cup \{p\} & \text{if } p \notin A \text{ and } A \cup \{p\} \text{ remains an antichain;} \\ A & \text{otherwise.} \end{cases}$$


Toggling for antichains

Our first new result gives another expression for the LK involution in terms of certain “local” involutions called **toggles**.

Let P be a poset and $A \in \mathcal{A}(P)$ an antichain. Let $p \in P$ be any element. The **toggle of p in A** is the antichain $\tau_p(A) \in \mathcal{A}(P)$, where

$$\tau_p(A) := \begin{cases} A \setminus \{p\} & \text{if } p \in A; \\ A \cup \{p\} & \text{if } p \notin A \text{ and } A \cup \{p\} \text{ remains an antichain;} \\ A & \text{otherwise.} \end{cases}$$

In other words, we “toggle” the status of p in A , if possible:




Toggling for antichains

Our first new result gives another expression for the LK involution in terms of certain “local” involutions called **toggles**.

Let P be a poset and $A \in \mathcal{A}(P)$ an antichain. Let $p \in P$ be any element. The **toggle of p in A** is the antichain $\tau_p(A) \in \mathcal{A}(P)$, where

$$\tau_p(A) := \begin{cases} A \setminus \{p\} & \text{if } p \in A; \\ A \cup \{p\} & \text{if } p \notin A \text{ and } A \cup \{p\} \text{ remains an antichain;} \\ A & \text{otherwise.} \end{cases}$$

In other words, we “toggle” the status of p in A , if possible:




Toggling for antichains

Our first new result gives another expression for the LK involution in terms of certain “local” involutions called **toggles**.

Let P be a poset and $A \in \mathcal{A}(P)$ an antichain. Let $p \in P$ be any element. The **toggle of p in A** is the antichain $\tau_p(A) \in \mathcal{A}(P)$, where

$$\tau_p(A) := \begin{cases} A \setminus \{p\} & \text{if } p \in A; \\ A \cup \{p\} & \text{if } p \notin A \text{ and } A \cup \{p\} \text{ remains an antichain;} \\ A & \text{otherwise.} \end{cases}$$

In other words, we “toggle” the status of p in A , if possible:



$$\begin{aligned} \tau_x(\bullet \diagup \circ \diagdown \circ) &= \circ \diagup \circ \diagdown \circ \\ \tau_x(\circ \diagup \bullet \diagdown \circ) &= \bullet \diagup \circ \diagdown \bullet \end{aligned}$$


Toggling for antichains

Our first new result gives another expression for the LK involution in terms of certain “local” involutions called **toggles**.

Let P be a poset and $A \in \mathcal{A}(P)$ an antichain. Let $p \in P$ be any element. The **toggle of p in A** is the antichain $\tau_p(A) \in \mathcal{A}(P)$, where

$$\tau_p(A) := \begin{cases} A \setminus \{p\} & \text{if } p \in A; \\ A \cup \{p\} & \text{if } p \notin A \text{ and } A \cup \{p\} \text{ remains an antichain;} \\ A & \text{otherwise.} \end{cases}$$


In other words, we “toggle” the status of p in A , if possible:



$$\begin{aligned} \tau_x(\bullet \diagup \circ \diagdown \circ) &= \circ \diagup \circ \diagdown \circ \\ \tau_x(\circ \diagup \bullet \diagdown \bullet) &= \bullet \diagup \circ \diagdown \bullet \\ \tau_x(\circ \diagup \bullet \diagdown \circ) &= \circ \diagup \bullet \diagdown \circ \end{aligned}$$


Toggling in ranked posets

A poset P is **ranked** if we can write $P = P_1 \sqcup P_2 \sqcup \cdots \sqcup P_r$ so that all the edges of the Hasse diagram of P are from P_i (below) to P_{i+1} (above):



Toggling in ranked posets

A poset P is **ranked** if we can write $P = P_1 \sqcup P_2 \sqcup \cdots \sqcup P_r$ so that all the edges of the Hasse diagram of P are from P_i (below) to P_{i+1} (above):




Since τ_p and τ_q commute if p and q are incomparable, and all the elements within a rank are incomparable, we can define

$$\tau_i := \prod_{p \in P_i} \tau_p$$

to be the composition of all toggles at rank i , for $i = 1, \dots, r$:

Toggling in ranked posets

A poset P is **ranked** if we can write $P = P_1 \sqcup P_2 \sqcup \cdots \sqcup P_r$ so that all the edges of the Hasse diagram of P are from P_i (below) to P_{i+1} (above):



Since τ_p and τ_q commute if p and q are incomparable, and all the elements within a rank are incomparable, we can define

$$\tau_i := \prod_{p \in P_i} \tau_p$$

to be the composition of all toggles at rank i , for $i = 1, \dots, r$:



The LK involution as a composition of toggles

Theorem (H.-Joseph, 2021)

The LK involution $\text{LK}: \mathcal{A}(\Delta^{n-1}) \rightarrow \mathcal{A}(\Delta^{n-1})$ can be written as the following composition of toggles:

$$\text{LK} = (\tau_{n-1})(\tau_{n-1}\tau_{n-2}) \cdots (\tau_{n-1} \cdots \tau_3\tau_2)(\tau_{n-1} \cdots \tau_2\tau_1)$$

The LK involution as a composition of toggles


Theorem (H.-Joseph, 2021)

The LK involution $\text{LK}: \mathcal{A}(\Delta^{n-1}) \rightarrow \mathcal{A}(\Delta^{n-1})$ can be written as the following composition of toggles:



$$\text{LK} = (\tau_{n-1})(\tau_{n-1}\tau_{n-2}) \cdots (\tau_{n-1} \cdots \tau_3\tau_2)(\tau_{n-1} \cdots \tau_2\tau_1)$$

Remark: for a ranked poset P , the composition of toggles $\tau_r \cdots \tau_2\tau_1$ “from bottom to top” is called **rowmotion** and has been studied by many authors (Cameron–Fon-Der-Flaass, Striker–Williams, Propp–Roby, Joseph, etc...) in the emerging subfield of **dynamical algebraic combinatorics**.



The LK involution as a composition of toggles: example





The LK involution as a composition of toggles: example





The LK involution as a composition of toggles: example





The LK involution as a composition of toggles: example





The LK involution as a composition of toggles: example





The LK involution as a composition of toggles: example



The LK involution as a composition of toggles: example



The LK involution as a composition of toggles: example



Section 4

Piecewise linear and birational lifts

Convex polytopes

Why did we want to write the LK involution as a composition of toggles?
In order to **extend** it to the **piecewise linear** realm...

Convex polytopes

Why did we want to write the LK involution as a composition of toggles?
In order to **extend** it to the **piecewise linear** realm...

A **convex polytope** in \mathbb{R}^n can be defined either as

- a convex hull of finitely many points (**vertices**);
- a bounded intersection of finitely many linear inequalities (**facets**).


Convex polytopes

Why did we want to write the LK involution as a composition of toggles?
In order to **extend** it to the **piecewise linear** realm...

A **convex polytope** in \mathbb{R}^n can be defined either as

- a convex hull of finitely many points (**vertices**);
- a bounded intersection of finitely many linear inequalities (**facets**).

In dimensions 2 and 3, these are familiar shapes:




Convex polytopes

Why did we want to write the LK involution as a composition of toggles?
In order to **extend** it to the **piecewise linear** realm...

A **convex polytope** in \mathbb{R}^n can be defined either as

- a convex hull of finitely many points (**vertices**);
- a bounded intersection of finitely many linear inequalities (**facets**).

In dimensions 2 and 3, these are familiar shapes:



There is a rich interplay between combinatorics and convex geometry,
because combinatorial objects can often be “realized” polytopally: e.g.,
the subsets of $\{1, 2, \dots, n\}$ correspond to the vertices of the n -hypercube.

The chain polytope of a poset

In 1986, Richard Stanley associated to any poset P two polytopes in \mathbb{R}^P , the **order polytope** $\mathcal{O}(P)$ and the **chain polytope** $\mathcal{C}(P)$.

The chain polytope of a poset

In 1986, Richard Stanley associated to any poset P two polytopes in \mathbb{R}^P , the **order polytope** $\mathcal{O}(P)$ and the **chain polytope** $\mathcal{C}(P)$.

The **chain polytope** $\mathcal{C}(P)$ has facets

$$0 \leq x_p, \quad \forall p \in P$$

$$\sum_{p \in C} x_p \leq 1, \quad \forall C \subseteq P \text{ a maximal chain.}$$

The chain polytope of a poset

In 1986, Richard Stanley associated to any poset P two polytopes in \mathbb{R}^P , the **order polytope** $\mathcal{O}(P)$ and the **chain polytope** $\mathcal{C}(P)$.

The **chain polytope** $\mathcal{C}(P)$ has facets

$$0 \leq x_p, \quad \forall p \in P$$

$$\sum_{p \in C} x_p \leq 1, \quad \forall C \subseteq P \text{ a maximal chain.}$$

Stanley proved that the **vertices** of $\mathcal{C}(P)$ are precisely the **indicator functions of antichains** $A \in \mathcal{A}(P)$:

The chain polytope of a poset


In 1986, Richard Stanley associated to any poset P two polytopes in \mathbb{R}^P , the **order polytope** $\mathcal{O}(P)$ and the **chain polytope** $\mathcal{C}(P)$.

The **chain polytope** $\mathcal{C}(P)$ has facets

$$0 \leq x_p, \quad \forall p \in P$$

$$\sum_{p \in C} x_p \leq 1, \quad \forall C \subseteq P \text{ a maximal chain.}$$

Stanley proved that the **vertices** of $\mathcal{C}(P)$ are precisely the **indicator functions of antichains** $A \in \mathcal{A}(P)$:



The chain polytope of a poset




In 1986, Richard Stanley associated to any poset P two polytopes in \mathbb{R}^P , the **order polytope** $\mathcal{O}(P)$ and the **chain polytope** $\mathcal{C}(P)$.

The **chain polytope** $\mathcal{C}(P)$ has facets

$$0 \leq x_p, \quad \forall p \in P$$

$$\sum_{p \in C} x_p \leq 1, \quad \forall C \subseteq P \text{ a maximal chain.}$$

Stanley proved that the **vertices** of $\mathcal{C}(P)$ are precisely the **indicator functions of antichains** $A \in \mathcal{A}(P)$:



Piecewise linear toggling

In 2013, D. Einstein and J. Propp (c.f. Joseph) introduced a (continuous) **piecewise linear extension** of the toggles τ_p .

Piecewise linear toggling

In 2013, D. Einstein and J. Propp (c.f. Joseph) introduced a (continuous) **piecewise linear extension** of the toggles τ_p .

For $p \in P$, the **PL toggle** $\tau_p^{\text{PL}}: \mathcal{C}(P) \rightarrow \mathcal{C}(P)$ is defined by

$$\tau_p^{\text{PL}}(\pi)(q) := \begin{cases} \pi(q) & \text{if } q \neq p; \\ 1 - \max \left\{ \sum_{r \in C} \pi(r) : \begin{array}{l} C \subseteq P \text{ a maximal} \\ \text{chain with } p \in C \end{array} \right\} & \text{if } p = q. \end{cases}$$

Piecewise linear toggling

In 2013, D. Einstein and J. Propp (c.f. Joseph) introduced a (continuous) **piecewise linear extension** of the toggles τ_p .

For $p \in P$, the **PL toggle** $\tau_p^{\text{PL}}: \mathcal{C}(P) \rightarrow \mathcal{C}(P)$ is defined by

$$\tau_p^{\text{PL}}(\pi)(q) := \begin{cases} \pi(q) & \text{if } q \neq p; \\ 1 - \max \left\{ \sum_{r \in C} \pi(r) : \begin{array}{l} C \subseteq P \text{ a maximal} \\ \text{chain with } p \in C \end{array} \right\} & \text{if } p = q. \end{cases}$$

Restricted to the vertices of the chain polytope $\mathcal{C}(P)$, it is the same as τ_p .

Piecewise linear toggling


In 2013, D. Einstein and J. Propp (c.f. Joseph) introduced a (continuous) **piecewise linear extension** of the toggles τ_p .

For $p \in P$, the **PL toggle** $\tau_p^{\text{PL}}: \mathcal{C}(P) \rightarrow \mathcal{C}(P)$ is defined by

$$\tau_p^{\text{PL}}(\pi)(q) := \begin{cases} \pi(q) & \text{if } q \neq p; \\ 1 - \max \left\{ \sum_{r \in C} \pi(r) : \begin{array}{l} C \subseteq P \text{ a maximal} \\ \text{chain with } p \in C \end{array} \right\} & \text{if } p = q. \end{cases}$$

Restricted to the vertices of the chain polytope $\mathcal{C}(P)$, it is the same as τ_p .

Geometrically, τ_p **reflects** π within line segment in $\mathcal{C}(P)$ in direction x_p :



The PL LK involution

As before, for a ranked poset P we use $\tau_i^{\text{PL}} := \prod_{p \in P_i} \tau_p^{\text{PL}}$ to denote the composition of all toggles at rank i .

The PL LK involution

As before, for a ranked poset P we use $\tau_i^{\text{PL}} := \prod_{p \in P_i} \tau_p^{\text{PL}}$ to denote the composition of all toggles at rank i .

We define the **PL LK involution** $\text{LK}^{\text{PL}} : \mathcal{C}(\Delta^{n-1}) \rightarrow \mathcal{C}(\Delta^{n-1})$ to be

$$\text{LK}^{\text{PL}} := (\tau_{n-1}^{\text{PL}})(\tau_{n-1}^{\text{PL}} \tau_{n-2}^{\text{PL}}) \cdots (\tau_{n-1}^{\text{PL}} \cdots \tau_3^{\text{PL}} \tau_2^{\text{PL}})(\tau_{n-1}^{\text{PL}} \cdots \tau_2^{\text{PL}} \tau_1^{\text{PL}})$$

By prior theorem, it's same as LK when restricted to the vertices of $\mathcal{C}(P)$.

The PL LK involution

As before, for a ranked poset P we use $\tau_i^{\text{PL}} := \prod_{p \in P_i} \tau_p^{\text{PL}}$ to denote the composition of all toggles at rank i .

We define the **PL LK involution** $\text{LK}^{\text{PL}} : \mathcal{C}(\Delta^{n-1}) \rightarrow \mathcal{C}(\Delta^{n-1})$ to be

$$\text{LK}^{\text{PL}} := (\tau_{n-1}^{\text{PL}})(\tau_{n-1}^{\text{PL}} \tau_{n-2}^{\text{PL}}) \cdots (\tau_{n-1}^{\text{PL}} \cdots \tau_3^{\text{PL}} \tau_2^{\text{PL}})(\tau_{n-1}^{\text{PL}} \cdots \tau_2^{\text{PL}} \tau_1^{\text{PL}})$$

By prior theorem, it's same as LK when restricted to the vertices of $\mathcal{C}(P)$.

Theorem (H.-Joseph, 2021)

- (1) LK^{PL} is an involution.
- (2) For any $\pi \in \mathcal{C}(\Delta^{n-1})$, $\sum_{p \in P} \pi(p) + \sum_{p \in P} \text{LK}^{\text{PL}}(\pi)(p) = n - 1$.

The PL LK involution

As before, for a ranked poset P we use $\tau_i^{\text{PL}} := \prod_{p \in P_i} \tau_p^{\text{PL}}$ to denote the composition of all toggles at rank i .

We define the **PL LK involution** $\text{LK}^{\text{PL}} : \mathcal{C}(\Delta^{n-1}) \rightarrow \mathcal{C}(\Delta^{n-1})$ to be

$$\text{LK}^{\text{PL}} := (\tau_{n-1}^{\text{PL}})(\tau_{n-1}^{\text{PL}} \tau_{n-2}^{\text{PL}}) \cdots (\tau_{n-1}^{\text{PL}} \cdots \tau_3^{\text{PL}} \tau_2^{\text{PL}})(\tau_{n-1}^{\text{PL}} \cdots \tau_2^{\text{PL}} \tau_1^{\text{PL}})$$


By prior theorem, it's same as LK when restricted to the vertices of $\mathcal{C}(P)$.

Theorem (H.-Joseph, 2021)


- (1) LK^{PL} is an involution.
- (2) For any $\pi \in \mathcal{C}(\Delta^{n-1})$, $\sum_{p \in P} \pi(p) + \sum_{p \in P} \text{LK}^{\text{PL}}(\pi)(p) = n - 1$.

Observe that (2) is an extension of the fact that LK combinatorially exhibits the symmetry of the Narayana numbers.

The PL LK involution: example



The PL LK involution: example



We can check that

$$(0.5 + 0 + 0 + 0.1 + 0 + 0.3 + 0.1 + 0.1 + 0.2 + 0.4) + (0 + 0.1 + 0.1 + 0.2 + 0.4 + 0.7 + 0.5 + 0.1 + 0.2 + 0) =$$

$$1.7 + 2.3 = 4$$

Tropical geometry

Algebraic geometry studies
polynomial expressions like


$$x^3y + y^3z + z^3x$$

Tropical geometry

Algebraic geometry studies
polynomial expressions like

$$x^3y + y^3z + z^3x$$

which lead to “curvy” hypersurfaces
like



Tropical geometry


Algebraic geometry studies
polynomial expressions like

$$x^3y + y^3z + z^3x$$

Tropical geometry studies
piecewise linear expressions like

$$\max(3x + y, 3y + z, 3z + x)$$

which lead to “curvy” hypersurfaces
like




Tropical geometry

Algebraic geometry studies
polynomial expressions like

$$x^3y + y^3z + z^3x$$


which lead to “curvy” hypersurfaces
like



Tropical geometry studies
piecewise linear expressions like

$$\max(3x + y, 3y + z, 3z + x)$$

which lead to “flat” polytopal
complexes like



“De-tropicalization”

The process of replacing $(\times, +)$ with $(+, \max)$ in a polynomial expression is called **tropicalization**:

$$x^3y + y^3z + z^3x \mapsto \max(3x + y, 3y + z, 3z + x)$$

It lead to important interactions between algebraic & convex geometry.

“De-tropicalization”

The process of replacing $(\times, +)$ with $(+, \max)$ in a polynomial expression is called **tropicalization**:

$$x^3y + y^3z + z^3x \mapsto \max(3x + y, 3y + z, 3z + x)$$

It lead to important interactions between algebraic & convex geometry.

(Adjective “tropical” comes from fact that computer scientist & pioneer of tropical geometry Imre Simon worked at University of São Paulo,  Sam Hopkins (2021)

“De-tropicalization”

The process of replacing $(\times, +)$ with $(+, \max)$ in a polynomial expression is called **tropicalization**:

$$x^3y + y^3z + z^3x \mapsto \max(3x + y, 3y + z, 3z + x)$$

It lead to important interactions between algebraic & convex geometry.

(Adjective “tropical” comes from fact that computer scientist & pioneer of tropical geometry Imre Simon worked at University of São Paulo,  

The process of replacing $(+, \max)$ with $(\times, +)$ in a piecewise linear expression is called **de-tropicalization***:

$$\max(3x + y, 3y + z, 3z + x) \mapsto x^3y + y^3z + z^3x$$

It is often interesting to try to de-tropicalize PL maps, like those coming from classical combinatorial constructions.

Birational toggling

Einstein–Propp (c.f. Joseph–Roby) also introduced a **birational extension** of the toggles τ_p , via de-tropicalization.

Birational toggling

Einstein–Propp (c.f. Joseph–Roby) also introduced a **birational extension** of the toggles τ_p , via de-tropicalization.

For $p \in P$, the **birational toggle** $\tau_p^B: \mathbb{C}^P \dashrightarrow \mathbb{C}^P$ is

$$\tau_p^B(\pi)(q) := \begin{cases} \pi(q) & \text{if } q \neq p; \\ \kappa \cdot \left(\prod_{\substack{C \subseteq P \\ \text{max. chain,} \\ p \in C}} \sum_{r \in C} \pi(r) \right)^{-1} & \text{if } p = q, \end{cases}$$

where $\kappa \in \mathbb{C}$ is some fixed constant.

Birational toggling

Einstein–Propp (c.f. Joseph–Roby) also introduced a **birational extension** of the toggles τ_p , via de-tropicalization.

For $p \in P$, the **birational toggle** $\tau_p^B : \mathbb{C}^P \dashrightarrow \mathbb{C}^P$ is

$$\tau_p^B(\pi)(q) := \begin{cases} \pi(q) & \text{if } q \neq p; \\ \kappa \cdot \left(\prod_{\substack{C \subseteq P \\ \text{max. chain,} \\ p \in C}} \sum_{r \in C} \pi(r) \right)^{-1} & \text{if } p = q, \end{cases}$$

where $\kappa \in \mathbb{C}$ is some fixed constant.

The birational toggle τ_p^B tropicalizes to the PL toggle τ_p^{PL} .

The birational LK involution

As before, if P is ranked we set $\tau_i^B := \prod_{p \in P_i} \tau_p^B$.

The birational LK involution

As before, if P is ranked we set $\tau_i^B := \prod_{p \in P_i} \tau_p^B$.

We define the birational LK involution $\text{LK}^B : \mathbb{C}^{\Delta^{n-1}} \dashrightarrow \mathbb{C}^{\Delta^{n-1}}$ by

$$\text{LK}^B := (\tau_{n-1}^B)(\tau_{n-1}^B \tau_{n-2}^B) \cdots (\tau_{n-1}^B \cdots \tau_3^B \tau_2^B)(\tau_{n-1}^B \cdots \tau_2^B \tau_1^B)$$

It tropicalizes to LK^{PL} .

The birational LK involution

As before, if P is ranked we set $\tau_i^B := \prod_{p \in P_i} \tau_p^B$.

We define the birational LK involution $\text{LK}^B : \mathbb{C}^{\Delta^{n-1}} \dashrightarrow \mathbb{C}^{\Delta^{n-1}}$ by

$$\text{LK}^B := (\tau_{n-1}^B)(\tau_{n-1}^B \tau_{n-2}^B) \cdots (\tau_{n-1}^B \cdots \tau_3^B \tau_2^B)(\tau_{n-1}^B \cdots \tau_2^B \tau_1^B)$$

It tropicalizes to LK^{PL} .

Theorem (H.-Joseph, 2021)

- (1) LK^B is an involution.
- (2) For any $\pi \in \mathbb{C}^{\Delta^{n-1}}$, $\prod_{p \in P} \pi(p) \cdot \prod_{p \in P} \text{LK}^B(\pi)(p) = \kappa^{n-1}$.

The birational LK involution

As before, if P is ranked we set $\tau_i^B := \prod_{p \in P_i} \tau_p^B$.

We define the birational LK involution $\text{LK}^B : \mathbb{C}^{\Delta^{n-1}} \dashrightarrow \mathbb{C}^{\Delta^{n-1}}$ by

$$\text{LK}^B := (\tau_{n-1}^B)(\tau_{n-1}^B \tau_{n-2}^B) \cdots (\tau_{n-1}^B \cdots \tau_3^B \tau_2^B)(\tau_{n-1}^B \cdots \tau_2^B \tau_1^B)$$


It tropicalizes to LK^{PL} .

Theorem (H.-Joseph, 2021)


- (1) LK^B is an involution.
- (2) For any $\pi \in \mathbb{C}^{\Delta^{n-1}}$, $\prod_{p \in P} \pi(p) \cdot \prod_{p \in P} \text{LK}^B(\pi)(p) = \kappa^{n-1}$.

(2) is the birational analog of the fact that LK combinatorially exhibits the symmetry of the Narayana numbers.

The birational LK involution: example




The birational LK involution: example



We can check that this operation really is an involution; e.g.,

$$\frac{z'(x' + y')}{y'} = \frac{\frac{xy}{x+y} \cdot \left(\frac{z(x+y)}{y} + \frac{z(x+y)}{x} \right)}{\frac{z(x+y)}{x}} = \frac{zx + zy}{\frac{z(x+y)}{x}} = \frac{z(x+y)}{\frac{z(x+y)}{x}} = x.$$

The birational LK involution: example



We can check that this operation really is an involution; e.g.,

$$\frac{z'(x' + y')}{y'} = \frac{\frac{xy}{x+y} \cdot \left(\frac{z(x+y)}{y} + \frac{z(x+y)}{x} \right)}{\frac{z(x+y)}{x}} = \frac{zx + zy}{\frac{z(x+y)}{x}} = \frac{z(x+y)}{\frac{z(x+y)}{x}} = x.$$

And if we multiply together all the above values, we get κ^3 .

Thank you!

these slides are available on my website
and the paper on the arXiv: arXiv:2012.15795

Exercises

231

- 6.24.** [?] Explain the significance of the following sequence:

un, dos, tres, quatre, cinc, sis, set, vuit, nou, deu, . . .

R. Stanley, *Enumerative Combinatorics*, Vol. 2