

Howard Math 274, HW# 2,

Spring 2022; Instructor: Sam Hopkins; Due: Friday, March 25th

1. Let $\lambda = (\lambda_1, \lambda_2, \dots), \mu = (\mu_1, \mu_2, \dots) \vdash n$ be partitions of n . Recall that the *lexicographic order* \prec on partitions of n is given by $\mu \prec \lambda$ iff there is some j such that $\mu_i = \lambda_i$ for all $i < j$ and $\mu_j < \lambda_j$. It is a total order: we either have $\mu \prec \lambda$ or $\lambda \prec \mu$ or $\lambda = \mu$.

A different order on partitions of n is the dominance order. The *dominance order* \leq is defined by $\mu \leq \lambda$ iff $\mu_1 + \mu_2 + \dots + \mu_j \leq \lambda_1 + \lambda_2 + \dots + \lambda_j$ for all j . The dominance order is only partial order: we might have neither $\mu \leq \lambda$ nor $\lambda \leq \mu$.

- (a) Show that the lexicographic order *extends* the dominance order in the sense that if we have partitions $\mu, \nu \vdash n$ with $\mu \leq \lambda$ and $\mu \neq \lambda$ then necessarily $\mu \prec \lambda$.
 - (b) Give an example of partitions $\mu, \nu \vdash n$ with $\mu \prec \lambda$ but $\mu \not\leq \lambda$.
2. Show that we could've used dominance order instead of lexicographic order in our arguments about the triangularity of the transition matrices from p_λ or e_λ to m_μ . That is, show that

$$p_\lambda = \sum_{\lambda \leq \mu} \alpha_\mu^\lambda m_\mu \quad \text{and} \quad e_\lambda = \sum_{\mu \leq \lambda^t} \beta_\mu^\lambda m_\mu \quad \text{for coefficients } \alpha_\mu^\lambda, \beta_\mu^\lambda \in \mathbb{C}$$

for any $\lambda \vdash n$, where \leq is dominance order and λ^t is the transpose (a.k.a. conjugate) of λ .

3. Let $\lambda \vdash n$ and define f^λ to be the coefficient of $x_1 x_2 \dots x_n$ in the Schur function $s_\lambda(x_1, x_2, \dots)$. Explain why $f^\lambda = f^{\lambda^t}$. Give an example showing that this is not true for other coefficients of Schur functions (i.e., that $s_\lambda \neq s_{\lambda^t}$ in general).
4. The Cauchy–Binet formula says that if $A = (A_{i,j})$ is an $m \times n$ matrix and $B = (B_{i,j})$ is an $n \times m$ matrix, then the determinant of the $m \times m$ matrix AB can be computed by

$$\det(AB) = \sum_{I \subseteq [n], \#I=m} \det(A|_{\text{cols}=I}) \det(B|_{\text{rows}=I}).$$

Here, as always, $[n] := \{1, 2, \dots, n\}$, and $A|_{\text{cols}=I}$ (resp., $B|_{\text{rows}=I}$) means the $m \times m$ matrix we get by restricting A to the columns in I (resp., by restricting B to the rows in I).

Deduce the Cauchy–Binet formula from the Lindström–Gessel–Viennot formula.

Hint: Consider the network with source vertices s_1, \dots, s_m , target vertices t_1, \dots, t_m , and internal vertices k_1, \dots, k_n , and edges $s_i \rightarrow k_j$ with weight $A_{i,j}$ and $k_i \rightarrow t_j$ with weight $B_{i,j}$.

5. Let $\lambda = (\lambda_1, \lambda_2, \dots)$ be a partition and k a positive integer. Give a formula for $m_\lambda(\overbrace{1, 1, \dots, 1}^{k \text{ 1's}})$.

Hint: Your formula can use the *length* $\ell(\lambda) := \max\{i: \lambda_i > 0\}$ of the partition λ , as well as the *multiplicities* $m_i(\lambda) := \{j: \lambda_j = i\}$ for $i \geq 1$.