

# UPHO LATTICES I: EXAMPLES AND NON-EXAMPLES OF CORES

SAM HOPKINS

*Dedicated to Richard Stanley on the occasion of his 80th birthday*

**ABSTRACT.** A poset is called upper homogeneous, or “upho,” if every principal order filter of the poset is isomorphic to the whole poset. We study (finite type  $\mathbb{N}$ -graded) upho lattices, with an eye towards their classification. Any upho lattice has associated to it a finite graded lattice called its core, which determines its rank generating function. We investigate which finite graded lattices arise as cores of upho lattices, providing both positive and negative results. On the one hand, we show that many well-studied finite lattices do arise as cores, and we present combinatorial and algebraic constructions of the upho lattices into which they embed. On the other hand, we show there are obstructions which prevent many finite lattices from being cores.

## 1. INTRODUCTION

Symmetry is a fundamental theme in mathematics. A close cousin of symmetry is self-similarity, where a part resembles the whole. In this paper, we study certain partially ordered sets that are self-similar in a precise sense. Namely, a poset is called *upper homogeneous*, or “*upho*,” if every principal order filter of the poset is isomorphic to the whole poset. In other words, a poset  $\mathcal{P}$  is upho if, looking up from each element  $p \in \mathcal{P}$ , we see another copy of  $\mathcal{P}$ . Upho posets were introduced recently by Stanley [19, 21]. We believe they are a natural and rich class of posets which deserve further attention.

Upho posets are infinite. In order to be able to apply the tools of enumerative and algebraic combinatorics, we need to impose some finiteness condition on the posets we consider. Thus, we restrict our attention to finite type  $\mathbb{N}$ -graded posets. These are the infinite posets  $\mathcal{P}$  that possess a rank function  $\rho: \mathcal{P} \rightarrow \mathbb{N}$  for which we can form the rank generating function

$$F(\mathcal{P}; x) := \sum_{p \in \mathcal{P}} x^{\rho(p)}.$$

Henceforth, upho posets are assumed finite type  $\mathbb{N}$ -graded unless otherwise specified.

The big problem concerning upho posets that we are interested in is the following.

**Problem 1.1.** Classify upho lattices.

Problem 1.1 is likely a hard problem, perhaps even impossible. But let us explain why there is some hope of making progress on this problem. It was shown by Gao et al. [8] that there are uncountably many different rank generating functions of

(finite type  $\mathbb{N}$ -graded) upho posets. This prevents us from being able to say much about upho posets in general. However, the situation is different for *lattices*: in [10] we showed that the rank generating function of a upho lattice is the inverse of a polynomial with integer coefficients.

More precisely, we made the following observation about the rank generating function of an upho lattice. Let  $\mathcal{L}$  be an upho lattice, and let  $L := [\hat{0}, s_1 \vee \cdots \vee s_r] \subseteq \mathcal{L}$  denote the interval in  $\mathcal{L}$  from its minimum  $\hat{0}$  to the join of its atoms  $s_1, \dots, s_r$ . We refer to the finite graded lattice  $L$  as the *core* of the upho lattice  $\mathcal{L}$ . We showed in [10] that

$$(1.1) \quad F(\mathcal{L}; x) = \chi^*(L; x)^{-1},$$

where  $\chi^*(L; x) = \sum_{p \in L} \mu(\hat{0}, p) x^{\rho(p)}$  is the (reciprocal) characteristic polynomial of  $L$ . In this way, the core of an upho lattice determines its rank generating function.<sup>1</sup>

The core does not determine the upho lattice completely. That is, there are different upho lattices with the same core. Nevertheless, to resolve Problem 1.1 we would certainly need to answer the following question.

**Question 1.2.** Which finite graded lattices are cores of upho lattices?

Question 1.2 can be thought of as a kind of tiling problem: our goal is to tile an infinite, fractal lattice  $\mathcal{L}$  using copies of some fixed finite lattice  $L$ , or show that no such tiling is possible. Question 1.2 is the main problem we pursue in this paper. In addressing Question 1.2 here, we provide both positive and negative results.

On the positive side, we show that many well-studied families of finite graded lattices are cores of upho lattices. Our first major result is the following.

**Theorem 1.3.** *Any member of a uniform sequence of supersolvable geometric lattices is the core of some upho lattice.*

Supersolvable lattices were introduced by Stanley in [16], and uniform sequences of lattices were introduced by Dowling in [6]. These two notions represent two different ways that a finite lattice can have a recursive structure. Examples of uniform sequences of supersolvable geometric lattices include:

- the finite Boolean lattices  $B_n$ , i.e., the lattices of subsets of  $\{1, 2, \dots, n\}$ ;
- the  $q$ -analogues  $B_n(q)$  of  $B_n$ , i.e., the lattices of  $\mathbb{F}_q$ -subspaces of  $\mathbb{F}_q^n$ ;
- the partition lattices  $\Pi_n$ , i.e., the lattices of set partitions of  $\{1, 2, \dots, n\}$ ;
- the Type B partition lattices  $\Pi_n^B$ , i.e., the intersection lattices of the Type  $B_n$  Coxeter hyperplane arrangements;
- (generalizing the previous two items) the Dowling lattices [7, 6]  $Q_n(G)$  associated to any finite group  $G$ .

Hence, these are all cores of upho lattices. We discuss these examples in detail, providing explicit descriptions of the upho lattices for which they are cores. Figure 1 depicts one upho lattice produced via this construction.

---

<sup>1</sup>In fact, since the flag  $f$ -vector of any upho poset is determined by its rank generating function (see [21, §3]), the core of an upho lattice determines its entire flag  $f$ -vector.



FIGURE 1. Partitions of sets of the form  $\{1, 2, \dots, n\}$  into 2 blocks, ordered by refinement. This is an upho lattice with core  $\Pi_3$ .



FIGURE 2. The dual braid monoid  $\langle a, b, c \mid ab = bc = ca \rangle$  associated to the symmetric group  $S_3$ . This is an upho lattice with core the noncrossing partition lattice of  $S_3$ .

In addition to combinatorial constructions, we also explore algebraic constructions of upho lattices. Monoids provide one algebraic source of upho lattices, as the following lemma explains.

**Lemma 1.4.** (*c.f.* [8, Lemma 5.1]) *Let  $M$  be a finitely generated monoid whose defining relations are homogeneous. If  $M$  is left cancellative and every pair of elements in  $M$  have a least common right multiple, then  $(M, \leq_L)$  is an upho lattice, where  $\leq_L$  denotes the partial order of left divisibility.*

A class of monoids satisfying the conditions of Lemma 1.4 are the (homogeneous) Garside monoids [4, 3]. The core of a Garside monoid consists of its simple elements. Examples of lattices of simple elements in Garside monoids include:

- the weak order of a finite Coxeter group  $W$ ;
- the noncrossing partition lattice of a finite Coxeter group  $W$ .

Hence, these are also cores of upho lattices. Figure 2 depicts an upho lattice of this form. We review these examples coming from Garside and Coxeter theory in detail.

On the negative side, we show that there are various obstructions which prevent arbitrary finite graded lattices from being realized as cores of upho lattices. There are restrictions on the characteristic polynomial of the lattice coming from

the equation (1.1). There are also some structural obstructions, requiring the lattice to be partly self-similar. These obstructions allow us to show, for instance, that the following plausible candidates cannot in fact be realized as cores:

- the face lattice of the  $n$ -dimensional cross polytope, and the face lattice of the  $n$ -dimensional hypercube, for  $n \geq 3$ ;
- the bond lattice of the cycle graph  $C_n$ , for  $n \geq 4$ ;
- (generalizing the previous item) the lattice of flats of the uniform matroid  $U(n, k)$ , for  $2 < k < n$ .

The upshot is that Question 1.2 is quite subtle: it can be difficult to recognize when a given finite graded lattice is the core of an upho lattice. Many well-behaved finite lattices are cores of upho lattices, but many too are not.

To conclude this introduction, let us also discuss future directions we are pursuing. A question naturally suggested by our work here is the following.

**Question 1.5.** For a finite graded lattice  $L$ , let  $\kappa(L)$  denote the cardinality of the collection of (isomorphism classes of) upho lattices  $\mathcal{L}$  with core  $L$ . How does  $\kappa(L)$  behave? For example, is  $\kappa(L)$  finite for each  $L$ ? Or, is  $\kappa(L)$  bounded by a function of the rank of  $L$ ?

Notice that Question 1.5 essentially asks how big the difference between answering Question 1.2 and resolving Problem 1.1 is. In work in progress joint with Joel Lewis [12], we address Question 1.5. On the one hand, we will show that  $\kappa(L)$  is finite if  $L$  has no automorphisms, suggesting that possibly  $\kappa(L)$  is finite for all lattices  $L$ . On the other hand, we will show that  $\kappa(L)$  is unbounded even when we restrict to lattices  $L$  of rank two.

Finally, if completely resolving Problem 1.1 is too difficult, we might instead hope to classify some subvarieties of upho lattices. Two of the most important subvarieties of lattices are the distributive lattices and the modular lattices. In planned future work, we will explore distributive and modular upho lattices.

It is easy to see from the representation theorem for locally finite distributive lattices that the only upho distributive lattices are  $\mathbb{N}^d$ . Upho modular lattices are more interesting. Stanley observed that if  $R$  is a discrete valuation ring with finite residue field, then the lattice of full rank submodules of the free module  $R^d$  gives an upho modular lattice. Slightly extending Stanley's observation, we can show that any (sufficiently symmetric) affine building gives rise to an upho modular lattice. Conjecturally (c.f. [8, Conjecture 1.1]) all upho modular lattices are of this form.

The rest of the paper is structured as follows. In Section 2, we go over some definitions and preliminary results. In Section 3, we construct upho lattices from uniform sequences of supersolvable lattices. In Section 4, we explain how monoids give rise to upho lattices. Finally, in Section 5, we discuss obstructions to realizing a finite graded lattice as the core of an upho lattice.

**Acknowledgments.** I thank the following people for useful comments related to this work: Yibo Gao, Joel Lewis, Vic Reiner, David Speyer, Richard Stanley, Benjamin Steinberg, Nathan Williams, and Gjergji Zaimi. Sage mathematical software [22] was an important computational aid for this research.

## 2. PRELIMINARIES

In this section, we review some basics regarding posets and upho posets. We generally stick to standard notation for posets, as laid out of instance in [18, §3]. We use  $\mathbb{N} := \{0, 1, \dots\}$  to denote the natural numbers,  $\mathbb{Z}$  to denote the integers,  $\mathbb{Q}$  the rationals, and  $\mathbb{R}$  the reals.

**2.1. Poset basics.** Let  $P = (P, \leq)$  be a poset. When working with multiple posets we write  $\leq_P$  for clarity. We use standard conventions like writing  $y \geq x$  to mean  $x \leq y$ , writing  $x < y$  to mean  $x \leq y$  and  $x \neq y$ , and so on. We also routinely identify any subset  $S \subseteq P$  with the corresponding induced subposet  $S = (S, \leq)$ .

**2.1.1. Basic terminology.** An *interval* of  $P$  is a subset  $[x, y] := \{z \in P : x \leq z \leq y\}$  for  $x \leq y \in P$ . The poset  $P$  is *locally finite* if every interval of  $P$  is finite.

For  $x, y \in P$ , we say  $x$  is *covered* by  $y$ , written  $x < y$ , if  $x < y$  and there is no  $z \in P$  with  $x < z < y$ . The *Hasse diagram* of  $P$  is the directed graph whose vertices are the elements of  $P$  with an edge from  $x$  to  $y$  when  $x < y$ . We draw the Hasse diagram of  $P$  in the plane, with  $x$  below  $y$  if there is an edge from  $x$  to  $y$  (and therefore we do not draw the arrows on the edges). Any locally finite poset is determined by its Hasse diagram.

A *chain* of  $P$  is a totally ordered subset, i.e., a subset  $C \subseteq P$  for which any two elements in  $C$  are comparable. An *antichain* of  $P$  is a subset  $A \subseteq P$  for which any two elements in  $A$  are incomparable. We say a chain is *maximal* if it is maximal by inclusion among chains, and similarly for antichains.

An *order filter* of  $P$  is an upwards-closed subset, i.e., a subset  $F \subseteq P$  such that if  $x \in F$  and  $x \leq y$  then  $y \in F$ . Dually, an *order ideal* of  $P$  is a downwards-closed subset, i.e., a subset  $I \subseteq P$  such that if  $y \in I$  and  $x \leq y$  then  $x \in I$ . An order filter (respectively, order ideal) is *principal* if it is of the form  $V_p := \{q \in P : p \leq q\}$  (resp., of the form  $\Lambda_p := \{q \in P : q \leq p\}$ ) for some  $p \in P$ .

A *minimum* of  $P$ , which we always denote by  $\hat{0} \in P$ , is an element with  $\hat{0} \leq x$  for all  $x \in P$ . Dually, a *maximum*, denoted  $\hat{1} \in P$ , is an element with  $x \leq \hat{1}$  for all  $x \in P$ . Clearly, minimums and maximums are unique if they exist. If  $P$  has a minimum  $\hat{0}$ , then we call  $s \in P$  an *atom* if  $\hat{0} < s$ . Dually, if  $P$  has a maximum  $\hat{1}$ , then we call  $t \in P$  a *coatom* if  $t < \hat{1}$ .

**2.1.2. New posets from old.** The *dual poset*  $P^*$  of  $P$  is the poset with the same set of elements but with the opposite order, i.e.,  $x \leq_{P^*} y$  if and only if  $y \leq_P x$ . We say  $P$  is *self-dual* if it is isomorphic to its dual poset.

Now let  $Q$  be another poset. The *direct sum*  $P + Q$  of  $P$  and  $Q$  is the poset whose set of elements is the (disjoint) union  $P \sqcup Q$ , with  $x \leq_{P+Q} y$  if either  $x, y \in P$  and  $x \leq_P y$ , or  $x, y \in Q$  and  $x \leq_Q y$ . We say  $P$  is *connected* if it cannot be written as a nontrivial direct sum. For a positive integer  $n \geq 1$ , we denote the direct sum of  $n$  copies of  $P$  by  $nP$ .

Meanwhile, the *ordinal sum*  $P \oplus Q$  of  $P$  and  $Q$  is the poset whose elements are the (disjoint) union  $P \sqcup Q$ , with  $x \leq_{P \oplus Q} y$  if either  $x, y \in P$  and  $x \leq_P y$ , or  $x, y \in Q$

and  $x \leq_Q y$ , or  $x \in P$  and  $y \in Q$ . In other words,  $P + Q$  is obtained by placing  $P$  and  $Q$  “side-by-side,” while  $P \oplus Q$  is obtained by placing  $P$  “below”  $Q$ .

The *direct product*  $P \times Q$  of  $P$  and  $Q$  is the poset whose set of elements is the (Cartesian) product  $P \times Q$ , with  $(p_1, q_1) \leq_{P \times Q} (p_2, q_2)$  if  $p_1 \leq_P p_2$  and  $q_1 \leq_Q q_2$ . We say  $P$  is *indecomposable* if it cannot be written as a nontrivial direct product. For a positive integer  $n \geq 1$ , we denote the direct product of  $n$  copies of  $P$  by  $P^n$ .

**2.1.3. Möbius functions.** Suppose for the moment that  $P$  is locally finite, and let  $\text{Int}(P)$  denote the set of intervals of  $P$ . The *Möbius function*  $\mu: \text{Int}(P) \rightarrow \mathbb{Z}$  of  $P$  is defined recursively by

$$\mu(x, x) = 1 \text{ for all } x \in P; \quad \mu(x, y) = - \sum_{x \leq z < y} \mu(x, z) \text{ for all } x < y \in P,$$

where we use the standard notational shorthand  $\mu(x, y) = \mu([x, y])$ . The most important application of Möbius functions is the Möbius inversion formula (see [18, §3.7]), a kind of generalization of the principle of inclusion-exclusion to any poset.

The Möbius function of a product of posets decomposes as a product of Möbius functions. In other words, we have  $\mu_{P \times Q}((p_1, q_1), (p_2, q_2)) = \mu_P(p_1, p_2) \cdot \mu_Q(q_1, q_2)$  for all  $(p_1, q_1) \leq (p_2, q_2) \in P \times Q$  (see [18, Proposition 3.8.2]).

**2.1.4. Lattices.** For  $x, y \in P$ , an *upper bound* of  $x$  and  $y$  is a  $z \in P$  with  $x \leq z$  and  $y \leq z$ , and the *join* (or *least upper bound*) of  $x$  and  $y$ , denoted  $x \vee y$ , is the minimum among all upper bounds of  $x$  and  $y$ , if such a minimum exists. Dually, a *lower bound* of  $x$  and  $y$  is a  $z \in P$  with  $z \leq x$  and  $z \leq y$ , and the *meet* (or *greatest lower bound*) of  $x$  and  $y$ , denoted  $x \wedge y$ , is the maximum among all lower bounds of  $x$  and  $y$ , if such a maximum exists. If  $x \vee y$  exists for every  $x, y \in P$ , then  $P$  is a *join semilattice*. Dually, if  $x \wedge y$  exists for every  $x, y \in P$ , then  $P$  is a *meet semilattice*. The poset  $P$  is a *lattice* if it is both a join and meet semilattice.

Now let  $L$  be a lattice. The operations of  $\vee$  and  $\wedge$  are associative and commutative, and therefore for any finite, nonempty subset  $S = \{x_1, \dots, x_n\} \subseteq L$  we can set  $\bigvee S := x_1 \vee \dots \vee x_n$  and  $\bigwedge S := x_1 \wedge \dots \wedge x_n$ . If  $L$  has a minimum  $\hat{0}$  then by convention we set  $\bigvee \emptyset := \hat{0}$ , and dually if  $L$  has a maximum  $\hat{1}$  we set  $\bigwedge \emptyset := \hat{1}$ . A *finite* lattice  $L$  always has a minimum  $\hat{0} = \bigwedge L$  and a maximum  $\hat{1} = \bigvee L$ .

If  $L$  has a minimum  $\hat{0}$  and a maximum  $\hat{1}$ , then a *complement* of an element  $x \in L$  is a  $y \in L$  with  $x \wedge y = \hat{0}$  and  $x \vee y = \hat{1}$ .

The lattice  $L$  is *distributive* if the operation of meet distributes over that of join, i.e.,  $x \wedge (y \vee z) = (x \wedge y) \vee (x \wedge z)$  for all  $x, y, z \in L$ . It is well-known that this is also equivalent to join distributing over meet, i.e.,  $x \vee (y \wedge z) = (x \vee y) \wedge (x \vee z)$  for all  $x, y, z \in L$ .

**Example 2.1.** For any poset  $P$ , we use  $J(P)$  to denote the poset of order ideals of  $P$ , ordered by inclusion. Then,  $J(P)$  is always a distributive lattice, where the operations of join and meet are union and intersection, respectively. As a variant of this construction, we use  $J_{\text{fin}}(P)$  to denote the finite order ideals of  $P$ , which again always gives a distributive lattice. We note that  $J(P + Q) = J(P) \times J(Q)$  and similarly  $J_{\text{fin}}(P + Q) = J_{\text{fin}}(P) \times J_{\text{fin}}(Q)$ .

The lattice  $L$  is *modular* if whenever  $x \leq y \in L$ , we have  $x \vee (z \wedge y) = (x \vee z) \wedge y$  for all  $z \in L$ . Observe that modularity is a weaker condition than distributivity: all distributive lattices are modular, but most modular lattices are not distributive.

We note that the product  $L_1 \times L_2$  of two lattices  $L_1$  and  $L_2$  remains a lattice, and similarly if we append the adjectives “distributive” or “modular.” Also, any interval in a lattice is a lattice, and similarly if we append the adjectives “distributive” or “modular.” These properties follow from the fact that lattices, distributive lattices, and modular lattices, are varieties in the sense of universal algebra.

**2.1.5. Convention for finite versus infinite posets.** We will routinely work with both finite and infinite posets, although the posets will always be at least locally finite. For clarity, we now use the following convention: normal script letters (like  $P$  or  $L$ ) denote finite posets, while caligraphic letters (like  $\mathcal{P}$  or  $\mathcal{L}$ ) denote infinite posets.

**2.2. Finite graded posets.** Let  $P$  be a finite poset. For a nonnegative integer  $n \geq 0$ , we say that  $P$  is *n-graded* if  $P$  has a minimum  $\hat{0}$ , a maximum  $\hat{1}$ , and we can write  $P = \bigsqcup_{i=0}^n P_i$  as a disjoint union such that every maximal chain of  $P$  is of the form  $\hat{0} = x_0 < x_1 < \cdots < x_n = \hat{1}$  with  $x_i \in P_i$ . In this case, the *rank function*  $\rho: P \rightarrow \mathbb{N}$  is defined by setting  $\rho(x) := i$  for  $x \in P_i$ . Equivalently,  $\rho(\hat{0}) = 0$  and  $\rho(y) = \rho(x) + 1$  whenever  $x < y \in P$ .

**Example 2.2.** For any positive integer  $n \geq 1$ , we let  $[n] := \{1, 2, \dots, n\}$ . We view  $[n]$  as a poset, with the usual total order. This chain poset  $[n]$  is the most basic example of a finite  $(n - 1)$ -graded poset.

We say that the finite poset  $P$  is *graded* if it is  $n$ -graded for some  $n$ . In this case, we say that the *rank* of  $P$  is  $n$  and, slightly abusing notation, write  $\rho(P) := n$ . If  $P$  and  $Q$  are two finite graded posets, then their product  $P \times Q$  is also graded of rank  $\rho(P \times Q) = \rho(P) + \rho(Q)$ . Also, any interval in a finite graded poset is graded.

**2.2.1. Generating polynomials for finite graded posets.** Now assume that  $P$  is graded. The *rank generating polynomial* of  $P$  is

$$F(P; x) := \sum_{p \in P} x^{\rho(p)}.$$

The *characteristic polynomial* of  $P$  is

$$\chi(P; x) := \sum_{p \in P} \mu(\hat{0}, p) x^{\rho(P) - \rho(p)}.$$

The exponent of  $x$  in each term of the characteristic polynomial  $\chi(P; x)$  records the corank  $\rho(P) - \rho(p)$  of the element  $p \in P$ . Using the corank in the characteristic polynomial is very standard, but, for reasons that will become clear soon, we need a version of the characteristic polynomial where the exponent records the usual rank instead. Hence, we define the *reciprocal characteristic polynomial* of  $P$  to be

$$\chi^*(P; x) := \sum_{p \in P} \mu(\hat{0}, p) x^{\rho(p)}.$$

Observe that  $\chi^*(P; x) = x^{\rho(P)} \cdot \chi(P; x^{-1})$ .

These invariants of finite graded posets all play nicely with products. Namely,  $F(P \times Q; x) = F(P; x) \cdot F(Q; x)$  and  $\chi^*(P \times Q; x) = \chi^*(P; x) \cdot \chi^*(Q; x)$ .

**2.2.2. Finite graded lattices.** Here we are most interested in finite graded lattices.

**Example 2.3.** The rank  $n$  (*finite*) *Boolean lattice*  $B_n$  is the poset of subsets of  $[n]$ , ordered by inclusion.  $B_n$  is a finite graded lattice, with  $\rho(S) = \#S$  for all  $S \in B_n$ . Its Möbius function is given by  $\mu(S, T) = (-1)^{\#T \setminus S}$  for all  $S \leq T \in B_n$ . Hence,  $F(B_n; x) = \sum_{k=0}^n \binom{n}{k} x^k = (1+x)^n$  and  $\chi^*(B_n; x) = \sum_{k=0}^n (-1)^k \binom{n}{k} x^k = (1-x)^n$ . These formulas can also be seen from the fact that  $B_n = J(n[1]) = [2]^n$ .

**Example 2.4.** Birkhoff's representation theorem for finite distributive lattices says that every finite distributive lattice  $L$  has the form  $L = J(P)$  for a unique finite poset  $P$  (the subposet of *join-irreducible* elements of  $L$ ); see [1, §3.3] or [18, §3.4]. So let  $L = J(P)$  be a finite distributive lattice. Then  $L$  is graded, with  $\rho(I) = \#I$  for  $I \in J(P)$ . Its Möbius function is given by

$$\mu(I, I') = \begin{cases} (-1)^{\#I' \setminus I} & \text{if } I' \setminus I \text{ is an antichain of } P; \\ 0 & \text{otherwise,} \end{cases}$$

for  $I \leq I' \in J(P)$  (see [18, Example 3.9.6]). Hence,  $F(L; x) = \sum_{I \in J(P)} x^{\#I}$  and  $\chi^*(L; x) = \sum_{I \in \min(P)} (-x)^{\#I} = (1-x)^{\#\min(P)}$ , where  $\min(P)$  is the set of minimal elements of  $P$ . Observe how this example generalizes Example 2.3.

Example 2.4 explains that all finite distributive lattices are graded. More generally, all finite modular lattices are graded. In fact, a finite lattice  $L$  is modular if and only if  $L$  is graded and  $\rho(p) + \rho(q) = \rho(p \vee q) + \rho(p \wedge q)$  for all  $p, q \in L$  (see, e.g., [1, §2.8] or [18, §3.3]).

**Example 2.5.** Let  $q$  be a prime power, and  $\mathbb{F}_q$  the finite field with  $q$  elements. We denote by  $B_n(q)$  the poset of  $\mathbb{F}_q$ -subspaces of the vector space  $\mathbb{F}_q^n$ , ordered by inclusion. This *subspace lattice*  $B_n(q)$ , also known as the  *$q$ -analogue* of  $B_n$ , is a finite modular lattice. Its rank function is  $\rho(U) = \dim(U)$  for  $U \in B_n(q)$ , and its Möbius function is  $\mu(U, V) = (-1)^k q^{\binom{k}{2}}$ , where  $k = \dim(V) - \dim(U)$ , for  $U \leq V \in B_n(q)$  (see [18, Example 3.10.2]). Hence,  $F(B_n(q); x) = \sum_{k=0}^n \begin{bmatrix} n \\ k \end{bmatrix}_q x^k$  and  $\chi^*(B_n(q); x) = \sum_{k=0}^n (-1)^k q^{\binom{k}{2}} \begin{bmatrix} n \\ k \end{bmatrix}_q x^k = (1-x)(1-qx)(1-q^2x) \cdots (1-q^{n-1}x)$ . Here we used the standard notation for the  *$q$ -binomial coefficient*  $\begin{bmatrix} n \\ k \end{bmatrix}_q := \frac{[n]_q!}{[k]_q! [n-k]_q!}$ , where  $[n]_q := \frac{1-q^n}{1-q} = 1 + q + q^2 + \cdots + q^{n-1}$  and  $[n]_q! := [n]_q \cdot [n-1]_q \cdots [1]_q$ .

There are variants of the modular property for finite lattices that are very interesting from a combinatorial point of view. A finite lattice  $L$  is (*upper*) *semimodular* if it is graded and satisfies  $\rho(p) + \rho(q) \geq \rho(p \vee q) + \rho(p \wedge q)$  for all  $p, q \in L$ . The lattice  $L$  is *atomic* (or, in some references, *atomistic*) if every element is a join of atoms. Finally,  $L$  is *geometric* if it is both semimodular and atomic. For example, the modular lattices  $B_n$  and  $B_n(q)$  are geometric, since they are atomic. However, most geometric lattices are not modular. Geometric lattices are intensely studied, because they are precisely the lattices of flats of matroids (see [18, §3.3]). Any



interval in a semimodular lattice is semimodular, and, less obviously, any interval in a geometric lattice is geometric (see, e.g., [1, §8.9, Exercise 7] and [18, Proposition 3.3.3]).

**Example 2.6.** Recall that a *partition* of a set  $X$  is a collection  $\pi = \{B_1, \dots, B_k\}$  of nonempty subsets of  $X$  ( $\emptyset \subsetneq B_1, \dots, B_k \subseteq X$ ) that are pairwise disjoint ( $B_i \cap B_j = \emptyset$  for  $i \neq j$ ) and whose union is  $X$  ( $\cup_{i=1}^k B_i = X$ ). The sets  $B_i \in \pi$  are called the *blocks* of  $\pi$ . The *partition lattice*  $\Pi_n$  is the poset of partitions of  $[n]$ , ordered by refinement. In other words, for two partitions  $\pi$  and  $\pi'$  of  $[n]$ , we have  $\pi \leq \pi'$  if every block  $B \in \pi$  satisfies  $B \subseteq B'$  for some block  $B' \in \pi'$ . The partition lattice is a geometric lattice of rank  $\rho(\Pi_n) = n - 1$ , with  $\rho(\pi) = n - \#\pi$  for  $\pi \in \Pi_n$ . Hence,  $F(\Pi_n; x) = \sum_{k=0}^n S(n, n-k)x^k$ , where  $S(n, k)$  are the *Stirling number of the second kind*. Furthermore, it is well-known that  $\chi^*(\Pi_n; x) = \sum_{k=0}^n s(n, n-k)x^k = (1-x)(1-2x) \cdots (1-(n-1)x)$ , where  $s(n, k)$  are the *(signed) Stirling number of the first kind* (see [18, Example 3.10.4]).

**2.3. Infinite graded posets.** Let  $\mathcal{P}$  be an infinite poset. We say  $\mathcal{P}$  is  *$\mathbb{N}$ -graded* if  $\mathcal{P}$  has a minimum  $\hat{0}$  and we can write  $\mathcal{P} = \bigsqcup_{i=0}^{\infty} P_i$  as a disjoint union such that every maximal chain of  $\mathcal{P}$  is of the form  $\hat{0} = x_0 \leq x_1 \leq x_2 \leq \cdots$  with  $x_i \in P_i$ . In this case, the *rank function*  $\rho: \mathcal{P} \rightarrow \mathbb{N}$  is defined by setting  $\rho(x) := i$  for  $x \in P_i$ . Equivalently,  $\rho(\hat{0}) = 0$  and  $\rho(y) = \rho(x) + 1$  whenever  $x \leq y \in \mathcal{P}$ .

If  $\mathcal{P}$  and  $\mathcal{Q}$  are  $\mathbb{N}$ -graded, then  $\mathcal{P} \times \mathcal{Q}$  is  $\mathbb{N}$ -graded. Also, any interval in a *locally finite*  $\mathbb{N}$ -graded poset is a finite graded poset.

**2.3.1. Generating functions for infinite graded posets.** Let  $\mathcal{P}$  be an  $\mathbb{N}$ -graded poset. In order to define sensible analogs of the rank generating and characteristic polynomials for  $\mathcal{P}$ , we need to make a further finiteness assumption. So let us say that  $\mathcal{P}$  is *finite type* if  $\{p \in \mathcal{P}: \rho(p) = i\}$  is finite for each  $i \in \mathbb{N}$ . Observe that a finite type  $\mathbb{N}$ -graded poset is locally finite (but finite type is a stronger requirement than just being locally finite).

So now assume that  $\mathcal{P}$  is a finite type  $\mathbb{N}$ -graded poset. Then we define the *rank generating function* of  $\mathcal{P}$  to be

$$F(\mathcal{P}; x) = \sum_{p \in \mathcal{P}} x^{\rho(p)},$$

a formal power series in the variable  $x$ . And we define the *characteristic generating function* of  $\mathcal{P}$  to be

$$\chi^*(\mathcal{P}; x) = \sum_{p \in \mathcal{P}} \mu(\hat{0}, p) x^{\rho(p)},$$

again, a formal power series. We write  $\chi^*(\mathcal{P}; x)$  with an asterisk to emphasize that the characteristic generating function of an infinite poset  $\mathcal{P}$  uses rank in the exponent, like the *reciprocal* characteristic polynomial  $\chi^*(P; x)$  of a finite poset  $P$ .

Again, these invariants play nicely with products:  $F(\mathcal{P} \times \mathcal{Q}; x) = F(\mathcal{P}; x) \cdot F(\mathcal{Q}; x)$  and  $\chi^*(\mathcal{P} \times \mathcal{Q}; x) = \chi^*(\mathcal{P}; x) \cdot \chi^*(\mathcal{Q}; x)$ .

**Example 2.7.** The set of natural numbers  $\mathbb{N} = \{0, 1, \dots\}$ , with their usual total order, is the most basic example of an  $\mathbb{N}$ -graded poset. In fact,  $\mathbb{N}$  is a finite type  $\mathbb{N}$ -graded lattice with  $F(\mathbb{N}; x) = \sum_{k=0}^{\infty} x^k = \frac{1}{1-x}$  and  $\chi^*(\mathbb{N}; x) = 1 - x$ . Hence, for any positive integer  $n \geq 1$ ,  $\mathbb{N}^n$  is a finite type  $\mathbb{N}$ -graded lattice with  $F(\mathbb{N}^n; x) = \frac{1}{(1-x)^n}$  and  $\chi^*(\mathbb{N}^n; x) = (1 - x)^n$ .

**2.4. Upho posets.** Let  $\mathcal{P}$  be an infinite poset.  $\mathcal{P}$  is *upper homogeneous*, or “*upho*,” if for every  $p \in \mathcal{P}$ , the principal order filter  $V_p = \{q \in \mathcal{P} : q \geq p\}$  is isomorphic to  $\mathcal{P}$ . To avoid trivialities, let us also require that  $\mathcal{P}$  has at least two elements.

**Example 2.8.** The natural numbers  $\mathbb{N}$  form an upho poset. Similarly, the nonnegative rational numbers  $\{x \in \mathbb{Q} : x \geq 0\}$ , with their usual order, form an upho poset. And ditto for the nonnegative real numbers  $\{x \in \mathbb{R} : x \geq 0\}$ .

**Example 2.9.** Let  $X$  be any infinite set. Then the poset of finite subsets of  $X$ , ordered by inclusion, is upho.

In this paper we are primarily concerned with upho posets (in fact, upho lattices). In order to be able to apply the tools of enumerative and algebraic combinatorics to study these posets, we must impose some finiteness conditions on them. Hence, from now on, *all upho posets are assumed finite type  $\mathbb{N}$ -graded unless otherwise specified*. Of the preceding examples, only  $\mathbb{N}$  is finite type  $\mathbb{N}$ -graded.

The product  $\mathcal{P} \times \mathcal{Q}$  of two upho posets  $\mathcal{P}$  and  $\mathcal{Q}$  remains upho. So, for instance,  $\mathbb{N}^n$  is an upho lattice for any  $n \geq 1$ . We will soon see many more examples of upho lattices, but for now  $\mathbb{N}^n$  is a good prototypical example to have in mind.

**Remark 2.10.** Upho posets were introduced by Stanley [19, 21]. Stanley was mainly interested in *planar* upho posets (i.e., those with planar Hasse diagrams). In particular, he was interested in counting the maximal chains in  $[\hat{0}, p]$  for all  $p \in \mathcal{P}$ , for various planar upho posets  $\mathcal{P}$ . When  $\mathcal{P} = \mathbb{N}^2$ , these numbers form Pascal’s triangle. Thus, Stanley used these chain counts for other planar upho posets to produce analogues of Pascal’s triangle [20, 21]. All planar upho posets are meet semilattices, but most are not lattices. In fact, it is not hard to see that  $\mathbb{N}$  and  $\mathbb{N}^2$  are the only planar upho lattices. Planar upho posets have a rather simple structure, as described in [8]. We will see that upho lattices can have a very intricate structure.

**2.4.1. Rank and characteristic generating functions of upho posets.** The following important result on rank generating functions of upho posets can be proved by a simple application of Möbius inversion.

**Theorem 2.11** ([10, Theorem 1]). *For any upho poset  $\mathcal{P}$ ,  $F(\mathcal{P}; x) = \chi^*(\mathcal{P}; x)^{-1}$ .*

**2.4.2. Upho lattices and their cores.** Now suppose that  $\mathcal{L}$  is an upho lattice. Then we define the *core* of  $\mathcal{L}$  to be  $L := [\hat{0}, s_1 \vee s_2 \vee \dots \vee s_r] \subseteq \mathcal{L}$ , where  $s_1, \dots, s_r$  are the atoms of  $\mathcal{L}$ . Evidently, the core of an upho lattice is a finite graded lattice. The point of the core is the following corollary, which can be proved for instance using Rota’s cross-cut theorem (see [18, Corollary 3.9.4]).

**Corollary 2.12** ([10, Corollary 6]). *Let  $\mathcal{L}$  be an upho lattice with core  $L$ . Then  $\chi^*(\mathcal{L}; x) = \chi^*(L; x)$ . Hence, from Theorem 2.11, we conclude  $F(\mathcal{L}; x) = \chi^*(L; x)^{-1}$ .*

Note that Corollary 2.12 was stated as equation (1.1) in the introduction.

**Example 2.13.** For any  $n \geq 1$ ,  $\mathbb{N}^n$  is an upho lattice with core  $B_n$ , and indeed  $F(\mathbb{N}^n; x) = \frac{1}{(1-x)^n} = \chi^*(B_n; x)^{-1}$ .

With all the terminology and preliminary results fully explained, we now return to Question 1.2: which finite graded lattices  $L$  arise as cores of upho lattices  $\mathcal{L}$ ? For example, we just saw that the Boolean lattices  $B_n$  do. We will explore this question in the next three sections.

### 3. UPHO LATTICES FROM SEQUENCES OF FINITE LATTICES

In this section, we will develop a method for producing upho lattices from limits of sequences of finite graded lattices which are appropriately embedded in one another. In order to make “appropriately embedded in one another” precise, we will need two notions from the theory of finite lattices: supersolvability, as introduced by Stanley in [16], and uniformity, as introduced by Dowling in [6].

**3.1. Supersolvable semimodular lattices.** Let  $L$  be a (finite) semimodular lattice. An element  $p \in L$  is *modular* if  $\rho(p) + \rho(q) = \rho(p \vee q) + \rho(p \wedge q)$  for all  $q \in L$ . For example, if  $L$  is modular, then every element is modular. An important property of modularity is that it is transitive on “lower” intervals:

**Proposition 3.1** (See [15] or [17, Proposition 4.10(b)]). *Let  $L$  be a semimodular lattice. If  $x$  is modular in  $L$  and  $y$  is modular in  $[\hat{0}, x]$ , then  $y$  is modular in  $L$ .*

The semimodular lattice  $L$  is *supersolvable* if it possesses a maximal chain  $\hat{0} = x_0 < x_1 < \dots < x_n = \hat{1}$  of modular elements. As suggested by Proposition 3.1, we can build up a maximal chain of modular elements from the top down, coatom-by-coatom. For this reason, supersolvable lattices have a recursive structure. Moreover, as shown by Stanley [16], they enjoy many remarkable enumerative properties, the most prominent being that their characteristic polynomials factor.

**Theorem 3.2** (Stanley [15] [16, Theorem 4.1]). *Let  $L$  be a supersolvable semimodular lattice with maximal chain of modular elements  $x_0 < x_1 < \dots < x_n$ . For  $i = 1, \dots, n$ , set*

$$a_i := \#\{\text{atoms } s \in L : s \leq x_i, s \not\leq x_{i-1}\}.$$


*Then  $\chi^*(L; x) = (1 - a_1x)(1 - a_2x) \cdots (1 - a_nx)$ .*

**Example 3.3.** The partition lattice  $\Pi_n$  is a supersolvable geometric lattice. Indeed,  $\pi_0 < \pi_1 < \dots < \pi_{n-1} \in \Pi_n$  is a maximal chain of modular elements, where

$$\pi_i := \{\{1, 2, \dots, i+1\}, \{i+2\}, \{i+3\}, \dots, \{n\}\}$$

for  $i = 0, 1, \dots, n-1$ . Here  $a_i = i$  for  $i = 1, \dots, n-1$ , so Theorem 3.2 gives that  $\chi^*(\Pi_n; x) = (1-x)(1-2x) \cdots (1-(n-1)x)$ , in agreement with what we said earlier.

**Remark 3.4.** In [16], Stanley defined the notion of supersolvability more generally for any finite lattice, not necessarily semimodular. It is possible that the results in this section could be extended to this broader class of (not necessarily semimodular) supersolvable lattices. However, the examples that we know all involve geometric lattices, so it is unclear, at the moment, what this greater generality would buy us.

FIGURE 3. The “trimmed” partition lattice  $\Pi_4^{(2)}$ .

Next, we describe a procedure for “trimming” a supersolvable semimodular lattice to produce another one. Although this procedure is quite natural, we have not seen it anywhere in the literature. We use this trimming procedure later to guarantee that a certain limit poset is finite type N-graded.

So let  $L$  be a supersolvable semimodular lattice with a fixed maximal chain of modular elements  $x_0 < x_1 < \dots < x_n$ . Relative to this chain, we define  $\nu: L \rightarrow \mathbb{N}$  by  $\nu(x) := \min\{i: x \leq x_i\}$  for all  $x \in L$ . And then for a positive integer  $k \geq 1$ , we define

$$L^{(k)} := \{x \in L: \nu(x) - \rho(x) < k\}.$$

For instance,  $L^{(1)} = \{x_0, x_1, \dots, x_n\}$  and  $L^{(n)} = L$ . Note that  $L^{(i)} \subseteq L^{(j)}$  for  $i \leq j$ .

**Lemma 3.5.** *For any  $k \geq 1$ , the subposet  $L^{(k)}$  of  $L$  is again a supersolvable semimodular lattice, with maximal chain of modular elements  $x_0 < x_1 < \dots < x_n$ .*

The rank of an element in  $L^{(k)}$  is the same as its rank in  $L$ . And the join of two elements in  $L^{(k)}$  is the same as their join in  $L$ . But the meet of two elements in  $L^{(k)}$  may be different than their meet in  $L$ , so  $L^{(k)}$  is not in general a sublattice of  $L$ .

**Example 3.6.** Again consider the partition lattice  $\Pi_n$ , with maximal chain of modular elements  $\pi_0 < \pi_1 < \dots < \pi_{n-1}$  as described above. Then, for any  $k \geq 1$ ,

$$\Pi_n^{(k)} = \{\pi \in \Pi_n: \max\{i: i \text{ is in a non-singleton block of } \pi\} < n + k - \#\pi\}.$$

(Recall that a block is called a *singleton* if it has one element. Also, above we use the convention  $\max(\emptyset) := 0$ .) For instance, Figure 3 depicts  $\Pi_4^{(2)}$ . In this figure we use the shorthand of writing a partition  $\pi = \{B_1, \dots, B_k\}$  as  $B_1 | B_2 | \dots | B_k$ . The perceptive reader may compare this figure to Figure 1.

*Proof of Lemma 3.5.* We start with two important claims.

**Claim 1:** For  $x, y \in L^{(k)}$ , the join  $x \vee y$ , taken in  $L$ , is in  $L^{(k)}$ .

**Claim 2:** For  $x \in L^{(k)}$  and  $i = 0, 1, \dots, n$ , the meet  $x \wedge x_i$ , taken in  $L$ , is in  $L^{(k)}$ .

*Proof of Claim 1:* Let  $x, y \in L^{(k)}$ . Set  $c := \max(\nu(x), \nu(y))$ ,  $d := \max(\rho(x), \rho(y))$ . Since  $\nu(x) - \rho(x) < k$  and  $\nu(y) - \rho(y) < k$ , we have  $c + d < k$ . And since  $x_c$  is an upper bound for  $x$  and  $y$ , we have  $\nu(x \vee y) \leq c$ . Similarly, since  $x \vee y$  is greater than or equal to  $x$  and to  $y$ , we have  $\rho(x \vee y) \geq d$ . Thus,  $\nu(x \vee y) - \rho(x \vee y) \leq c - d < k$ .

*Proof of Claim 2:* Let  $x \in L^{(k)}$  and  $i \in \{0, 1, \dots, n-1\}$ . If  $x \leq x_i$ , then  $x \wedge x_i = x$  and the claim is clear. So suppose that  $x \not\leq x_i$ . Since  $x_{\nu(x)}$  is an upper bound for  $x$

and  $x_i$ , we have  $\rho(x \vee x_i) \leq \nu(x)$ . And since  $x_i$  is modular, we have  $\rho(x) + \rho(x_i) = \rho(x \vee x_i) + \rho(x \wedge x_i)$ , i.e.,  $\rho(x \wedge x_i) = \rho(x) + \rho(x_i) - \rho(x \vee x_i) \geq \rho(x) + i - \nu(x)$ . Meanwhile, since  $x_i$  is an upper bound for  $x \wedge x_i$ , we have  $\nu(x \wedge x_i) \leq i$ . Thus,  $\nu(x \wedge x_i) - \rho(x \wedge x_i) \leq \nu(x) - \rho(x) < k$ .

We proceed to prove the lemma. First, let us explain why  $L^{(k)}$  is a lattice. Notice Claim 1 already implies that  $L^{(k)}$  is a sub-join semilattice of  $L$ . Then, since  $L^{(k)}$  is a finite join semilattice with a  $\hat{0}$ , it is a lattice (see, e.g., [18, Proposition 3.3.1]).

Next, let us explain why  $L^{(k)}$  is graded. Let  $y_0 < y_1 < y_2 < \dots < y_m \in L^{(k)}$  be any chain. We will show that we can extend this chain to a chain of length  $n$ . To that end, let  $L'$  denote the sublattice of  $L$  generated by  $\{x_0, \dots, x_n, y_0, \dots, y_m\}$ . An important result of Stanley [16, Proposition 2.1] says that  $L'$  is a distributive lattice (in fact, this leads to an alternative characterization of supersolvable lattices). Since  $L'$  is distributive, every element in  $L'$  can be written as a join of elements of the form  $x_i \wedge y_j$ . Hence, Claims 1 and 2 combine to imply that  $L' \subseteq L^{(k)}$ . Then, again since  $L'$  is distributive and hence graded, the chain  $y_0 < y_1 < y_2 < \dots < y_m$  can be extended in  $L'$  (and thus in  $L^{(k)}$ ) to a maximal chain of length  $n$ . So indeed  $L^{(k)}$  is graded, and in fact each element in  $L^{(k)}$  has the same rank as it does in  $L$ .

Next, we show that  $L^{(k)}$  is semimodular. Let  $x, y \in L^{(k)}$ . As mentioned, the join of  $x$  and  $y$  in  $L^{(k)}$  is the same as its join in  $L$ . On the other hand, the meet of  $x$  and  $y$  in  $L^{(k)}$  can only be lower than in  $L$ . So, the quantity  $\rho(x \vee y) + \rho(x \wedge y)$  is smaller when computed in  $L^{(k)}$  versus  $L$ . Thus, the inequality  $\rho(x) + \rho(y) \geq \rho(x \vee y) + \rho(x \wedge y)$  remains true in  $L^{(k)}$ , and  $L^{(k)}$  is semimodular.

Finally, we explain why  $L^{(k)}$  is supersolvable, with maximal chain of modular elements  $x_0 < x_1 < \dots < x_n$ . Let  $y \in L^{(k)}$  and  $i \in \{0, 1, \dots, n-1\}$ . Claims 1 and 2 imply that  $x_i \vee y$  and  $x_i \wedge y$  are the same in  $L^{(k)}$  as they are in  $L$ , so  $\rho(x_i) + \rho(y) = \rho(x_i \vee y) + \rho(x_i \wedge y)$  remains true, and thus  $x_i$  is modular in  $L^{(k)}$ .  $\square$

**3.2. Uniform sequences of geometric lattices.** A sequence of finite posets  $P_0, P_1, P_2, \dots$  is a *uniform sequence* if (for each  $n = 0, 1, \dots$ ):

- $P_n$  is  $n$ -graded;
- $[a, \hat{1}_{P_n}]$  is isomorphic to  $P_{n-1}$  for all atoms  $a \in P_n$ .

From now on in this section,  $L_0, L_1, \dots$  is a uniform sequence of geometric lattices.

Uniform sequences also enjoy many remarkable enumerative properties, as first observed by Dowling [6]. For  $0 \leq j \leq i$ , define the numbers  $V(i, j)$  and  $v(i, j)$  by

$$\sum_{j=0}^i V(i, j) x^{i-j} = F(L_i; x), \quad \sum_{j=0}^i v(i, j) x^{i-j} = \chi^*(L_i; x).$$

These numbers  $V(i, j)$  and  $v(i, j)$  are called the *Whitney numbers of the second and first kind*, respectively, for our sequence of lattices  $L_n$ . By convention, let us also declare  $V(i, j) := v(i, j) := 0$  for  $j > i$ .

**Theorem 3.7** (Dowling [6, Theorem 6]; see also [18, Exercise 3.130]). *The infinite, lower unitriangular matrices  $[V(i, j)]_{0 \leq i, j \leq \infty}$  and  $[v(i, j)]_{0 \leq i, j \leq \infty}$  are inverses.*

**Example 3.8.** Taking  $L_n = \Pi_{n+1}$  gives a uniform sequence of geometric lattices. In this case, we have  $V(i, j) = S(i+1, j+1)$  and  $v(i, j) = s(i+1, j+1)$ , the Stirling numbers of the second and first kind. Thus, the two infinite, lower unitriangular in Theorem 3.7 are:

$$[S(n, k)]_{1 \leq n, k \leq \infty} = \begin{pmatrix} 1 & 0 & 0 & 0 & \cdots \\ 1 & 1 & 0 & 0 & \cdots \\ 1 & 3 & 1 & 0 & \cdots \\ 1 & 7 & 6 & 1 & \cdots \\ \vdots & \vdots & \vdots & \vdots & \ddots \end{pmatrix} [s(n, k)]_{1 \leq n, k \leq \infty} = \begin{pmatrix} 1 & 0 & 0 & 0 & \cdots \\ -1 & 1 & 0 & 0 & \cdots \\ 2 & -3 & 1 & 0 & \cdots \\ -6 & 11 & -6 & 1 & \cdots \\ \vdots & \vdots & \vdots & \vdots & \ddots \end{pmatrix}$$

It is a classical result that these matrices of Stirling numbers are inverses.

When the geometric lattices  $L_n$  in our uniform sequence are also supersolvable, Theorems 3.2 and 3.7 combine to yield a very strong enumerative corollary, as we now explain. First, we need a preparatory result about symmetric polynomials. Recall that the *complete homogeneous* and *elementary symmetric polynomials* in variables  $x_1, \dots, x_n$  are, respectively, given by

$$h_k(x_1, \dots, x_n) := \sum_{1 \leq i_1 \leq i_2 \leq \dots \leq i_k \leq n} x_{i_1} x_{i_2} \cdots x_{i_k};$$

$$e_k(x_1, \dots, x_n) := \sum_{1 \leq i_1 < i_2 < \dots < i_k \leq n} x_{i_1} x_{i_2} \cdots x_{i_k},$$

for  $k > 0$ . We also use the conventions  $h_0(x_1, \dots, x_n) := e_0(x_1, \dots, x_n) := 1$  and  $h_k(x_1, \dots, x_n) := e_k(x_1, \dots, x_n) := 0$  for  $k < 0$ .

**Proposition 3.9.** Define infinite, lower unitriangular matrices  $A = [a_{i,j}]_{0 \leq i,j \leq \infty}$  and  $B = [b_{i,j}]_{0 \leq i,j \leq \infty}$  by letting

$$a_{i,j} := h_{i-j}(x_1, \dots, x_{j+1});$$

$$b_{i,j} := (-1)^{i-j} e_{i-j}(x_1, \dots, x_i),$$

for all  $0 \leq i, j$ . Then  $A$  and  $B$  are inverses.

*Proof.* Define the matrix  $C = [c_{i,j}]_{0 \leq i,j \leq \infty}$  by  $C = BA$ . Clearly,  $c_{n,m} = 0$  for  $m > n$ . For  $m \leq n$ ,

$$c_{n,m} = \sum_{i=0}^{n-m} (-1)^i e_i(x_1, \dots, x_n) h_{n-m-i}(x_1, \dots, x_{m+1}).$$

If  $m = n$ , this certainly equals 1. So now suppose  $m < n$ . Then  $c_{n,m}$  is the coefficient of  $z^{n-m}$  in

$$\left( \sum_{k=0}^{\infty} (-1)^k e_k(x_1, \dots, x_n) z^k \right) \left( \sum_{k=0}^{\infty} h_k(x_1, \dots, x_{m+1}) z^k \right) = \prod_{i=1}^n (1 - x_i z) \prod_{i=1}^{m+1} \frac{1}{1 - x_i z},$$

which is 0 (since the right-hand side is a polynomial in  $z$  of degree  $n - m - 1$ ). Therefore,  $C$  is the identity matrix, and  $A$  and  $B$  are inverses.  $\square$

We return to our uniform sequence of geometric lattices  $L_0, L_1, \dots$ . Set

$$a_n := \#\{\text{atoms } s \in L_n\} - \#\{\text{atoms } s \in L_{n-1}\},$$

for  $n = 1, 2, \dots$ .

**Corollary 3.10.** *Suppose that all the geometric lattices  $L_n$  in our uniform sequence are supersolvable. Then, their Whitney numbers of the second and first kind are*

$$\begin{aligned} V(i, j) &= h_{i-j}(a_1, \dots, a_{j+1}); \\ v(i, j) &= (-1)^{i-j} e_{i-j}(a_1, \dots, a_i), \end{aligned}$$

for  $0 \leq j \leq i$ . In particular,  $\chi^*(L_n; x) = (1 - a_1x)(1 - a_2x) \cdots (1 - a_nx)$  for all  $n$ .

*Proof.* Fix  $n \geq 1$ . Our first goal is to show  $\chi^*(L_n; x) = (1 - a_1x)(1 - a_2x) \cdots (1 - a_nx)$ . So let  $\hat{0} = x_0 < x_1 < \cdots < x_n = \hat{1}$  be a maximal chain of modular elements in  $L_n$ . Consider the modular coatom  $x_{n-1}$ . Because  $\hat{1}$  is the join of the atoms in  $L_n$  (since  $L_n$  is atomic), there must be some atom  $s \in L_n$  with  $s \not\leq x_{n-1}$ . This  $s$  is a complement of  $x_{n-1}$ , i.e.,  $s \vee x_{n-1} = \hat{1}$  and  $s \wedge x_{n-1} = \hat{0}$ . Since  $x_{n-1}$  is modular, there is thus a canonical isomorphism from  $[s, \hat{1}]$  to  $[0, x_{n-1}]$  given by  $x \mapsto x \wedge x_{n-1}$  (see [1, §4.2]). But, by assumption,  $[s, \hat{1}]$  is isomorphic to  $L_{n-1}$ , so  $[0, x_{n-1}]$  is isomorphic to  $L_{n-1}$  as well. Then by induction, each  $[0, x_i]$  is isomorphic to  $L_i$ . Thus,  $\#\{\text{atoms } s \in L_n : s \leq x_i, s \not\leq x_{i-1}\} = a_i$ . So by Theorem 3.2, we indeed have  $\chi^*(L_n; x) = (1 - a_1x)(1 - a_2x) \cdots (1 - a_nx)$ .

The previous paragraph tells us that  $v(i, j) = (-1)^{i-j} e_{i-j}(a_1, \dots, a_i)$  for all  $i, j$ . We know from Theorem 3.7 that the matrices  $[V(i, j)]$  and  $[v(i, j)]$  are inverses. But we also know from Proposition 3.9 that the inverse of the matrix  $[v(i, j)]$  is the matrix  $[h_{i-j}(a_1, \dots, a_{j+1})]$ . We conclude  $V(i, j) = h_{i-j}(a_1, \dots, a_{j+1})$  for all  $i, j$ .  $\square$

**Example 3.11.** Let us continue our running example with  $L_n = \Pi_{n+1}$ . As we saw earlier, these partition lattices are supersolvable and we have  $a_n = n$  in this case. So Corollary 3.10 tells us that

$$\begin{aligned} S(n, k) &= h_{n-k}(1, 2, \dots, k); \\ s(n, k) &= (-1)^{n-k} e_{n-k}(1, 2, \dots, n-1). \end{aligned}$$

These are classical formulas for the Stirling numbers.

**3.3. Uniform sequences of supersolvable geometric lattices.** Continue to fix a uniform sequence of geometric lattices  $L_0, L_1, \dots$ . Observe that, by induction, the interval  $[x, \hat{1}] \subseteq L_n$  is isomorphic to  $L_{n-\rho(x)}$  for all  $x \in L_n$ . So, in a sense,  $L_n$  is as close to being upho as a finite graded lattice can be: the principal order filters for all elements of the same rank are isomorphic. It is therefore reasonable to try to build an upho lattice by taking a limit of the  $L_n$  in some way. This is indeed what we will do, but to do it correctly requires some technical precision.

As we have already hinted, in order to take a limit of the  $L_n$  we will need to combine the notion of uniformity with that of supersolvability. But we will also need to make sure that the way our lattices are supersolvable is compatible with the way they form a uniform sequence. We saw in the proof of Corollary 3.10 that when

the  $L_n$  are supersolvable there are many isomorphic copies of  $L_{n-1}$  sitting inside of  $L_n$ : for each atom  $s \in L_n$ , the “upper” interval  $[s, \hat{1}_{L_n}]$  is isomorphic to  $L_{n-1}$ , and also for each modular coatom  $t \in L_n$ , the “lower” interval  $[\hat{0}_{L_n}, t]$  is isomorphic to  $L_{n-1}$ . We will need to *fix* all of these isomorphisms and make sure they are compatible with one another.

So, abusing terminology, by a *uniform sequence of supersolvable geometric lattices* we will mean a uniform sequence of geometric lattices  $L_0, L_1, \dots$  together with (for each  $n = 0, 1, \dots$ ):

- an isomorphism  $\theta_s: [s, \hat{1}_{L_n}] \rightarrow L_{n-1}$  for each atom  $s \in L_n$ ;
- an *embedding* (isomorphism onto its image)  $\iota_n: L_n \rightarrow L_{n+1}$ ,

satisfying (for each  $n = 0, 1, \dots$ ):

- $(\iota_{n-1} \circ \theta_s)(x) = (\theta_{\iota_n(s)} \circ \iota_n)(x)$  for each atom  $s \in L_n$  and all  $x \in [s, \hat{1}_{L_n}]$ ;
- the image of  $\iota_n$  is  $[\hat{0}_{L_{n+1}}, t_n]$ , where  $t_n \in L_{n+1}$  is a modular coatom.

The requirement that the image of  $\iota_n$  is  $[\hat{0}_{L_{n+1}}, t_n]$  implies in particular that all the embeddings  $\iota_n$  are *rank preserving*, i.e.,  $\rho(\iota_n(x)) = \rho(x)$  for all  $x \in L_n$ .

**Remark 3.12.** Notice how in the definition of uniform sequence we required the “upper” intervals  $[s, \hat{1}_{L_n}]$  to be isomorphic to  $L_{n-1}$  for *all* atoms  $s \in L_n$ , whereas here we only require that there be *some* (modular) coatom  $t_{n-1} \in L_n$  for which the “lower” interval  $[\hat{0}_{L_n}, t_{n-1}]$  is isomorphic to  $L_{n-1}$ . This is a crucial distinction!

**Example 3.13.** Let us continue to examine the case  $L_n = \Pi_{n+1}$ . We can upgrade this sequence to a uniform sequence of supersolvable geometric lattices by defining embeddings  $\iota_n: \Pi_{n+1} \rightarrow \Pi_{n+2}$  and isomorphisms  $\theta_s: [s, \hat{1}_{\Pi_{n+1}}] \rightarrow \Pi_n$  as follows. First of all, we set  $\iota_n(\pi) := \pi \cup \{n+2\}$  for all  $\pi \in \Pi_{n+1}$ . Next, note that any atom  $s \in \Pi_{n+1}$  has a unique non-singleton block, of the form  $\{i, j\}$  for  $1 \leq i < j \leq n+1$ . Let us denote this atom by  $s_{i,j}$ . Then for  $\pi = \{B_1, \dots, B_m\} \in [s_{i,j}, \hat{1}_{\Pi_{n+1}}]$  we set  $\theta_{s_{i,j}}(\pi) := \bigcup_{\ell=1}^m \{f(k) : k \in B_\ell \setminus \{j\}\}$ , where  $f(k)$  is  $k$  if  $k < j$  and  $k-1$  if  $k > j$ . In other words, to obtain  $\theta_{s_{i,j}}(\pi)$  from  $\pi$ , we delete  $j$  from whichever block it appears in (necessarily together with  $i$ ) and then re-index by subtracting one from all numbers greater than  $j$ . It is straightforward to verify that these  $\iota_n$  and  $\theta_s$  satisfy the requirements listed above.

From now on in this section, let us assume moreover that our sequence  $L_0, L_1, \dots$  is a uniform sequence of *supersolvable* geometric lattices. So we now also have fixed embeddings  $\iota_n: L_n \rightarrow L_{n+1}$ , and isomorphisms  $\theta_s: [s, \hat{1}_{L_n}] \rightarrow L_{n-1}$  for each atom  $s \in L_n$ . First of all, let us justify our terminology by explaining how this additional structure indeed forces the  $L_n$  to be supersolvable geometric lattices.

**Proposition 3.14.** *For  $n = 0, 1, \dots$ , the geometric lattice  $L_n$  is supersolvable.*

*Proof.* Let  $x_i := \iota_{n-1} \circ \dots \circ \iota_{i+1}(t_i) \in L_n$  for  $i = 0, 1, \dots, n-1$ , and  $x_n := \hat{1}_{L_n}$ , where  $t_m \in L_{m+1}$  is the distinguished modular coatom for which the image of  $\iota_m$  is  $[\hat{0}_{L_{m+1}}, t_m]$ . Since the  $t_m$  are modular, and the embeddings  $\iota_m$  are rank preserving, repeated application of Proposition 3.1 shows that  $x_0 \leq x_1 \leq \dots \leq x_n \in L_n$  is indeed a maximal chain of modular elements.  $\square$



Henceforth, each supersolvable geometric lattice  $L_n$  comes together with the maximal chain of modular elements  $x_0 < x_1 < \dots < x_n$  from the proof of Proposition 3.14. This lets us, for instance, speak of  $\nu: L_n \rightarrow \mathbb{N}$  and  $L_n^{(k)}$ , as defined in Section 3.1. Notice that in addition to preserving rank, the embeddings  $\iota_n$  preserve the  $\nu$  statistic, i.e.,  $\nu(\iota_n(x)) = \nu(x)$  for all  $x \in L_n$ . Therefore, the embeddings  $\iota_n$  restrict to embeddings  $\iota_n: L_n^{(k)} \rightarrow L_{n+1}^{(k)}$  for any  $k \geq 1$ .

**3.4. Limits of uniform sequences of supersolvable geometric lattices.** Continue to fix a uniform sequence of supersolvable geometric lattices  $L_0, L_1, \dots$ . We will now take the limit of this sequence, which we will denote by  $\mathcal{L}_\infty$ . Since we have distinguished embeddings  $\iota_n: L_n \rightarrow L_{n+1}$ , it makes sense to define  $\mathcal{L}_\infty := \bigcup_{n=1}^\infty L_n$ .

More precisely, we define  $\mathcal{L}_\infty := \varinjlim L_n$ , the direct limit of the directed system formed by the  $L_n$  together with the  $\iota_n$ . That is, we let  $\mathcal{L}_\infty := \bigsqcup_{n=1}^\infty L_n / \sim$ , the disjoint union of all the finite lattices  $L_n$  modulo the equivalence relation  $\sim$  generated by  $x \sim \iota_n(x)$  for all  $x \in L_n$  and all  $n = 0, 1, \dots$ . Denoting the equivalence class of an element  $x \in \bigsqcup_{n=1}^\infty L_n$  by  $[x]$ , the partial order on  $\mathcal{L}_\infty$  is  $[x] \leq [y]$  if  $x' \leq y'$  in some  $L_n$  for some  $x' \in [x], y' \in [y]$ .

Because the  $L_n$  are graded and the embeddings  $\iota_n$  are rank preserving, their limit  $\mathcal{L}_\infty$  is  $\mathbb{N}$ -graded. Similarly, since each  $L_n$  is a lattice, their limit  $\mathcal{L}_\infty$  is a lattice. And the uniformity of the sequence  $L_n$  can be used to show that every principal order filter in  $\mathcal{L}_\infty$  is isomorphic to  $\mathcal{L}_\infty$ . But  $\mathcal{L}_\infty$  is *not* finite type  $\mathbb{N}$ -graded: for instance, it has infinitely many atoms. To summarize, the limit poset  $\mathcal{L}_\infty$  is an upho lattice, except for the fact that it is not finite type  $\mathbb{N}$ -graded.

To resolve this final wrinkle, we need to make a “thinner” poset out of the “wide” poset  $\mathcal{L}_\infty$ . This is where the trimming procedure described in Section 3.1 comes into play. So, define  $\nu: \mathcal{L}_\infty \rightarrow \mathbb{N}$  by  $\nu([x]) := \min\{n: x' \in L_n \text{ for some } x' \in [x]\}$  for all  $[x] \in \mathcal{L}_\infty$ . And then for any positive integer  $k \geq 1$ , define

$$\mathcal{L}_\infty^{(k)} := \{[x] \in \mathcal{L}_\infty: \nu([x]) - \rho([x]) < k\}.$$

Since  $\rho([x]) = \rho(x)$  and  $\nu([x]) = \nu(x)$  for any  $[x] \in \mathcal{L}_\infty$ , we equivalently have that  $\mathcal{L}_\infty^{(k)} = \bigcup_{n=1}^\infty L_n^{(k)}$ , the direct limit of the trimmed finite lattices  $L_n^{(k)}$  with respect to the embeddings  $\iota_n: L_n^{(k)} \rightarrow L_{n+1}^{(k)}$ . Yet another way to think of  $\mathcal{L}_\infty^{(k)}$  is that it consists of all the elements of rank  $n - k + 1$  in each  $L_n$ , for  $n \geq k - 1$ . It is this  $\mathcal{L}_\infty^{(k)}$  which is a proper (i.e., finite type  $\mathbb{N}$ -graded) upho lattice.

**Example 3.15.** We continue with the example of  $L_n = \Pi_{n+1}$ . Then  $\mathcal{L}_\infty$  consists of all partitions of the set  $\{1, 2, \dots\}$  for which all but finitely many blocks are singletons, partially ordered by refinement. This poset  $\mathcal{L}_\infty$  is  $\mathbb{N}$ -graded: we have  $\rho(\pi) = \sum_{B \in \pi} (\#B - 1)$ . But  $\mathcal{L}_\infty$  is *not* finite type  $\mathbb{N}$ -graded. Meanwhile,  $\mathcal{L}_\infty^{(k)}$  can be viewed as the collection of all partitions of a set of the form  $[n] = \{1, 2, \dots, n\}$  (for some  $n \geq k$ ) into  $k$  blocks. The partitions in  $\mathcal{L}_\infty^{(k)}$  are not all partitions of the same set, but the partial order can be described in the same way: for  $\pi, \pi' \in \mathcal{L}_\infty^{(k)}$ , we have  $\pi \leq \pi'$  if for every  $B \in \pi$  there exists a  $B' \in \pi'$  with  $B \subseteq B'$ . And now  $\mathcal{L}_\infty^{(k)}$

is finite type  $\mathbb{N}$ -graded: we have  $\rho(\pi) = n - k$  if  $\pi$  is a partition of  $[n]$ , so there are only finitely many elements of each rank. Figure 1 depicts  $\mathcal{L}_\infty^{(2)}$  for this example.

The following theorem is the main result of this section. (In the introduction, it was stated, less precisely, as Theorem 1.3.)

**Theorem 3.16.** *For any  $k \geq 1$ ,  $\mathcal{L}_\infty^{(k)}$  is an upho lattice whose core is  $L_k$ .*

*Proof.* Let us first explain why  $\mathcal{L}_\infty^{(k)}$  is a finite type  $\mathbb{N}$ -graded lattice. As mentioned above,  $\mathcal{L}_\infty^{(k)} = \bigcup_{n=1}^\infty L_n^{(k)}$ . We know from Lemma 3.5 that the  $L_n^{(k)}$  are graded posets, and the embeddings  $\iota_n$  are rank preserving, so the limit  $\mathcal{L}_\infty^{(k)}$  is  $\mathbb{N}$ -graded. Similarly, we know from Lemma 3.5 that the  $L_n^{(k)}$  are lattices, so the limit  $\mathcal{L}_\infty^{(k)}$  is a lattice as well. Finally, as mentioned, for each  $n \geq k - 1$ , the elements of rank  $n - k + 1$  in  $\mathcal{L}_\infty^{(k)}$  are precisely  $[x]$  for  $x \in L_n$  with  $\rho(x) = n - k + 1$ . Hence, there are only finitely many elements of each rank in  $\mathcal{L}_\infty^{(k)}$ , i.e.,  $\mathcal{L}_\infty^{(k)}$  is finite type  $\mathbb{N}$ -graded.

Next, let us explain why  $\mathcal{L}_\infty^{(k)}$  is upho. Actually, it is convenient to first work with  $\mathcal{L}_\infty$ . We will show that every principal order filter in  $\mathcal{L}_\infty$  is isomorphic to  $\mathcal{L}_\infty$ . It clearly suffices to do this for principal order filters  $V_{[s]} \subseteq \mathcal{L}_\infty$  corresponding to atoms  $[s] \in \mathcal{L}_\infty$ . So let  $[s] \in \mathcal{L}_\infty$  be an atom. Recall the distinguished modular coatoms  $t_n \in L_{n+1}$  for which  $\iota_n(L_n) = [\hat{0}_{L_{n+1}}, t_n]$ . Let  $n_0 := \nu([s])$  and notice that  $n_0$  is also the smallest  $n_0$  for which  $[s] \leq [t_{n_0}]$ . We will define a series of isomorphisms  $\eta_n: [[s], [t_n]] \rightarrow [\hat{0}, [t_{n-1}]]$ , for  $n \geq n_0$ , with isomorphism  $\eta: V_{[s]} \rightarrow \mathcal{L}_\infty$  then obtained as the limit of  $\eta_n$ .

To define  $\eta_n$ , assume we have chosen the representative  $s$  of the equivalence class  $[s]$  so that  $s \in L_n$ . Then the elements of  $[[s], [t_n]]$  are  $[x]$  for  $x \in [s, \hat{1}_{L_n}]$ . By supposition, we have an isomorphism  $\theta_s: [s, \hat{1}_{L_n}] \rightarrow L_{n-1}$ . But also, the elements of  $[\hat{0}, [t_{n-1}]]$  are  $[x]$  for  $x \in L_{n-1}$ . The isomorphism  $\eta_n: [[s], [t_n]] \rightarrow [\hat{0}, [t_{n-1}]]$  is thus defined by composing  $\theta_s$  with the identifications of  $[[s], [t_n]]$  and  $[s, \hat{1}_{L_n}]$ , and of  $L_{n-1}$  and  $[\hat{0}, [t_{n-1}]]$ . Crucially, the compatibility requirement  $\iota_{n-1} \circ \theta_s = \theta_{\iota_n(s)} \circ \iota_n$  implies that the restriction of  $\eta_{n+1}$  to the domain of  $\eta_n$  agrees with  $\eta_n$ .

So we get a sequence of isomorphisms  $\eta_n: [[s], [t_n]] \rightarrow [\hat{0}, [t_{n-1}]]$ , for  $n \geq n_0$ , which have the property that the restriction of  $\eta_n$  to the domain of  $\eta_m$  agrees with  $\eta_m$  for all  $m \leq n$ . Moreover, each  $[x] \in V_{[s]}$  (respectively,  $[x] \in \mathcal{L}_\infty$ ) belongs to  $[[s], [t_n]]$  (resp.,  $[\hat{0}, [t_{n-1}]]$ ) for sufficiently large  $n$ . It therefore makes sense to define the isomorphism  $\eta: V_{[s]} \rightarrow \mathcal{L}_\infty$  by  $\eta := \bigcup_{n=n_0}^\infty \eta_n$ , i.e.,  $\eta([x]) := \eta_n([x])$  if  $[x]$  belongs to the domain of  $\eta_n$ .

Now we return to  $\mathcal{L}_\infty^{(k)}$ . We want to show that every principal order filter in  $\mathcal{L}_\infty^{(k)}$  is isomorphic to  $\mathcal{L}_\infty^{(k)}$ , and again it is sufficient to only consider the principal order filters for atoms. So let  $[s] \in \mathcal{L}_\infty^{(k)}$  be an atom, and consider the corresponding principal order filter, which for clarity we denote by  $V_{[s]}^{(k)} \subseteq \mathcal{L}_\infty^{(k)}$ . We claim that the isomorphism  $\eta: V_{[s]} \rightarrow \mathcal{L}_\infty$  above restricts to an isomorphism  $\eta: V_{[s]}^{(k)} \rightarrow \mathcal{L}_\infty^{(k)}$ . To see this, observe that, for each  $n \geq k - 1$ , the elements of rank  $n - k + 1$  in  $V_{[s]}^{(k)}$  are the elements of rank  $n - k + 1$  in  $[[s], [t_{n+1}]]$ . By construction,  $\eta_{n+1}$  maps these

elements to the elements of rank  $n - k + 1$  in  $[\hat{0}, [t_n]]$ , which are precisely the elements of rank  $n - k + 1$  in  $\mathcal{L}_\infty^{(k)}$ . So indeed,  $\eta$  restricts to an isomorphism  $\eta: V_{[s]}^{(k)} \rightarrow \mathcal{L}_\infty^{(k)}$ .

Finally, to see why the core of  $\mathcal{L}_\infty^{(k)}$  is  $L_k$ , notice that the atoms of  $\mathcal{L}_\infty^{(k)}$  are  $[s]$  for atoms  $s \in L_k$ . Joins in  $\mathcal{L}_\infty^{(k)}$  are joins of the representative elements in the  $L_n$ , so the join of the atoms of  $\mathcal{L}_\infty^{(k)}$  is  $[t_k]$ , and its core is  $[\hat{0}, [t_k]]$ , i.e., a copy of  $L_k$ .  $\square$

**Corollary 3.17.** *For any  $k \geq 1$ , we have*

$$F(\mathcal{L}_\infty^{(k)}; x) = \frac{1}{(1 - a_1 x)(1 - a_2 x) \cdots (1 - a_k x)}.$$

*Proof.* We know that  $\chi^*(L_k; x) = (1 - a_1 x)(1 - a_2 x) \cdots (1 - a_k x)$  (see Corollary 3.10). The corollary then follows from Theorem 3.16 and Corollary 2.12.  $\square$

**Remark 3.18.** I thank David Speyer and Gjergji Zaimi for explaining the following to me in answers to a question I posted to MathOverflow [9]. Suppose that  $A = [a_{ij}]_{0 \leq i, j \leq \infty}$  and  $B = [b_{ij}]_{0 \leq i, j \leq \infty}$  are two infinite, lower unitriangular matrices satisfying:

- $A$  and  $B$  are inverses;
- $\sum_{i=k}^{\infty} a_{i,k} x^{i-k} = \left( \sum_{i=0}^{k+1} b_{k+1, k+1-i} x^i \right)^{-1}$  for all  $k \geq 0$ .

Let  $a_k := a_{k, k-1} - a_{k-1, k-2}$  for all  $k \geq 1$  (where by convention  $a_{0, -1} := 0$ ). Then the entries of  $A$  and  $B$  are determined by this sequence  $a_1, a_2, \dots$ . Specifically, we must have that

$$\begin{aligned} a_{i,j} &= h_{i-j}(a_1, \dots, a_{j+1}); \\ b_{i,j} &= (-1)^{i-j} e_{i-j}(a_1, \dots, a_i). \end{aligned}$$

Because of this, we could alternatively have deduced Corollary 3.17 by combining Theorem 3.16 and Corollary 2.12 with Dowling's Theorem 3.7, without any appeal to Stanley's Theorem 3.2.

### 3.5. Examples of uniform sequences of supersolvable geometric lattices.

The construction presented above in this section would not be interesting if there were not any interesting examples of uniform sequences of supersolvable geometric lattices. Fortunately, there are many interesting examples. We now review all examples that we know of.

**3.5.1. Boolean lattices.** Let  $L_n = B_n$  be the sequence of Boolean lattices.

We define the auxiliary data  $\iota_n: B_n \rightarrow B_{n+1}$  and  $\theta_s: [s, \hat{1}_{B_n}] \rightarrow B_{n-1}$  as follows. First, the embedding  $\iota_n$  comes from the inclusion of sets  $[n] \subseteq [n+1]$ . That is, we set  $\iota_n(S) := S$  for all  $S \in B_n$ . Next, noting that any atom in  $B_n$  has the form  $\{i\}$  for  $1 \leq i \leq n$ , we set  $\theta_{\{i\}}(S) := \{f(j): j \in S \setminus \{i\}\}$  for all  $S \in [\{i\}, \hat{1}_{B_n}]$ , where  $f(j)$  is  $j$  if  $j < i$  and  $j - 1$  if  $j > i$ . In other words, to obtain  $\theta_{\{i\}}(S)$  from  $S$ , we delete  $i$  and then re-index by subtracting one from all numbers greater than  $i$ .

It is straightforward to verify that this gives a uniform sequence of supersolvable geometric lattices. Here  $a_n = 1$  for all  $n$ , and  $V(i, j) = \binom{i}{j}$  and  $v(i, j) = (-1)^{i-j} \binom{i}{j}$ .

Let us use the notation  $\mathcal{B}_\infty := \mathcal{L}_\infty$  and  $\mathcal{B}_\infty^{(k)} := \mathcal{L}_\infty^{(k)}$  for this sequence  $L_n = B_n$ . Then,  $\mathcal{B}_\infty$  is the poset of finite subsets of  $\{1, 2, \dots\}$ , ordered by inclusion. And for any  $k \geq 1$ ,  $\mathcal{B}_\infty^{(k)} = \{\text{finite } S \subseteq \{1, 2, \dots\} : \max(S) < \#S + k\}$ , ordered by inclusion. Equivalently,  $\mathcal{B}_\infty^{(k)} = \{\text{finite } S \subseteq \{1, 2, \dots\} : S \subseteq [\#S + k - 1]\}$ . Since the core of this upho lattice  $\mathcal{B}_\infty^{(k)}$  is  $B_k$ , we have  $F(\mathcal{B}_\infty^{(k)}; x) = 1/(1-x)^k$ .

Although the core of  $\mathcal{B}_\infty^{(k)}$  is  $B_k$ , we note that  $\mathcal{B}_\infty^{(k)}$  is *not* isomorphic to  $\mathbb{N}^k$  (for any  $k \geq 2$ ). Indeed, the element  $\{1, 2, \dots, n\}$  in  $\mathcal{B}_\infty^{(k)}$  covers  $n$  elements. But in  $\mathbb{N}^k$ , each element covers at most  $k$  elements. This gives the simplest example showing that an upho lattice is not determined by its core.

**3.5.2. Subspace lattices.** Now fix a prime power  $q$ , and let  $L_n = B_n(q)$  be the sequence of subspace lattices over  $\mathbb{F}_q$ . In order to define the various maps between the  $B_n(q)$ , it will be necessary to concretely represent elements of the vector spaces  $\mathbb{F}_q^n$ . Thus, let each  $\mathbb{F}_q^n$  come with a distinguished basis  $\{e_1, \dots, e_n\}$ . This gives a canonical inclusion  $\mathbb{F}_q^n \subseteq \mathbb{F}_q^{n+1}$ . It also means that each subspace  $U \in B_n(q)$  can be represented uniquely by the matrix in reduced column echelon form whose column space is  $U$ . Similarly, elements  $g \in \text{GL}(\mathbb{F}_q^n)$  of the general linear group of  $\mathbb{F}_q^n$  can also now be represented by matrices. These matrix representations are useful because we can order matrices lexically, by reading them row-by-row.

We define the auxiliary data  $\iota_n : B_n(q) \rightarrow B_{n+1}(q)$  and  $\theta_s : [s, \hat{1}_{B_n(q)}] \rightarrow B_{n-1}(q)$  as follows. First,  $\iota_n$  comes from the inclusion of vector spaces  $\mathbb{F}_q^n \subseteq \mathbb{F}_q^{n+1}$ . That is, to get the representing matrix of  $\iota_n(U)$  from that of  $U \in B_n(q)$ , we append a row of zeros. Next, noting that any atom (1-dimensional subspace)  $S \in B_n(q)$  has several complementary  $(n-1)$ -dimensional subspaces, we choose the complement  $T \in B_n(q)$  of  $S$  whose representing matrix is first in lexical order. Similarly, there are several  $g \in \text{GL}(\mathbb{F}_q^n)$  with  $g \cdot T = \text{Span}\{e_1, \dots, e_{n-1}\}$ , so we choose the  $g$  whose representing matrix is first in lexical order. Then, set  $\theta_s(U) := \iota_{n-1}^{-1}(g(T \cap U))$  for  $U \in [S, \hat{1}_{B_n(q)}]$ .

It is again an straightforward check that we get a uniform sequence of supersolvable geometric lattices. Here  $a_n = q^n$ ,  $V(i, j) = \begin{bmatrix} i \\ j \end{bmatrix}_q$  and  $v(i, j) = (-1)^{i-j} q^{\binom{i-j}{2}} \begin{bmatrix} i \\ j \end{bmatrix}_q$ .

We use notation  $\mathcal{B}_\infty(q) := \mathcal{L}_\infty$  and  $\mathcal{B}_\infty^{(k)}(q) := \mathcal{L}_\infty^{(k)}$  for this sequence  $L_n = B_n(q)$ . Let  $\mathbb{F}_q^\infty$  denote the (infinite-dimensional)  $\mathbb{F}_q$ -vector space with basis  $\{e_1, e_2, \dots\}$ . Then,  $\mathcal{B}_\infty(q)$  is the poset of finite-dimensional subspaces of  $\mathbb{F}_q^\infty$ , ordered by inclusion. And  $\mathcal{B}_\infty^{(k)}(q) = \{\text{finite-dimensional } U \subseteq \mathbb{F}_q^\infty : U \subseteq \text{Span}(\{e_1, e_2, \dots, e_{\dim(U)+k-1}\})\}$  for any  $k \geq 1$ . We have  $F(\mathcal{B}_\infty^{(k)}(q); x) = 1/((1-x)(1-qx) \cdots (1-q^{n-1}x))$ , since the core of  $\mathcal{B}_\infty^{(k)}(q)$  is  $B_k(q)$ . Evidently,  $\mathcal{B}_\infty^{(k)}(q)$  is a  $q$ -analogue of  $\mathcal{B}_\infty^{(k)}$ .

**3.5.3. Partition lattices.** The partition lattices were our running example above, but for completeness we repeat everything here. It is well-known that the partition lattices  $\Pi_n$  are supersolvable geometric lattices, and taking  $L_n = \Pi_{n+1}$  gives a uniform sequence. The appropriate auxiliary data  $\iota_n$  and  $\theta_s$  were defined in Example 3.13.

As a reminder,  $\iota_n: \Pi_{n+1} \rightarrow \Pi_{n+2}$  is given by  $\iota_n(\pi) = \pi \cup \{\{n+2\}\}$ . Here  $a_n = n$ , and  $V(i, j) = S(i+1, j+1)$  and  $v(i, j) = s(i+1, j+1)$ , the Stirling numbers of the second and first kind.

Let us use the notation  $\Pi_\infty := \mathcal{L}_\infty$  and  $\Pi_\infty^{(k)} := \mathcal{L}_\infty^{(k)}$  for this sequence  $L_n = \Pi_{n+1}$ . Then,  $\Pi_\infty$  is the poset of all partitions of the set  $\{1, 2, \dots\}$  for which all but finitely many blocks are singletons, ordered by refinement. And for any  $k \geq 1$ ,  $\Pi_\infty^{(k)}$  can be identified with the collection of partitions of  $[n]$  into  $k$  blocks, for some  $n \geq k$ . The partial order on  $\Pi_\infty^{(k)}$  is still refinement in the sense that for  $\pi, \pi' \in \Pi_\infty^{(k)}$  we have  $\pi \leq \pi'$  if for every  $B \in \pi$  there exists  $B' \in \pi'$  with  $B \subseteq B'$ . We have  $F(\Pi_\infty^{(k)}; x) = 1/((1-x)(1-2x)\cdots(1-kx))$ , since the core of  $\Pi_\infty^{(k)}$  is  $\Pi_{k+1}$ . As mentioned above, Figure 1 depicts  $\Pi_\infty^{(2)}$ .

**3.5.4. Type B partition lattices.** We now describe a Type B variant of the previous example. For an integer  $i \in \mathbb{Z}$ , let us use the shorthand  $\bar{i} := -i$ , and for a subset of integers  $S \subseteq \mathbb{Z}$ , let  $\bar{S} := \{\bar{i} : i \in S\}$ . For  $n \geq 1$ , a partition  $\pi$  of the set  $[n] \cup \bar{[n]}$  is called a *Type B partition* if:

- for every block  $B \in \pi$ , we also have  $\bar{B} \in \pi$ ;
- there is at most one block  $B \in \pi$  (called the *zero block*) with  $B = \bar{B}$ .

Let the *Type B partition lattice*  $\Pi_n^B$  be the poset of Type B partitions of  $[n] \cup \bar{[n]}$ , ordered by refinement.

It is well-known that  $\Pi_n^B$  is a geometric lattice. In fact, just as  $\Pi_{n+1}$  is the lattice of flats of the Coxeter arrangement of Type  $A_n$ ,  $\Pi_n^B$  is the lattice of flats of the Coxeter arrangement of Type  $B_n$  (see [23, 13, 17]). Moreover, it is known that  $\Pi_n^B$  is a *supersolvable* geometric lattice, and taking  $L_n = \Pi_n^B$  gives a uniform sequence. (This also follows from a more general result of Dowling [6] discussed below.) Note that the rank function on  $\Pi_n^B$  is given by  $\rho(\pi) = n - k$  if  $\pi$  has  $2k$  non-zero blocks.

We define  $\iota_n: \Pi_n^B \rightarrow \Pi_{n+1}^B$  and  $\theta_s: [s, \hat{1}_{\Pi_n^B}] \rightarrow \Pi_{n-1}^B$  for this sequence as follows. First, we set  $\iota_n(\pi) := \pi \cup \{\{n+1\}, \{\bar{n}+1\}\}$  for all  $\pi \in \Pi_n^B$ . Next, consider an atom  $s \in \Pi_n^B$ . The atom  $s$  could have a single non-singleton block of the form  $\{i, \bar{i}\}$  for  $1 \leq i \leq n$ ; denote this kind of atom by  $s_i$ . Or, the atom  $s$  could have two non-singleton blocks of the form  $\{i, j\}, \{\bar{i}, \bar{j}\}$  (respectively,  $\{i, \bar{j}\}, \{\bar{i}, j\}$ ) for  $1 \leq i < j \leq n$ ; denote this kind of atom by  $s_{i,j}$  (resp.,  $s_{i,\bar{j}}$ ). In order to not be bogged down by notation, let us describe how to obtain  $\theta_s(\pi)$  from  $\pi$  in words only. In the case where  $s = s_i$ , we obtain  $\theta_s(\pi)$  from  $\pi$  by deleting  $i$  and  $\bar{i}$  from  $\pi$ , and re-indexing by decreasing by one the absolute value of numbers greater in absolute value than  $i$ . In the case where  $s = s_{i,j}$  or  $s_{i,\bar{j}}$ , we obtain  $\theta_s(\pi)$  from  $\pi$  by deleting  $j$  and  $\bar{j}$  from  $\pi$ , and re-indexing by decreasing by one the absolute value of numbers greater in absolute value than  $j$ . These are slight variations of the maps we used for the Type A partition lattices, and it is again a straightforward, albeit tedious, check that they satisfy the requirements.

For this sequence,  $a_n = 2n - 1$ , and  $V(i, j) = S_B(i, j)$  and  $v(i, j) = s_B(i, j)$ , where  $S_B(n, k)$  and  $s_B(n, k)$  are the *Type B Stirling numbers of the second and first kind*.

These Type B Stirling numbers are defined by

$$\begin{aligned} S_B(n, k) &:= h_{n-k}(1, 3, \dots, 2k+1); \\ s_B(n, k) &:= (-1)^{n-k} e_{n-k}(1, 3, \dots, 2n-1). \end{aligned}$$

(See the recent paper [14] for more about the Type B Stirling numbers.) In particular, we have that  $\chi^*(\Pi_n^B; x) = (1-x)(1-3x) \cdots (1-(2n-1)x)$ .

We use the notation  $\prod_\infty^B := \mathcal{L}_\infty$  and  $\prod_\infty^{B,(k)} := \mathcal{L}_\infty^{(k)}$  for this sequence  $L_n = \Pi_n^B$ . Let us call a partition of the set  $\mathbb{Z} \setminus \{0\}$  a Type B partition if it satisfies the same two conditions in the bulleted list above. Then  $\prod_\infty^B$  is the poset of all Type B partitions of  $\mathbb{Z} \setminus \{0\}$  where all but finitely many blocks are singletons, ordered by refinement. And for any  $k \geq 1$ , we can view  $\prod_\infty^{B,(k)}$  as the collection of Type B partitions of a set of the form  $[n] \cup \overline{[n]}$  (for  $n \geq k-1$ ) which have  $2(k-1)$  non-zero blocks. The partial order on  $\prod_\infty^{B,(k)}$  is still refinement in the sense that for  $\pi, \pi' \in \prod_\infty^{B,(k)}$  we have  $\pi \leq \pi'$  if for every  $B \in \pi$  there exists  $B' \in \pi'$  with  $B \subseteq B'$ . We have  $F(\prod_\infty^{B,(k)}; x) = 1/((1-x)(1-3x) \cdots (1-(2k-1)x))$ , since the core of  $\prod_\infty^{B,(k)}$  is  $\Pi_k^B$ .

**3.5.5. Dowling lattices.** The most sophisticated example of a uniform sequence of supersolvable geometric lattices is due to Dowling [7, 6]. The *Dowling lattice*  $Q_n(G)$  depends on the choice of a finite group  $G$ . Choosing  $G$  to be the trivial group gives  $Q_n(G) = \Pi_{n+1}$ , and choosing  $G = \mathbb{Z}/2\mathbb{Z}$  gives  $Q_n(G) = \Pi_n^B$ . In this way, the Dowling lattices recover the previous two examples as special cases. Dowling first defined  $Q_n(G)$  in [7] for  $G$  the multiplicative group of a finite field, and then in [6] for any finite group  $G$ . See also [5, §5.3] for a graphical description of  $Q_n(G)$ .

We now review the construction of Dowling lattices. Thus, fix a finite group  $G$ , say with  $m$  elements. The construction of  $Q_n(G)$  requires several technical definitions, so please bear with us.

A *partial partition* of a set  $X$  is a collection  $\pi = \{B_1, \dots, B_k\}$  of nonempty subsets of  $X$  ( $\emptyset \subsetneq B_1, \dots, B_k \subseteq X$ ) that are pairwise disjoint ( $B_i \cap B_j = \emptyset$  for  $i \neq j$ ). In other words, a partial partition of  $X$  is a partition of a subset of  $X$ . Note that we allow  $k = 0$ , i.e., the partition with no blocks. There is a canonical bijection between the partial partitions of  $[n]$  and the (usual) partitions of  $[n+1]$  which takes the partial partition  $\pi = \{B_1, \dots, B_k\}$  to  $\pi' := \pi \cup \{[n+1] \setminus \cup_{i=1}^k B_i\}$ .

A  *$G$ -labeled set* is a map  $\alpha: A \rightarrow G$  from a set  $A$  to  $G$ . We also denote such a  $G$ -labeled set by the pair  $(\alpha, A)$ . We say that two  $G$ -labeled sets  $(\alpha, A)$  and  $(\beta, B)$  are equivalent if  $A = B$  and there is  $g \in G$  such that  $\alpha(x) = g \cdot \beta(x)$  for all  $x \in A$ . We denote the equivalence class of  $(\alpha, A)$  by  $[\alpha, A]$ .

A *partial  $G$ -partition* of a set  $X$  is a collection  $\alpha = \{[\alpha_1, A_1], [\alpha_2, A_2], \dots, [\alpha_k, A_k]\}$  of equivalence classes of  $G$ -labeled sets for which the underlying sets  $\{A_1, \dots, A_k\}$  form a partial partition of  $X$ . We continue to refer to the  $A_i$  as the blocks of  $\alpha$ .

We can now define  $Q_n(G)$ . The elements of  $Q_n(G)$  are all partial  $G$ -partitions of  $[n]$ . And the partial order is: for partial  $G$ -partitions  $\alpha = \{[\alpha_1, A_1], \dots, [\alpha_k, A_k]\}$  and  $\beta = \{[\beta_1, B_1], \dots, [\beta_\ell, B_\ell]\}$ , we have  $\alpha \leq \beta$  if

- each block  $B_j$  in  $\beta$  is a union  $B_j = A_{i_1} \cup \dots \cup A_{i_r}$  of blocks  $A_{i_1}, \dots, A_{i_r}$  in  $\alpha$ ;

- for any block  $A_i$  in  $\alpha$  with  $A_i \subseteq B_j$  for some block  $B_j$  in  $\beta$ , we have that the restriction  $\beta_j|_{A_i}: A_i \rightarrow G$  of the  $G$ -labeled set  $\beta_j: B_j \rightarrow G$  to  $A_i$  is equivalent to the  $G$ -labeled set  $\alpha_i: A_i \rightarrow G$ .

For instance, the maximum element of  $Q_n(G)$  is the partial  $G$ -partition  $\alpha = \emptyset$  with no blocks. And the minimum element of  $Q_n(G)$  is  $\alpha = \{[*], \{1\}], [*, \{2\}], \dots, [*, \{n\}]\}$ , where  $*$  denotes the map to  $G$  which is constantly equal to the identity  $e \in G$ .

Dowling [6] proved that  $Q_n(G)$  is a supersolvable geometric lattice of rank  $n$ , and that taking  $L_n = Q_n(G)$  gives a uniform sequence (see also [18, Exercise 3.131]). Note that the rank function on  $Q_n(G)$  is given by  $\rho(\alpha) = n - k$  if  $\alpha$  has  $k$  blocks.

We define  $\iota_n: Q_n(G) \rightarrow Q_{n+1}(G)$  and  $\theta_s: [s, \hat{1}_{Q_n(G)}] \rightarrow Q_{n-1}(G)$  for this sequence as follows. First, we set  $\iota_n(\alpha) := \alpha \cup \{[*], \{n+1\}]\}$  for all  $\alpha \in Q_n(G)$ . Next, consider an atom  $s \in Q_n(G)$ . This atom  $s$  could have all singleton blocks, being of the form  $s = \{[*], \{1\}], \dots, [*, \{n\}]\} \setminus \{[*], i]\}$  for some  $1 \leq i \leq n$ ; denote such an atom by  $s_i$ . Or, the  $s$  could have a single non-singleton block of the form  $\{i, j\}$  for some  $1 \leq i < j \leq n$ ; denote such an atom by  $s_{i,j}$ .<sup>2</sup> We describe how to obtain  $\theta_s(\alpha)$  from  $\alpha$  in words. In the case where  $s = s_i$ , to obtain  $\theta_s(\alpha)$  from  $\alpha$  we re-index by subtracting one from all numbers greater than  $i$ . In the case where  $s = s_{i,j}$ , to obtain  $\theta_s(\alpha)$  from  $\alpha$  we delete  $j$  and re-index by subtracting one from all numbers greater than  $j$ . It is again a straightforward, albeit tedious, check that these satisfy the requirements.

For this sequence we have  $a_n = 1 + (n-1)m$ , and hence

$$\begin{aligned} V(i, j) &= h_{i-j}(1, 1+m, 1+2m, \dots, 1+jm); \\ v(i, j) &= (-1)^{i-j} e_{i-j}(1, 1+m, 1+2m, \dots, 1+(i-1)m). \end{aligned}$$

In particular,  $\chi^*(Q_n(G); x) = (1-x)(1-(1+m)x) \cdots (1-(1+(n-1)m)x)$ .

Let us use  $\mathcal{Q}_\infty(G) := \mathcal{L}_\infty$  and  $\mathcal{Q}_\infty^{(k)}(G) := \mathcal{L}_\infty^{(k)}$  for this sequence  $L_n = Q_n(G)$ . Then,  $\mathcal{Q}_\infty(G)$  is the poset of partial  $G$ -partitions  $\alpha = \{[\alpha_1, A_1], [\alpha_2, A_2], \dots\}$  of the set  $\{1, 2, \dots\}$  for which:

- all but finitely many blocks  $A_1, A_2, \dots$  are singletons;
- the union  $A_1 \cup A_2 \cup \dots$  is cofinite, i.e.,  $\{1, 2, \dots\} \setminus \cup_{i=1}^\infty A_i$  is finite.

The partial order  $\alpha \leq \beta$  for such partial  $G$ -partitions is exactly as described above.

Unfortunately, the best description of  $\mathcal{Q}_\infty^{(k)}(G)$  we have is slightly ugly. For  $k \geq 1$ , we can represent the elements of  $\mathcal{Q}_\infty^{(k)}(G)$  by pairs  $(\alpha, n)$  where  $n \in \mathbb{N}$  is a nonnegative integer and  $\alpha$  is a partial  $G$ -partition of  $[n]$  into  $k-1$  blocks. The partial order is  $(\alpha, i) \leq (\beta, j)$  if  $i \leq j$  and  $\alpha \cup \{[*], \{i+1\}], \dots, [*, \{j\}]\} \leq \beta$  according to the partial order on partial  $G$ -partitions described above. Alternatively, we can say that  $(\alpha, i) \leq (\beta, j)$  if  $i \leq j$  and  $\alpha \leq \beta'$ , where  $\beta'$  is the result of deleting  $i+1, \dots, j$  from  $\beta$ . We have  $F(\mathcal{Q}_\infty^{(k)}(G); x) = 1/((1-x)(1-(1+m)x) \cdots (1-(1+(k-1)m)x))$ , since the core of  $\mathcal{Q}_\infty^{(k)}(G)$  is  $Q_k(G)$ .

<sup>2</sup>In this notation, we suppress the choice of the  $G$ -labelling, but the  $G$ -labelling is irrelevant.

#### 4. UPHO LATTICES FROM MONOIDS

In this section we explore an algebraic source of upho lattices: monoids. Already Gao et al. [8, §5] observed that cancellative monoids yield upho posets. (In fact, the connection between enumeration in monoids, especially cancellative monoids, and Möbius function values appeared much earlier in the work of Cartier and Foata [2] on free partially commutative monoids.) But to get an upho *lattice* in this way requires a very special kind of monoid. The most significant source of monoids with the lattice property are the Garside monoids [4, 3], whose theory and main examples we review below. We note that the upho lattices we obtain in this section are quite different from the ones we saw above. For instance, whereas all the characteristic polynomials in the previous section factored into linear factors over the integers, that will not be the case for the lattices in this section.

##### 4.1. Monoid basics.

##### 4.2. Upo posets from cancellative monoids.

4.2.1. *A monoid for rank two cores.*

##### 4.3. Garside monoids and Coxeter groups.

4.3.1. *The classical positive braid monoid and weak order.*

4.3.2. *The dual braid monoid and noncrossing partition lattice.*

#### 5. OBSTRUCTIONS FOR CORES OF UPHO LATTICES

##### 5.1. Characteristic polynomial obstructions.

5.1.1. *The face lattice of the octahedron.*

5.1.2. *The bond lattice of the four cycle.*

##### 5.2. Structural obstructions.

5.2.1. *The face lattices of the cross polytope and hypercube.*

5.2.2. *The lattice of flats of the uniform matroid.*

#### REFERENCES

- [1] G. Birkhoff. *Lattice theory*, volume Vol. XXV. American Mathematical Society, Providence, R.I., third edition, 1967.
- [2] P. Cartier and D. Foata. *Problèmes combinatoires de commutation et réarrangements*, volume No. 85 of *Lecture Notes in Mathematics*. Springer-Verlag, Berlin-New York, 1969.
- [3] P. Dehornoy, F. Digne, E. Godelle, D. Krammer, and J. Michel. *Foundations of Garside theory*, volume 22 of *EMS Tracts in Mathematics*. European Mathematical Society (EMS), Zürich, 2015.
- [4] P. Dehornoy and L. Paris. Gaussian groups and Garside groups, two generalisations of Artin groups. *Proc. London Math. Soc. (3)*, 79(3):569–604, 1999.



- [5] P. Doubilet, G.-C. Rota, and R. Stanley. On the foundations of combinatorial theory. VI. The idea of generating function. In *Proceedings of the Sixth Berkeley Symposium on Mathematical Statistics and Probability (Univ. California, Berkeley, Calif., 1970/1971), Vol. II: Probability theory*, pages pp 267–318. Univ. California Press, Berkeley, Calif., 1972.
- [6] T. A. Dowling. A class of geometric lattices based on finite groups. *J. Combinatorial Theory Ser. B*, 14:61–86, 1973.
- [7] T. A. Dowling. A  $q$ -analog of the partition lattice. In *A survey of combinatorial theory (Proc. Internat. Sympos., Colorado State Univ., Fort Collins, Colo., 1971)*, pages 101–115. North-Holland, Amsterdam, 1973.
- [8] Y. Gao, J. Guo, K. Seetharaman, and I. Seidel. The rank-generating functions of upho posets. *Discrete Math.*, 345(1):Paper No. 112629, 14, 2022.
- [9] S. Hopkins. Matrices of combinatorial sequences that are inverse in two ways. MathOverflow. <https://mathoverflow.net/q/418266>.
- [10] S. Hopkins. A note on Möbius functions of upho posets. *Electron. J. Combin.*, 29(2):Paper No. 2.39, 6, 2022.
- [11] S. Hopkins and J. Lewis. Upho lattices II: ways of realizing a core, 2024. In preparation.
- [12] V. Reiner. Non-crossing partitions for classical reflection groups. *Discrete Math.*, 177(1-3):195–222, 1997.
- [13] B. E. Sagan and J. P. Swanson.  $q$ -Stirling numbers in type  $B$ . *European J. Combin.*, 118:Paper No. 103899, 2024.
- [14] R. P. Stanley. Modular elements of geometric lattices. *Algebra Universalis*, 1:214–217, 1971/72.
- [15] R. P. Stanley. Supersolvable lattices. *Algebra Universalis*, 2:197–217, 1972.
- [16] R. P. Stanley. An introduction to hyperplane arrangements. In *Geometric combinatorics*, volume 13 of *IAS/Park City Math. Ser.*, pages 389–496. Amer. Math. Soc., Providence, RI, 2007.
- [17] R. P. Stanley. *Enumerative combinatorics. Volume 1*, volume 49 of *Cambridge Studies in Advanced Mathematics*. Cambridge University Press, Cambridge, second edition, 2012.
- [18] R. P. Stanley. From Stern’s triangle to upper homogeneous posets. Talk transparencies, available at <https://math.mit.edu/~rstan/transparencies/stern-ml.pdf>, 2020.
- [19] R. P. Stanley. Some linear recurrences motivated by Stern’s diatomic array. *Amer. Math. Monthly*, 127(2):99–111, 2020.
- [20] R. P. Stanley. Theorems and conjectures on some rational generating functions. *Eur. J. Combin.*, 119:Paper No. 103814, 2024.
- [21] The Sage Developers. *SageMath, the Sage Mathematics Software System (Version 10.2)*, 2023. <https://www.sagemath.org>.
- [22] T. Zaslavsky. The geometry of root systems and signed graphs. *Amer. Math. Monthly*, 88(2):88–105, 1981.

Email address: [samuelfhopkins@gmail.com](mailto:samuelfhopkins@gmail.com)

DEPARTMENT OF MATHEMATICS, HOWARD UNIVERSITY, WASHINGTON, DC 20059