

3/25

Standard Young Tableaux

The product formula for the g.f. of r.p.p.'s of shape λ is nice, but what about counting a finite combinatorial set?

DEFIN A Standard Young Tableau of shape λ is a filling of boxes of λ w/ the numbers $1, 2, \dots, n := |\lambda|$, each appearing exactly once, so that numbers are strictly increasing along rows & down columns.

eg. The 2 SYT's of shape $\lambda = (2, 2)$ are

1	2
3	4

and

1	3
2	4

Let $f^\lambda := \# \text{ SYT's of shape } \lambda$. Note that also

$$f^\lambda = [x_1, x_2, \dots, x_n] S_\lambda(x_1, x_2, \dots) \leftarrow \text{"coefficient of } x_1^{a_1} x_2^{a_2} \dots x_n^{a_n} \text{ in Schur's"}$$

Since an SYT is the same as a semi standard tableau of content $= (1, 1, 1, \dots, 1)$.

Thm ("Hook Length Formula", Frame-Robinson-Thrall, 1954)

$$f^\lambda = n! \cdot \prod_{u \in \lambda} \frac{1}{h(u)} \quad \text{for any partition } \lambda \vdash n$$

eg. w/ $\lambda = (2, 2)$, hook lengths are:

3	2
2	1

$$\text{So that } f^\lambda = 4! / (3 \cdot 2 \cdot 2 \cdot 1) = 2 \quad \checkmark$$


Note that $n! = \# \text{ ways to fill boxes of } \lambda \text{ w/ numbers } 1, 2, \dots, n \text{ (each used once)}$
w/out any requirement on order of #'s

So ... HLF has a probabilistic interpretation:

It says that the probability a random filling is an SYT is exactly $\prod_{u \in \lambda} \frac{1}{h(u)}$.

Bogus probabilistic proof of HLF:

- A filling is an SYT iff each entry is smallest among #'s in its hook:
- In a random filling, the probability that box u has entry smallest in its hook is $1/h(u)$
- So prob. random filling is SYT = $\prod_{u \in \lambda} 1/h(u)$




PROBLEM: 1st two bullets are correct, but can only take products for probabilities of independent events, and these events are very much not independent!

There is a valid probabilistic proof of HLF based on construction of a random SYT via "hook walk"

- Choose a random box u in λ to start at,
- Unless we're at a SE border box, move to another random box in the hook of u
- when we hit a SE border box, put the number n there.

Then we repeat w/ where to put $n-1, n-2, \dots$ etc. down to 1.

e.g. In



we might start at u_1 , then go to u_2 , then go to u_3 , and put $n=15$ there


Main thing to show is that this procedure really produces each SYT w/ equal probability ($= \frac{1}{n!}$)

See Sagan §7.3 for proof of this... We'll give different proof...

3/28

Instead, we will deduce HLF for SYTs from g.f. of r.p.p.'s!
 Actually, it will be easiest to explain this deduction in the more general setting of (finite) posets!


Recall a poset is a set with a partial order. We draw posets using Hasse diagrams. reflexive,
antisymmetric,
transitive

$P =$  $\quad a < b, a < c, b < d, c < d$
 (implied: $a < d$)

DEFN A linear extension of a poset P is a list P_1, P_2, \dots, P_n of all elements s.t. $P_i \leq P_j \Rightarrow i \leq j$.
 We let $\mathcal{L}(P) := \{\text{lin. ext.'s of } P\}$.

e.g. w/ P as above, $\mathcal{L}(P) = \{abcd, acbd\}$

DEFN We say P is naturally labeled if elts are $P = \{1, 2, \dots, n\}$ and have $i \leq_P j \Rightarrow i \leq j$ (as numbers). In this case, we treat $\mathcal{L}(P) \subseteq S_n$ as a set of permutations.

e.g. $P =$  $$ is nat. labeled and $\mathcal{L}(P) = \{1234, 1324\}$

RMK: Note that the identity $12 \dots n$ is always in $\mathcal{L}(P)$.

Recall that a descent of a permutation $\sigma = \sigma_1, \sigma_2, \dots, \sigma_n$ is a position $1 \leq i \leq n-1$ s.t. $\sigma_i > \sigma_{i+1}$. Set $D(\sigma) = \{\text{descents of } \sigma\}$ and recall that the major index of σ is


$$\text{maj}(\sigma) := \sum_{i \in D(\sigma)} i$$

e.g. $D(1234) = \emptyset$ and $D(1324) = \{2\}$ so that

$$\sum_{\sigma \in \mathcal{L}(P)} q^{\text{maj}(\sigma)} = 1 + q^2, \text{ where } P \text{ is nat. labeled poset as above.}$$

DEFN A P-partition (for poset P) is a function $\pi: P \rightarrow \mathbb{N}$ that is order-reversing: i.e., $p \leq q \Rightarrow \pi(p) \geq \pi(q)$.

We use $|\pi| := \sum_{p \in P} \pi(p)$ (like w/ the v.p.p.'s).

e.g. One P -partition is  \square = value of π (as opposed to label)


Thm (G.f. for P -partitions)

For P naturally labeled, with $\#P = n$

$$\sum_{\pi \text{ P-partition}} q^{|\pi|} = \frac{\sum_{\sigma \in \mathcal{L}(P)} q^{\text{maj}(\sigma)}}{(1-q)(1-q^2) \cdots (1-q^n)}$$

e.g. w/ P as before, g.f. is

$$\frac{1+q^2}{(1-q)(1-q^2)(1-q^3)(1-q^4)} = 1 + q + 3q^2 + 4q^3 + 7q^4 + \dots$$



Rmk: w/ $P = \bullet \leq \bullet \leq \dots \leq \bullet$ an n -element antichain

thm says $\sum_{\text{all } f: [n] \rightarrow \mathbb{N}} q^{|f|} = \frac{\sum_{\text{all } \sigma \in S_n} q^{\text{maj}(\sigma)}}{(1-q)(1-q^2) \cdots (1-q^n)}$

$$\frac{1}{(1-q)^n} \iff \sum_{\sigma \in S_n} q^{\text{maj}(\sigma)} = [n]_q!$$

Something we proved last semester (maj and inv ~~equidistributed~~ ^{same distr.})

In fact, proof we give will be same as last semester.

Cor for any poset P ,

$\# \mathcal{L}(P) = \lim_{q \rightarrow 1} \left(\sum_{\pi \text{ P-partition}} q^{|\pi|} \right) \cdot (1-q)(1-q^2) \cdots (1-q^n)$

Pf: mult. both sides by $(1-q)(1-q^2) \cdots (1-q^n)$ in thm above, and take limit $q \rightarrow 1$ (or just plug in $q = 1$) $\therefore \square$

3/30

Before we prove the g.f. for P -partitions then, let's explain how this corollary implies the HLF for SYTs. Basically we just need to match up the various terms. To any partition $\lambda \vdash n$, associate poset P_λ ($\#P_\lambda = n!$ where elt's are boxes, and $u \geq v \Leftrightarrow u$ northwest of v).

e.g. $\lambda = \begin{array}{|c|c|} \hline \square & \square \\ \hline \square & \square \\ \hline \end{array} \Leftrightarrow P_\lambda = \begin{array}{c} a \\ \swarrow \quad \searrow \\ b \quad c \\ \swarrow \quad \searrow \\ d \quad e \end{array}$

With this construction, r.p.p.'s of $sh = \lambda = P_\lambda$ -partitions and \exists bij. between SYTs of $sh = \lambda$ and $\mathcal{L}(P_\lambda)$:

$$T \mapsto \text{box w/ } n, \text{ box w/ } n-1, \dots, \text{box w/ } 1$$

e.g. $\begin{array}{|c|c|c|} \hline 1 & 2 & 4 \\ \hline 3 & 5 & \\ \hline \end{array} \mapsto d, e, b, c, a \in \mathcal{L}(P_\lambda)$

So by cor have for any $\lambda \vdash n$ that

$$f_\lambda^q = \# \text{ SYTs of } sh = \lambda = \# \mathcal{L}(P_\lambda) = \lim_{q \rightarrow 1} \frac{(1-q)(1-q^2) \dots (1-q^n)}{\sum_{\pi \text{ r.p.p. of } sh = \lambda} q^{|\pi|}}$$

$$= \lim_{q \rightarrow 1} \frac{(1-q)(1-q^2) \dots (1-q^n)}{\prod_{u \in \lambda} (1-q^{h(u)})} \stackrel{\text{g.f. of r.p.p.'s a.k.a. Hillman-Grassl}}{=} \stackrel{\text{L'Hôpital's rule}}{=} n! \cdot \prod_{u \in \lambda} \frac{1}{h(u)} \quad \text{proving HLF!}$$

Rmk By choosing a particular natural labeling of P_λ , can also obtain a q -analog of the HLF for SYTs this way, giving maj-g.f. of tableaux.

So now to finish everything, need to prove P -partition g.f.:

Pf (of P-partition g.f. in terms of $\mathcal{L}(P)$);

The idea (which we saw last semester!) is to break ΣP -partitions into pieces corresponding to lin. ext.'s.

Lemma Every $f: [n] \rightarrow \mathbb{N}$ has a unique $\sigma \in S_n$ such that f is σ -compatible in sense that

- $f_{\sigma_1} \geq f_{\sigma_2} \geq \dots \geq f_{\sigma_n}$
- if $i \in D(\sigma)$ then $f_{\sigma_i} > f_{\sigma_{i+1}}$

This f is a P-partition $\Leftrightarrow \sigma \in \mathcal{L}(P)$.

Pf: Write $f_{\sigma_1} = f_{\sigma_2} = \dots = f_{\sigma_a} > f_{\sigma_{a+1}} = \dots = f_{\sigma_b} > f_{\sigma_{b+1}} = \dots$

So that $\sigma_1 < \sigma_2 < \dots < \sigma_a$ and $\sigma_{a+1} < \dots < \sigma_b$ and ...

e.g. $f = (2, 0, 5, 0, 3, 3, 2, 0)$ has $f_3 > f_5 = f_6 > f_1 = f_7 > f_2 = f_4 = f_8$

So $\sigma = 3, 5, 6, 1, 7, 2, 4, 8$ is unique perm. f is compatible with. The statement about P-partition \Leftrightarrow lin. ext. is clear. \square

Thus, $\sum_{\pi \text{ P-partition}} q^{|\pi|} = \sum_{\sigma \in \mathcal{L}(P)} \sum_{f: [n] \rightarrow \mathbb{N} \text{ } \sigma\text{-compatible}} q^{|f|}$

$$= \sum_{\sigma \in \mathcal{L}(P)} \sum_{\lambda: (\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_n \geq 0)} q^{\text{maj}(\sigma) + |\lambda|}$$

$$= \sum_{\sigma \in \mathcal{L}(P)} q^{\text{maj}(\sigma)} \cdot \sum_{\lambda: \ell(\lambda) \leq n} q^{|\lambda|}$$

$$= \sum_{\sigma \in \mathcal{L}(P)} q^{\text{maj}(\sigma)} \cdot \frac{1}{(1-q)(1-q^2)\dots(1-q^n)}$$

Subtract off the smallest σ -compatible $f_0: [n] \rightarrow \mathbb{N}$

$$\begin{array}{r} 3 \ 5 \ 6 \ 1 \ 7 \ 2 \ 4 \ 8 \\ (5, 3, 3, 2, 2, 0, 0, 0) = f \\ - (2, 2, 2, 1, 1, 0, 0, 0) = f_0 \\ \hline (3, 1, 1, 1, 1, 0, 0, 0) = \lambda \end{array}$$

NOTE: $|f_0| = \text{maj}(\sigma)$

Since $i \in D(\sigma)$

\Rightarrow have to increase by one values in 1st i spots to get a ~~decrease~~ strict decrease

\square