




Barcelona School of Economics

Assignment 4

Foundations of Econometrics

Group 7



October 21, 2025

Question 1

Class slide 3(33) (Unit 3) illustrated, via simulation, the effects of collinearity. The script used to generate the sample is included in file `data33.R`.

Part (a)

- (i) Estimate the regression model included in the slide, presenting OLS estimates and the 95% confidence intervals for each parameter; Include the output in your answer.

Answer:

Table 1: Regression Results with 95% CI

	Estimate	Std..Error	Lower	Upper
(Intercept)	7.9365	1.6426	4.5863	11.2867
x2	0.5953	0.0747	0.4429	0.7476
x3	0.2996	0.5264	-0.7740	1.3733
x4	0.7818	0.5277	-0.2945	1.8582

- (ii) Using `confidenceEllipse()` function, or equivalent, draw the 95% confidence region for parameters β_3 , β_4 . Include also in the drawing the confidence intervals for each parameter.

Answer:




Figure 1: 95% Confidence Region for β_3 and β_4

- (iii) Describe what the confidence region you just drew provides.

Answer:

The confidence region we drew illustrates the joint confidence region, while the red lines show the individual confidence intervals. It represents every combination of the two parameters that falls in the 95% confidence region. It also illustrates the correlation between the two parameters, as the confidence region is elongated along a diagonal line. The negative slope provides insight into the negative correlation between the two parameters.

- (iv) Use the figure of confidence intervals and confidence region to show the difference between testing statistical significance of regressors separately or jointly, and explain why this is so relevant under the presence of collinear regressors.

Answer:

The figure illustrates that joint testing can show us when collinear variables are jointly significant, even if they are not individually significant.

Individual tests: Testing $H_0 : \beta_3 = 0$ and $H_0 : \beta_4 = 0$ separately uses the rectangular region formed by the individual 95% confidence intervals (red dashed lines). From the regression output, both intervals include zero: x_3 CI = $[-0.7740, 1.3733]$ and x_4 CI = $[-0.2945, 1.8582]$. Since the rectangle contains the origin $(0, 0)$, we fail to reject both null hypotheses individually—neither x_3 nor x_4 appears statistically significant.

Joint test: Testing $H_0 : \beta_3 = \beta_4 = 0$ jointly uses the 95% confidence ellipse. The ellipse is much smaller than the rectangle due to the negative correlation between $\hat{\beta}_3$ and $\hat{\beta}_4$. If the origin $(0, 0)$ falls outside the ellipse, we reject the joint null hypothesis, meaning x_3 and x_4 are jointly significant even though neither is individually significant.

Collinearity: When regressors are collinear, their coefficients are negatively correlated because if one increases, the other must decrease to fit the same data. This creates the tilted ellipse. The individual tests ignore this correlation and use the wider rectangular region, making them overly conservative. The joint test correctly accounts for correlation, revealing that while we cannot precisely determine which variable drives the effect, we can confidently say that together they have a significant impact on y . This demonstrates why collinearity makes individual t -tests unreliable while joint F -tests remain valid.

Part (b)

Modify the script used to estimate now the same regression with data generated from the same dgp but now using a sample of 3500 observations.

- (i) Surprised with how the estimates have changed? Rigorously justify.

Answer:

- (ii) Surprised of the change of the 95% confidence intervals? Rigorously justify.

Answer:

- (iii) Surprised of the change of the 95% confidence region for parameters β_3, β_4 ? Rigorously justify.

Answer:

- (iv) Using the variance decomposition expression for $\text{var}(\hat{\beta}_3 | X)$, or $\text{var}(\hat{\beta}_4 | X)$, discuss why increasing n can explain the changes observed. Be specific.

Answer:

Part (c)

Now, go back to the original script generating 35 observations, and modify the script so that now $x_{i3} + 2x_{i4} = 0$.

- (i) Run the script again. Include the output in your answer.

Answer:

- (ii) How many estimates did you get an estimate for β_3 ? And for β_4 ? You should be able to show, using the proper derivation, that in fact you got an infinite number of estimates for β_3 and β_4 .

Answer:

Question 2

Data file `microsoft.csv` includes monthly data from May 1986 to April 2013 on RP_{msft} (excess return of Microsoft stock), $RP_{s\&p}$ (excess return on the S\&P500 portfolio), $Dprod$ (variation of Industrial production), $Dinflation$ (change in inflation rate), $Dterm$ (change in interest rate) and $m1$ (an indicator variable that takes value 1 if t is the month of January and 0 otherwise). The following regression is set to measure the reaction of the excess return of Microsoft stocks to changes in macroeconomic variables:

$$RP_{msft,t} = \beta_1 + \beta_2 RP_{s\&p,t} + \beta_3 Dprod_t + \beta_4 Dinflation_t + \beta_5 Dterm_t + \beta_6 m1_t + \epsilon_t$$

- (a) Estimate the model above by OLS. Present the complete output (estimates, standard errors, p-values) as your answer.

Answer:

- (b) The January effect states that on average, every else equal, the returns (or excess returns) are larger in the month of January than the rest of the months. Test, at $\alpha = 1\%$, the presence of the January effect using the exact t -test statistic. Would you say the data supports the presence of this effect?

Answer:

- (c) Aside from normality, list the assumptions needed to justify the use of the t test statistic. Justify your answer.

Answer:

- (d) For the test you performed in questions (2b), you were asked to use the t test statistic. Using this test requires, among others, for disturbances to be normally distributed. One of the available tests of normality of the distribution of a given random variable is the Jarque-Bera test. Under the null hypothesis of normality, the Jarque-Bera (JB) test statistic is:

$$JB \equiv \frac{n}{6} \left[sk^2 + \frac{(kur - 3)^2}{4} \right] \stackrel{a}{\sim} \chi^2(2),$$

where sk is the sample coefficient of skewness of the variable and kur is its sample coefficient of kurtosis. Using a significance level of 1%, draw (by hand absolutely fine) the distribution of JB under H_0 and the corresponding acceptance and rejection regions. Provide an intuition for the location of the acceptance region.

Answer:

- (e) Now, we want to test for the presence of normality in our disturbances using the JB test. Ideally, to test normality of disturbances, JB test should be applied to a sample of disturbances, but, given that they are unobservable, the JB test is usually applied to our OLS residuals. Explain how, if all the assumptions regarding the dgp

for consistency of OLS estimator are met, it would be justified for $\hat{\epsilon}_t$'s to take the place of ϵ_t 's to perform the test.

Answer:

- (f) Perform the JB test on the OLS residuals. What do you conclude? Comment.

Answer:

- (g) Repeat the test performed in (2b) using the asymptotic T -test statistic. Use a 1% significance level.

Answer:

- (h) Is the use of asymptotic tests justified in this case? Rigorously argue.

Answer:

- (i) Consider the following statement: "Using the exact t test statistic leads to slightly more conservative inference, because we get larger acceptance regions and larger p -values than if we used the asymptotic version." Do you agree? Rigorously argue.

Answer: