

# Chapter 1

## Logic

**Definition 1.0.1.** Proposition is a statement that is either true or false, but not both.

### 1.1 Logical operations

**Definition 1.1.1.** Definition of  $\neg$ .

$p$	$\neg p$
T	F
F	T

**Definition 1.1.2.** Definition of  $\wedge$ .

$p$	$q$	$p \wedge q$
T	T	T
T	F	F
F	T	F
F	F	F

**Definition 1.1.3.** Definition of  $\vee$ .

$p$	$q$	$p \vee q$
T	T	T
T	F	T
F	T	T
F	F	F

**Definition 1.1.4.** Definition of  $\Longleftrightarrow$ .

$p$	$q$	$p \iff q$
T	T	T
T	F	F
F	T	F
F	F	T

**Definition 1.1.5.** Definition of  $\implies$ .

$$p \implies q \\ \iff (\neg p) \vee q$$

## 1.2 Quantifiers

**Definition 1.2.1.** Universal quantifier is denoted by  $\forall$ .

$$\forall x, P(x)$$

**Definition 1.2.2.** Existential quantifier is denoted by  $\exists$ .

$$\exists x, P(x)$$

**Axiom 1.1.**

$$\forall x, (P(x) \wedge Q(x)) \iff (\forall x, P(x)) \wedge (\forall x, Q(x))$$

**Axiom 1.2.** P does not depend on x.

$$(\forall x, P(y)) \iff P(y)$$

**Axiom 1.3.** De Morgan's law

$$\neg(\forall x, P(x)) \iff \exists x, \neg(P(x))$$

**Axiom 1.4.** De Morgan's law

$$\neg(\exists x, P(x)) \iff \forall x, \neg(P(x))$$

**Definition 1.2.3.** Uniqueness quantifier is denoted by  $!\exists$ .

$$!\exists x, P(x) \\ \iff (\exists x, P(x)) \wedge (\forall x \forall y (P(x) \wedge P(y) \implies x = y))$$

## 1.3 Proof technique

**Proposition 1.3.1.**

$$(P \iff Q) \iff ((P \implies Q) \wedge (Q \implies P))$$

**Proposition 1.3.2.**

$$((P \implies Q) \wedge (Q \implies R)) \implies (P \implies R)$$

**Proposition 1.3.3.** contrapositive

$$(P \implies Q) \iff (\neg Q \implies \neg P)$$

## 1.4 Proposition

Let  $P = P(x_1, x_2, \dots, x_n)$ . Let  $Q = Q(x_1, x_2, \dots, x_n)$ . etc

**Proposition 1.4.1.** Double negation

$$\neg(\neg P) \iff P$$

**Proposition 1.4.2.** Reflexive property of iff.

$$P \iff P$$

Proof:

$P$	$P \iff P$
T	T
F	T

**Proposition 1.4.3.** Symmetric property of iff.

$$(P \iff Q) \iff (Q \iff P)$$

**Proposition 1.4.4.** Transitive property of iff.

$$((P \iff Q) \wedge (Q \iff R)) \implies (P \iff R)$$

**Proposition 1.4.5.** De Morgan's law

$$\neg(P \wedge Q) \iff (\neg P) \vee (\neg Q)$$

**Proposition 1.4.6.** De Morgan's law

$$\neg(P \vee Q) \iff (\neg P) \wedge (\neg Q)$$

**Proposition 1.4.7.**

$$(P \wedge Q) \implies (P \iff Q)$$

**Proposition 1.4.8.**

$$(\neg P \iff \neg Q) \iff (P \iff Q)$$

**Proposition 1.4.9.**

$$(P \wedge Q) \implies P$$

**Lemma 1.4.10.**

$$(P \wedge ((Q \wedge P) \implies R)) \implies (Q \implies R)$$

# Chapter 2

## Set theory

Set theory have one primitive notion, called set, and one binary relation, called set membership, denoted by  $\in$ .

**Definition 2.0.1.** Definition of  $\notin$ .

$$\begin{aligned} A \notin B \\ \iff \neg(A \in B) \end{aligned}$$

**Definition 2.0.2.**

$$\begin{aligned} \forall x \in S, P(x) \\ \iff \forall x(x \in S \implies P(x)) \end{aligned}$$

**Definition 2.0.3.**

$$\begin{aligned} \exists x \in S, P(x) \\ \iff \exists x(x \in S \wedge P(x)) \end{aligned}$$

**Proposition 2.0.4.**

$$\neg(\forall x \in S, P(x)) \iff \exists x \in S, \neg(P(x))$$

Proof:

$\neg(\forall x \in S, P(x))$	
$\iff \neg(\forall x(x \in S \implies P(x)))$	Definition <a href="#">2.0.2</a>
$\iff \neg(\forall x(\neg(x \in S) \vee P(x)))$	Definition <a href="#">1.1.5</a>
$\iff \exists x, \neg(\neg(x \in S) \vee P(x))$	Axiom <a href="#">1.3</a>
$\iff \exists x, \neg(\neg(x \in S)) \wedge \neg(P(x))$	Proposition <a href="#">1.4.6</a>
$\iff \exists x, x \in S \wedge \neg(P(x))$	Proposition <a href="#">1.4.1</a>
$\iff \exists x \in S, \neg(P(x))$	Definition <a href="#">2.0.3</a>

**Proposition 2.0.5.**

$$\neg(\exists x \in S, P(x)) \iff \forall x \in S, \neg(P(x))$$

Proof:

$\neg(\exists x \in S, P(x))$	
$\iff \neg(\exists x(x \in S \wedge P(x)))$	Definition <a href="#">2.0.3</a>
$\iff \forall x, \neg(x \in S \wedge P(x))$	Axiom <a href="#">1.4</a>
$\iff \forall x, (\neg(x \in S)) \vee (\neg(P(x)))$	Proposition <a href="#">1.4.5</a>
$\iff \forall x, x \in S \implies \neg(P(x))$	Definition <a href="#">1.1.5</a>
$\iff \forall x \in S, \neg(P(x))$	Definition <a href="#">2.0.2</a>

## 2.1 Equality of sets

**Definition 2.1.1.** Definition of  $=$ .

$$A = B$$

$$\iff \forall x(x \in A \iff x \in B)$$

**Definition 2.1.2.** Definition of  $\neq$ .

$$A \neq B$$

$$\iff \neg(A = B)$$

**Proposition 2.1.3.** Reflexive property of equality

$$\forall x(x = x)$$

Proof:

$$\begin{aligned} \forall x( & \\ & x = x \\ \iff & \forall y(y \in x \iff y \in x) \quad \text{Definition 2.1.1} \\ \iff & \text{True} \quad \text{Proposition 1.4.2} \\ & ) \end{aligned}$$

**Proposition 2.1.4.** Symmetric property of equality

$$\forall x \forall y((x = y) \implies (y = x))$$

Proof:

$$\begin{aligned} \forall x \forall y( & \\ & x = y \\ \implies & \forall z(z \in x \iff z \in y) \quad \text{Definition 2.1.1} \\ \implies & \forall z(z \in y \iff z \in x) \quad \text{Proposition 1.4.3} \\ \implies & y = x \quad \text{Definition 2.1.1} \\ & ) \end{aligned}$$

**Proposition 2.1.5.** Transitive property of equality

$$\forall x \forall y \forall z((x = y) \wedge (y = z) \implies (x = z))$$

Proof:

$$\begin{aligned} \forall x \forall y \forall z( & \\ & (x = y) \wedge (y = z) \\ \implies & (\forall w(w \in x \iff w \in y)) \wedge (\forall w(w \in y \iff w \in z)) \quad \text{Definition 2.1.1} \\ \implies & \forall w((w \in x \iff w \in y) \wedge (w \in y \iff w \in z)) \quad \text{Axiom 1.1} \\ \implies & \forall w(w \in x \iff w \in z) \quad \text{Proposition 1.4.4} \\ \implies & x = z \quad \text{Definition 2.1.1} \\ & ) \end{aligned}$$

**Axiom 2.1.** Axiom of Substitution

$$\begin{aligned} \forall x \forall y( & \\ x = y & \implies \forall A(x \in A \iff y \in A) \\ & ) \end{aligned}$$

**Axiom 2.2.** Existence of empty set

$$\exists x \forall y (y \notin x)$$

**Proposition 2.1.6.** Uniqueness of empty set.

$$!\exists x \forall y (y \notin x)$$

Proof:

Let  $P(x) = \forall y (y \notin x)$

$$\begin{aligned} & \exists x \forall y (y \notin x) && \text{Axiom 2.2} \\ \implies & \exists x, P(x) && \text{Definition of } P(x) \end{aligned}$$

$$\begin{aligned} & \forall x \forall y ( \\ & \quad P(x) \wedge P(y) \\ \implies & (\forall z (z \notin x)) \wedge (\forall z (z \notin y)) && \text{Definition of } P(x) \\ \implies & \forall z ((z \notin x) \wedge (z \notin y)) && \text{Axiom 1.1} \\ \implies & \forall z (z \notin x \iff z \notin y) && \text{Proposition 1.4.7} \\ \implies & \forall z (\neg(z \in x) \iff \neg(z \in y)) && \text{Definition 2.0.1} \\ \implies & \forall z (z \in x \iff z \in y) && \text{Proposition 1.4.8} \\ \implies & x = y && \text{Definition 2.1.1} \\ & ) \end{aligned}$$

$$\begin{aligned} & (\exists x, P(x)) \wedge \forall x \forall y ((P(x) \wedge P(y)) \implies (x = y)) \\ \implies & !\exists x, P(x) && \text{Definition 1.2.3} \\ \implies & !\exists x \forall y (y \notin x) && \text{Definition of } P(x) \end{aligned}$$

**Definition 2.1.7.** The unique empty set is denoted by  $\emptyset$ .

$$\forall x (x \notin \emptyset)$$

Proof:



Let  $P(x) = \forall y(y \notin x)$

$\neg \exists x \forall y(y \notin x)$	Proposition 2.1.6
$\implies \neg \exists x, P(x)$	Definition of P(x)
$\implies (\exists x, P(x)) \wedge \forall x \forall y((P(x) \wedge P(y)) \implies (x = y))$	Definition 1.2.3
$\implies P(\emptyset) \wedge \forall x((P(x) \wedge P(\emptyset)) \implies (x = \emptyset))$	Definition 2.1.7
$\implies P(\emptyset)$	Proposition 1.4.9
$\implies \forall y(y \notin \emptyset)$	Definition of P(x)

**Proposition 2.1.8.** Uniqueness of  $\emptyset$

$$\forall x(\forall y(y \notin x) \implies (x = \emptyset))$$

Proof:

Let  $P(x) = \forall y(y \notin x)$

$\neg \exists x \forall y(y \notin x)$	Proposition 2.1.6
$\implies \neg \exists x, P(x)$	Definition of P(x)
$\implies (\exists x, P(x)) \wedge \forall x \forall y((P(x) \wedge P(y)) \implies (x = y))$	Definition 1.2.3
$\implies P(\emptyset) \wedge \forall x((P(x) \wedge P(\emptyset)) \implies (x = \emptyset))$	Definition 2.1.7
$\implies (\forall x, P(\emptyset)) \wedge \forall x((P(x) \wedge P(\emptyset)) \implies (x = \emptyset))$	Axiom 1.2
$\implies \forall x(P(\emptyset) \wedge ((P(x) \wedge P(\emptyset)) \implies (x = \emptyset)))$	Axiom 1.1
$\implies \forall x(P(x) \implies (x = \emptyset))$	Lemma 1.4.10
$\implies \forall x(\forall y(y \notin x) \implies (x = \emptyset))$	Definition of P(x)

**Proposition 2.1.9.** Single choice

$$\forall x((x \neq \emptyset) \implies (\exists y, y \in x))$$

Proof:

$\forall x(\forall y(y \notin x) \implies (x = \emptyset))$	
$\implies \forall x(\neg(x = \emptyset) \implies \neg(\forall y(y \notin x)))$	Proposition 1.3.3
$\implies \forall x((x \neq \emptyset) \implies \neg(\forall y(y \notin x)))$	Definition 2.1.2
$\implies \forall x((x \neq \emptyset) \implies (\exists y, \neg(y \notin x)))$	Axiom 1.3
$\implies \forall x((x \neq \emptyset) \implies (\exists y, \neg(\neg(y \in x))))$	Definition 2.0.1
$\implies \forall x((x \neq \emptyset) \implies (\exists y, y \in x))$	Proposition 1.4.1