

Search for chargino and neutralino production in final states with two same-sign leptons, jets and missing transverse momentum at $\sqrt{s} = 13$ TeV with the ATLAS detector

Samuel Lo ¹

¹The University of Hong Kong

February 25, 2019

Introduction

Introduction

Standard Model : Fundamental Particles

- Standard Model(SM) is the current mainstream theory to describe the electromagnetic force, weak force and strong force.

Standard Model of Elementary Particles

	three generations of matter (fermions)				
	I	II	III		
mass charge spin	$\approx 2.4 \text{ MeV}/c^2$ 2/3 1/2 u up	$\approx 1.275 \text{ GeV}/c^2$ 2/3 1/2 c charm	$\approx 172.44 \text{ GeV}/c^2$ 2/3 1/2 t top	0 0 1 g gluon	$\approx 125.09 \text{ GeV}/c^2$ 0 0 0 H Higgs
QUARKS	$\approx 4.8 \text{ MeV}/c^2$ -1/3 1/2 d down	$\approx 95 \text{ MeV}/c^2$ -1/3 1/2 s strange	$\approx 4.18 \text{ GeV}/c^2$ -1/3 1/2 b bottom	0 0 1 γ photon	
	$\approx 0.511 \text{ MeV}/c^2$ -1 1/2 e electron	$\approx 105.67 \text{ MeV}/c^2$ -1 1/2 μ muon	$\approx 1.7768 \text{ GeV}/c^2$ -1 1/2 τ tau	0 1 1 Z Z boson	
LEPTONS	$< 2.2 \text{ eV}/c^2$ 0 1/2 ν_e electron neutrino	$< 1.7 \text{ MeV}/c^2$ 1/2 1/2 ν_μ muon neutrino	$< 15.5 \text{ MeV}/c^2$ 1/2 1/2 ν_τ tau neutrino	$\approx 80.39 \text{ GeV}/c^2$ ± 1 1 W W boson	SCALAR BOSONS
				GAUGE BOSONS	

Figure: The “periodic” table for all fundamental particles in SM.

Standard Model : Fundamental Interaction

Standard Model Interactions (Forces Mediated by Gauge Bosons)








Figure: All allowed fundamental higgs-related Feynman vertices in SM.

Figure: All allowed fundamental Feynman vertices in SM, except higgs-related vertices.

Standard Model : Limitation

- ▶ SM cannot explain gravity.
- ▶ SM cannot explain the nature of dark matter.
- ▶ The hierarchy problem
 - ▶ Why the weak force is stronger than the gravity by 10^{24} .
 - ▶ Why the Higgs mass is much lighter than the Planck mass.
 - ▶ At very high energy scale, the Higgs boson mass is strongly sensitive to quantum corrections.

Supersymmetry

- ▶ Supersymmetry(SUSY) is an extension of the Standard Model.
- ▶ It can solve the hierarchy problem of Higgs mass.
- ▶ It can explain the nature of dark matter.

Supersymmetry : MSSM

- ▶ Minimal Supersymmetric Standard Model(MSSM) is the simplest realization of the supersymmetry.
- ▶ It predicts that each particle in the Standard Model has its own partner particle, called the superpartner.
- ▶ The spin of the superpartner will differ from the Standard Model particle by $1/2$.
- ▶ A symmetry between the fermions and bosons.




Figure: The particles in Standard Model and their corresponding superpartners and their names.

Supersymmetry : Superpartners

- ▶ In the MSSM, one more neutral Higgs field H and two more charged Higgs fields H^+ , H^- needed to be introduced.
- ▶ In SM electro-weak bosons, there are in total 4 neutral bosons: γ , Z , h and H , and 4 charged bosons: W^+ , W^- , H^+ and H^- .
- ▶ The superpartners of the 4 neutral bosons together form 4 mass eigenstates, called neutralinos: $\tilde{\chi}_1^0$, $\tilde{\chi}_2^0$, $\tilde{\chi}_3^0$ and $\tilde{\chi}_4^0$.
- ▶ The superpartners of the 4 charged bosons together form two mass eigenstates with electric charge ± 1 , called charginos: $\tilde{\chi}_1^\pm$ and $\tilde{\chi}_2^\pm$.

Type	SM particle	Symbol	Spin	R-parity	Superpartner	Symbol	Spin	R-parity
Fermions	Quark	q	$\frac{1}{2}$	+1	Squark	\tilde{q}	0	-1
	Lepton	l	$\frac{1}{2}$	+1	Slepton	\tilde{l}	0	-1
Gluon	Gluon	g	1	+1	Gluino	\tilde{g}	$\frac{1}{2}$	-1
Neutral EW Bosons	Photon	γ	1	+1	Neutralinos	$\tilde{\chi}_1^0, \tilde{\chi}_2^0, \tilde{\chi}_3^0, \tilde{\chi}_4^0$	$\frac{1}{2}$	-1
	Z Boson	Z	1	+1				
	Neutral Higgs	h, H	0	+1				
Charged EW Bosons	W Boson	W^+, W^-	1	+1	Charginos	$\tilde{\chi}_1^\pm, \tilde{\chi}_2^\pm$	$\frac{1}{2}$	-1
	Charged Higgs	H^+, H^-	0	+1				

Table: The spin and R-parity for the Standard Model particles and their superpartners.

Supersymmetry : R-parity

- ▶ The baryon number B is defined by $\frac{1}{3}(n_q - n_{\bar{q}})$, where n_q is the number of quarks and $n_{\bar{q}}$ is the number of anti-quarks.
- ▶ The lepton number L is defined by $n_l - n_{\bar{l}}$, where n_l is the number of leptons and $n_{\bar{l}}$ is the number of anti-leptons.
- ▶ In SM, $(B - L)$ is conserved. But in MSSM, it is no longer conserved.
- ▶ To keep $(B - L)$ conservation and prevent the proton decay, the R-parity P_R is introduced.

$$P_R = (-1)^{3(B-L)-2s}$$


where s is the spin.

Supersymmetry : R-parity

- ▶ All SM particles have R-parity $+1$, while all SUSY particles have R-parity -1 .
- ▶ If the R-parity is conserved, the lightest supersymmetric particle (LSP) cannot decay and is stable.
- ▶ If the LSP is electrically neutral and interacts with matter only by the weak interaction and gravity, it could be a candidate for dark matter, for example the lightest neutralinos $\tilde{\chi}_1^0$.
- ▶ In this thesis, the R-parity is assumed to be conserved, and the lightest neutralino $\tilde{\chi}_1^0$ is assumed to be the LSP.
- ▶ Due to the conservation of R-parity, the supersymmetric particles can only be pair-produced, and will eventually decay into SM particles and the lightest neutralino $\tilde{\chi}_1^0$ (i.e. LSP).

Our Signal Scenario : Motivation

- ▶ In the recent searches for the squarks (\tilde{q}) and gluinos (\tilde{g}), the masses of gluinos and the first and second generation squarks are suggested to be larger than 1 TeV, while the masses of the third generation squarks are still allowed to be below 1 TeV.
- ▶ In this case, the direct pair production of $\tilde{\chi}_1^\pm$ and $\tilde{\chi}_2^0$ may be the dominant SUSY production process at the LHC, if the masses of them are below 1 TeV.
- ▶ In this thesis, their masses are assumed to be the same, and denoted by $m_{\tilde{\chi}_1^\pm, \tilde{\chi}_2^0}$.



Our Signal Scenario : Decay Processes

- If all the slepton (\tilde{l}) are heavier than $\tilde{\chi}_1^\pm$ and $\tilde{\chi}_2^0$:
 1. $\tilde{\chi}_1^\pm$ will decay to W boson and $\tilde{\chi}_1^0$:
 $\tilde{\chi}_1^\pm \rightarrow W^\pm + \tilde{\chi}_1^0$
 2. $\tilde{\chi}_2^0$ will decay to the lightest MSSM Higgs boson h and $\tilde{\chi}_1^0$:
 $\tilde{\chi}_2^0 \rightarrow h + \tilde{\chi}_1^0$





Figure: The Feynman diagram for our Wh same-sign signal scenario.

Our Signal Scenario : Decay Processes


- ▶ The W boson will further to one lepton (electron or muon) and one neutrino with the SM branching ratio :

$$W^\pm \rightarrow \ell^\pm + \nu$$
- ▶ The Higgs boson h will eventually decay to one lepton (electron or muon), quarks (i.e. jets) and neutrino(s) by various decay modes with the SM branching ratios. (For example, $h \rightarrow W^+W^-$ and $h \rightarrow \tau^+\tau^-$)
- ▶ From now on, a lepton ℓ^\pm only refer to an electron or muon, but not τ lepton or neutrino.



Our Signal Scenario : Signal Signature in the Final State

- ▶ In this thesis, we only search for two same-sign(SS) leptons, in order to suppress the SM backgrounds.
- ▶ A large missing energy is expected, due to the undetected neutralinos $\tilde{\chi}_1^0$ and neutrinos ν .
- ▶ Each quark will eventually become a particle shower within a narrow cone, called a jet, by the process of hadronization.
- ▶ The mass difference between the two lightest neutralinos ($m_{\tilde{\chi}_2^0} - m_{\tilde{\chi}_1^0}$) should be larger than the Higgs mass (~ 125 GeV).



Our Signal Scenario : Sensitive Region

- ▶ If the mass difference ($m_{\tilde{\chi}_1^\pm, \tilde{\chi}_2^0} - m_{\tilde{\chi}_1^0}$) is slightly larger than the Higgs mass, it is called the compressed region.
- ▶ In the compressed region, one of the lepton may have low energy, due to the low momentum of the Higgs boson, and hence it may not be detected.
- ▶ In this case, there may be originally 3 leptons, but only 2 leptons are detected.
- ▶ This allows more decay modes for Higgs boson, which produce two leptons. (For example, $h \rightarrow ZZ$)
- ▶ Hence, the signal will be more sensitive in the compressed region.