

Search for chargino and neutralino production in final states with two same-sign leptons, jets and missing transverse momentum at $\sqrt{s} = 13$ TeV with the ATLAS detector

Samuel Lo ¹

¹The University of Hong Kong

March 7, 2019

Introduction

Experimental Setup

Charge Flip Background

Fake Lepton Background

Signal Region

Validation Region

Results

Backup

Charge Flip Background : Likelihood Method

Method for measuring real and fake efficiencies

Validation plots for fake lepton background

pre-selection plots

“N-1” plots

Validation Region : Plots for different variables

Introduction

Standard Model : Fundamental Particles

- Standard Model(SM) is the current mainstream theory to describe the electromagnetic force, weak force and strong force.




Figure: The “periodic” table for all fundamental particles in SM.

Standard Model : Fundamental Interaction




Figure: All allowed fundamental higgs-related Feynman vertices in SM.

Figure: All allowed fundamental Feynman vertices in SM, except higgs-related vertices.

Standard Model : Limitation

- ▶ SM cannot explain gravity.
- ▶ SM cannot explain the nature of dark matter.
- ▶ The hierarchy problem
 - ▶ Why the weak force is stronger than the gravity by 10^{24} .
 - ▶ Why the Higgs mass is much lighter than the Planck mass.
 - ▶ At very high energy scale, the Higgs boson mass is strongly sensitive to quantum corrections.

Supersymmetry

- ▶ Supersymmetry(SUSY) is a theoretical extension of the Standard Model.
- ▶ It is one of the most promising theory.
- ▶ It can solve the hierarchy problem of Higgs mass.
- ▶ It can explain the nature of dark matter.

Supersymmetry : MSSM

- ▶ Minimal Supersymmetric Standard Model(MSSM) is the simplest realization of the supersymmetry.
- ▶ It predicts that each particle in the Standard Model has its own partner particle, called the superpartner.
- ▶ The spin of the superpartner will differ from the Standard Model particle by 1/2.
- ▶ A symmetry between the fermions and bosons.




Figure: The particles in Standard Model and their corresponding superpartners and their names.

Supersymmetry : Superpartners

- ▶ In the MSSM, one more neutral Higgs filed H and two more charged Higgs fileds H^+ , H^- needed to be introduced.
- ▶ In SM electro-weak bosons, there are in total 4 neutral bosons: γ , Z , h and H , and 4 charged bosons: W^+ , W^- , H^+ and H^- .
- ▶ The superpartners of the 4 neutral bosons together form 4 mass eigenstates, called neutralinos: $\tilde{\chi}_1^0$, $\tilde{\chi}_2^0$, $\tilde{\chi}_3^0$ and $\tilde{\chi}_4^0$.
- ▶ The superpartners of the 4 charged bosons together form two mass eigenstates with electric charge ± 1 , called charginos: $\tilde{\chi}_1^\pm$ and $\tilde{\chi}_2^\pm$.

Type	SM particle	Symbol	Spin	R-parity	Superpartner	Symbol	Spin	R-parity
Fermions	Quark	q	$\frac{1}{2}$	+1	Squark	\tilde{q}	0	-1
	Lepton	l	$\frac{1}{2}$	+1	Slepton	\tilde{l}	0	-1
Gluon	Gluon	g	1	+1	Gluino	\tilde{g}	$\frac{1}{2}$	-1
Neutral EW Bosons	Photon	γ	1	+1	Neutralinos	$\tilde{\chi}_1^0, \tilde{\chi}_2^0, \tilde{\chi}_3^0, \tilde{\chi}_4^0$	$\frac{1}{2}$	-1
	Z Boson	Z	1	+1				
	Neutral Higgs	h, H	0	+1				
Charged EW Bosons	W Boson	W^+, W^-	1	+1	Charginos	$\tilde{\chi}_1^\pm, \tilde{\chi}_2^\pm$	$\frac{1}{2}$	-1
	Charged Higgs	H^+, H^-	0	+1				

Table: The spin and R-parity for the Standard Model particles and their superpartners.

Supersymmetry : R-parity

- ▶ The baryon number B is defined by $\frac{1}{3}(n_q - n_{\bar{q}})$, where n_q is the number of quarks and $n_{\bar{q}}$ is the number of anti-quarks.
- ▶ The lepton number L is defined by $n_l - n_{\bar{l}}$, where n_l is the number of leptons and $n_{\bar{l}}$ is the number of anti-leptons.
- ▶ In SM, $(B - L)$ is conserved. But in MSSM, it is no longer conserved.
- ▶ To keep $(B - L)$ conservation and prevent the proton decay, the R-parity P_R is introduced.

$$P_R = (-1)^{3(B-L)-2s}$$


where s is the spin.

Supersymmetry : R-parity

- ▶ All SM particles have R-parity +1, while all SUSY particles have R-parity -1.
- ▶ If the R-parity is conserved, the lightest supersymmetric particle (LSP) cannot decay and is stable.
- ▶ If the LSP is electrically neutral and interacts with matter only by the weak interaction and gravity, it could be a candidate for dark matter, for example the lightest neutralinos $\tilde{\chi}_1^0$.
- ▶ In this thesis, the R-parity is assumed to be conserved, and the lightest neutralino $\tilde{\chi}_1^0$ is assumed to be the LSP.
- ▶ Due to the conservation of R-parity, the supersymmetric particles can only be pair-produced, and will eventually decay into SM particles and the lightest neutralino $\tilde{\chi}_1^0$ (i.e. LSP).

Our Signal Scenario : Motivation

- ▶ In the recent searches for the squarks (\tilde{q}) and gluinos (\tilde{g}), the masses of gluinos and the first and second generation squarks are suggested to be larger than 1 TeV, while the masses of the third generation squarks are still allowed to be below 1 TeV.
 - ▶ In this case, the direct pair production of $\tilde{\chi}_1^\pm$ and $\tilde{\chi}_2^0$ may be the dominant SUSY production process at the LHC, if the masses of them are below 1 TeV.
 - ▶ In this thesis, their masses are assumed to be the same, and denoted by $m_{\tilde{\chi}_1^\pm, \tilde{\chi}_2^0}$.



Our Signal Scenario : Decay Processes

- If all the slepton (\tilde{l}) are heavier than $\tilde{\chi}_1^\pm$ and $\tilde{\chi}_2^0$:

1. $\tilde{\chi}_1^\pm$ will decay to W boson and $\tilde{\chi}_1^0$:

$$\tilde{\chi}_1^\pm \rightarrow W^\pm + \tilde{\chi}_1^0$$

2. $\tilde{\chi}_2^0$ will decay to the lightest MSSM Higgs boson h and $\tilde{\chi}_1^0$:

$$\tilde{\chi}_2^0 \rightarrow h + \tilde{\chi}_1^0$$





Figure: The Feynman diagram for our Wh same-sign signal scenario.

Our Signal Scenario : Decay Processes


- ▶ The W boson will further decay to one lepton (electron or muon) and one neutrino with the SM branching ratio :

$$W^\pm \rightarrow \ell^\pm + \nu$$
- ▶ The Higgs boson h will eventually decay to one lepton (electron or muon), quarks (i.e. jets) and neutrino(s) by various decay modes with the SM branching ratios. (For example, $h \rightarrow W^+W^-$ and $h \rightarrow \tau^+\tau^-$)
- ▶ From now on, a lepton ℓ^\pm only refers to an electron or muon, but not τ lepton or neutrino.



Our Signal Scenario : Signal Signature in the Final State

- ▶ In this thesis, we only search for two same-sign(SS) leptons, in order to suppress the SM backgrounds.
- ▶ A large missing energy is expected, due to the undetected neutralinos $\tilde{\chi}_1^0$ and neutrinos ν .
- ▶ Each quark will eventually become a particle shower within a narrow cone, called a jet, by the process of hadronization.
- ▶ The mass difference between the two lightest neutralinos ($m_{\tilde{\chi}_2^0} - m_{\tilde{\chi}_1^0}$) should be larger than the Higgs mass (~ 125 GeV).



Our Signal Scenario : Sensitive Region

- ▶ If the mass difference ($m_{\tilde{\chi}_1^\pm, \tilde{\chi}_2^0} - m_{\tilde{\chi}_1^0}$) is slightly larger than the Higgs mass, it is called the compressed region.
- ▶ In the compressed region, one of the lepton may have low energy, due to the low momentum of the Higgs boson, and hence it may not be detected.
- ▶ In this case, there may be originally 3 leptons, but only 2 leptons are detected.
- ▶ This allows more decay modes for Higgs boson, which produce two leptons. (For example, $h \rightarrow ZZ$)
- ▶ Hence, the signal will be more sensitive in the compressed region.


Experimental Setup

Experimental Setup : LHC

- ▶ The Large Hadron Collider (LHC) is the most powerful circular particle accelerator in the world.
- ▶ Its circumference is 27 km.
- ▶ Two beams of protons will be accelerated in opposite direction to centre-of-mass energy 13 TeV.
- ▶ They will be finally collided at the ATLAS detector.




Experimental Setup : ATLAS detector



Experimental Setup : ATLAS detector

- The ATLAS detector consists of 3 main components:
 1. Inner detector: It is a particle tracker. It measures the tracks of charged particles.
 2. Calorimeter: It measures the energy of the particle and stops the particle. It consists of electromagnetic and hadronic calorimeters.
 3. Muon spectrometer: It is a particle tracker for muons.




Challenges of Background Estimation

- ▶ Before we search for the signal, we need to understand and estimate the background first.
- ▶ SM background from “real” same-sign dilepton pair is estimated by MC
 - ▶ DiBoson: WZ, WW, ZZ
 - ▶ ttV, VVV, multitop, Higgs
- ▶ SM background from “fake” same-sign dilepton pair, due to the limitation of our detector. They are due to the wrong identification of the particle. There are two types of these “fake” backgrounds.
 - ▶ Charge flip background
 - ▶ Fake lepton background

Basic kinematic variables

- ▶ The z-axis is in the proton beam direction.
- ▶ The direction of the momentum \mathbf{p} of the particle can be specified by the azimuthal angle ϕ and the polar angle θ , as usual in the spherical coordinate system.




Basic kinematic variables

- The transverse momentum \mathbf{p}_T is the projection of \mathbf{p} onto the x-y plane. The magnitude of \mathbf{p}_T is denoted by p_T

$$p_T = \sqrt{p_x^2 + p_y^2}$$

- p_T^1 and p_T^2 are p_T of the two leptons, with $p_T^1 \geq p_T^2$. The lepton with larger p_T is called the leading lepton, while the lepton with smaller p_T is called the sub-leading lepton.




Basic kinematic variables

- The pseudorapidity η is defined by

$$\eta = -\ln \left(\tan \frac{\theta}{2} \right)$$

- The values of pseudorapidity η have reflective symmetry about the x-y plane. (Negative value for $90^\circ < \theta < 180^\circ$)
- The angle separation ΔR of two particles is defined as:

$$\Delta R = \sqrt{(\Delta\phi)^2 + (\Delta\eta)^2} = \sqrt{(\phi_2 - \phi_1)^2 + (\eta_2 - \eta_1)^2}$$



Charge Flip Background

Charge Flip Background : Sources

- ▶ The charge flip background is due to the mis-identification of the sign of the charge of a lepton (mainly electron) in the reconstruction.
- ▶ The sign of the charge is determined by the direction of the curvature of the track. The charge may be reversed, or flipped.




Figure: Bremsstrahlung and γ conversion.




Figure: Track with high velocity
Oral Presentation

Charge Flip Background : charge flip rate

- ▶ The probability that the charge of an electron is flipped, called the charge flip rate $\epsilon(p_T, |\eta|)$, depends on the p_T and $|\eta|$ of the electron.
- ▶ The charge flip rates are measured by the likelihood method described in the backup slides.
- ▶ After getting the charge flip rates, the charge flip background can be found by applying the charge flip rates on the OS events in data.
- ▶ The probability that an OS data event becomes a SS data event, which is the charge of one of the leptons is flipped, is given by

$$\begin{aligned} P(p_T^1, |\eta^1|, p_T^2, |\eta^2|) \\ &= [1 - \epsilon(p_T^1, |\eta^1|)] \cdot \epsilon(p_T^2, |\eta^2|) + [1 - \epsilon(p_T^2, |\eta^2|)] \cdot \epsilon(p_T^1, |\eta^1|) \\ &= (1 - \epsilon_1)\epsilon_2 + (1 - \epsilon_2)\epsilon_1 \end{aligned}$$

- ▶ It is called the charge flip weight. The charge flip weight is applied on all OS data events to estimate the charge flip background.

Charge Flip Background : Result for Charge Flip Rate




Figure: The measured values of the charge-flip rate $|\epsilon_i|$ in data, with total uncertainties. Oral Presentation

Charge Flip Background : MC Validation Plots




Figure: The comparison between the weighted OS events and the SS events, with different variables.

Charge Flip Background : MC Validation Plots




Figure: The comparison between the weighted OS events and the SS events, with different variables.

Fake Lepton Background

Fake Lepton Background : Sources

The source of the fake lepton is that other object (like jet and photon) is mis-identified as a lepton. They can be mainly classified into 3 types:

- ▶ Heavy-flavor fakes:
 - ▶ It comes from semi-leptonic decays of heavy-quark (b or c) hadrons in jets.
- ▶ Light-flavor fakes:
 - ▶ It comes from semi-leptonic decays of light-quark hadrons in jets.
OR
 - ▶ It is due to mis-reconstructions of jets with light-quark hadrons.
- ▶ photon conversion:
 - ▶ It comes from the pair production from a photon

Fake Lepton Background : Matrix Method

- ▶ The fake leptons is estimated by the matrix method.
- ▶ The probability that a real lepton passes the tight selection (i.e. tight lepton) is denoted by the real efficiency ϵ .
- ▶ The probability that a real lepton does not pass the tight selection (i.e. loose lepton) is denoted by $\bar{\epsilon} = 1 - \epsilon$.
- ▶ The probability that a fake lepton passes the tight selection (i.e. tight lepton) is denoted by the fake efficiency f .
- ▶ The probability that a fake lepton does not pass the tight selection (i.e. loose lepton) is denoted by $\bar{f} = 1 - f$.

$$\begin{pmatrix} N_T \\ N_L \end{pmatrix} = \begin{pmatrix} \epsilon & f \\ \bar{\epsilon} & \bar{f} \end{pmatrix} \begin{pmatrix} N_R \\ N_F \end{pmatrix}$$

Fake Lepton Background : Matrix Method

Suppose we know $\begin{pmatrix} N_T \\ N_L \end{pmatrix}$, we can find N_F by inverting the matrix.

$$\begin{pmatrix} 0 \\ N_F \end{pmatrix} = \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} N_R \\ N_F \end{pmatrix} = \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} \epsilon & f \\ \bar{\epsilon} & \bar{f} \end{pmatrix}^{-1} \begin{pmatrix} N_T \\ N_L \end{pmatrix}$$

Then, we can find the number of tight lepton due to the fake lepton,
 N'_T

$$\begin{aligned} \begin{pmatrix} N'_T \\ N'_L \end{pmatrix} &= \begin{pmatrix} \epsilon & f \\ \bar{\epsilon} & \bar{f} \end{pmatrix} \begin{pmatrix} 0 \\ N_F \end{pmatrix} \\ &= \begin{pmatrix} \epsilon & f \\ \bar{\epsilon} & \bar{f} \end{pmatrix} \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} \epsilon & f \\ \bar{\epsilon} & \bar{f} \end{pmatrix}^{-1} \begin{pmatrix} N_T \\ N_L \end{pmatrix} \end{aligned}$$

Fake Lepton Background : Matrix Method

We can generalize the one-lepton case to the two-leptons case.

$$\begin{pmatrix} N_{TT} \\ N_{TL} \\ N_{LT} \\ N_{LL} \end{pmatrix} = \begin{pmatrix} \epsilon_1 \epsilon_2 & \epsilon_1 f_2 & f_1 \epsilon_2 & f_1 f_2 \\ \epsilon_1 \bar{\epsilon}_2 & \epsilon_1 \bar{f}_2 & f_1 \bar{\epsilon}_2 & f_1 \bar{f}_2 \\ \bar{\epsilon}_1 \epsilon_2 & \bar{\epsilon}_1 f_2 & \bar{f}_1 \epsilon_2 & \bar{f}_1 f_2 \\ \bar{\epsilon}_1 \bar{\epsilon}_2 & \bar{\epsilon}_1 \bar{f}_2 & \bar{f}_1 \bar{\epsilon}_2 & \bar{f}_1 \bar{f}_2 \end{pmatrix} \begin{pmatrix} N_{RR} \\ N_{RF} \\ N_{FR} \\ N_{FF} \end{pmatrix}$$

where ϵ_1 is the probability that a leading real lepton passes the signal selection (i.e. tight lepton), ϵ_2 is the probability that a subleading real lepton passes the signal selection (i.e. tight lepton), etc.

Fake Lepton Background : Matrix Method

We can find N_{RF} , N_{FR} , N_{FF} by inverting the matrix.

$$\begin{pmatrix} 0 \\ N_{RF} \\ N_{FR} \\ N_{FF} \end{pmatrix} = \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} N_{RR} \\ N_{RF} \\ N_{FR} \\ N_{FF} \end{pmatrix}$$

$$= \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} \epsilon_1\epsilon_2 & \epsilon_1f_2 & f_1\epsilon_2 & f_1f_2 \\ \epsilon_1\bar{\epsilon}_2 & \epsilon_1\bar{f}_2 & f_1\bar{\epsilon}_2 & f_1\bar{f}_2 \\ \bar{\epsilon}_1\epsilon_2 & \bar{\epsilon}_1f_2 & \bar{f}_1\epsilon_2 & \bar{f}_1f_2 \\ \bar{\epsilon}_1\bar{\epsilon}_2 & \bar{\epsilon}_1\bar{f}_2 & \bar{f}_1\bar{\epsilon}_2 & \bar{f}_1\bar{f}_2 \end{pmatrix}^{-1} \begin{pmatrix} N_{TT} \\ N_{TL} \\ N_{LT} \\ N_{LL} \end{pmatrix}$$

Fake Lepton Background : Matrix Method

Then, we can find the number of tight-tight leptons due to the fake leptons, N'_{TT}

$$\begin{pmatrix} N'_{TT} \\ N'_{TL} \\ N'_{LT} \\ N'_{LL} \end{pmatrix} = \begin{pmatrix} \epsilon_1\epsilon_2 & \epsilon_1f_2 & f_1\epsilon_2 & f_1f_2 \\ \epsilon_1\bar{\epsilon}_2 & \epsilon_1\bar{f}_2 & f_1\bar{\epsilon}_2 & f_1\bar{f}_2 \\ \bar{\epsilon}_1\epsilon_2 & \bar{\epsilon}_1f_2 & \bar{f}_1\epsilon_2 & \bar{f}_1f_2 \\ \bar{\epsilon}_1\bar{\epsilon}_2 & \bar{\epsilon}_1\bar{f}_2 & \bar{f}_1\bar{\epsilon}_2 & \bar{f}_1\bar{f}_2 \end{pmatrix} \begin{pmatrix} 0 \\ N_{RF} \\ N_{FR} \\ N_{FF} \end{pmatrix}$$

$$= \begin{pmatrix} \epsilon_1\epsilon_2 & \epsilon_1f_2 & f_1\epsilon_2 & f_1f_2 \\ \epsilon_1\bar{\epsilon}_2 & \epsilon_1\bar{f}_2 & f_1\bar{\epsilon}_2 & f_1\bar{f}_2 \\ \bar{\epsilon}_1\epsilon_2 & \bar{\epsilon}_1f_2 & \bar{f}_1\epsilon_2 & \bar{f}_1f_2 \\ \bar{\epsilon}_1\bar{\epsilon}_2 & \bar{\epsilon}_1\bar{f}_2 & \bar{f}_1\bar{\epsilon}_2 & \bar{f}_1\bar{f}_2 \end{pmatrix} \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}$$

$$\begin{pmatrix} \epsilon_1\epsilon_2 & \epsilon_1f_2 & f_1\epsilon_2 & f_1f_2 \\ \epsilon_1\bar{\epsilon}_2 & \epsilon_1\bar{f}_2 & f_1\bar{\epsilon}_2 & f_1\bar{f}_2 \\ \bar{\epsilon}_1\epsilon_2 & \bar{\epsilon}_1f_2 & \bar{f}_1\epsilon_2 & \bar{f}_1f_2 \\ \bar{\epsilon}_1\bar{\epsilon}_2 & \bar{\epsilon}_1\bar{f}_2 & \bar{f}_1\bar{\epsilon}_2 & \bar{f}_1\bar{f}_2 \end{pmatrix}^{-1} \begin{pmatrix} N_{TT} \\ N_{TL} \\ N_{LT} \\ N_{LL} \end{pmatrix}$$

Fake Lepton Background : Results of Real Efficiencies

The method for measuring the real efficiencies is described in the backup slides.




Figure: Electron real efficiency




Figure: Muon real efficiency

Fake Lepton Background : Results of Fake Efficiencies

The method for measuring the fake efficiencies is described in the backup slides.




Figure: Electron real efficiency




Figure: Muon real efficiency

Signal Region

Signal Region: Challenges

Estimation of the signal sensitivity

- ▶ Cross section for our signal $\sim 0.1 \text{ pb}$
- ▶ Total luminosity in one year $\sim 10^4 \text{ pb}^{-1}$
- ▶ Expected number of signal events in one year $\sim 10^3$
- ▶ Time interval between each collision = 25 ns
- ▶ Total number of events in one year $\sim 10^{15}$
- ▶ The probability to have a signal event $\sim 10^{-12}$
- ▶ Conclusion: the SUSY signal is very rare.

Signal Region : Strategy

In order to extract the signal, the following strategies are used.

- ▶ A hardware trigger system are used when taking the data (online), to only store the interested events in the disk.
(40 MHz → 100 kHz → 1 kHz)
- ▶ Some discriminant variables (based on the kinematic variables) are defined, to help distinguish the signal and the background (i.e. noise).
- ▶ Two dedicated signal regions are defined based on the discriminant variables, to maximize signal sensitivity (like the signal-to-noise ratio).

Signal Region : discriminant variable

- ▶ The missing momentum \mathbf{p}^{miss} due to undetected particle is defined by

$$\mathbf{p}^{\text{miss}} = - \sum_{\text{All detected particles}} \mathbf{p}$$

- ▶ The transverse missing energy, denoted by E_T^{miss} or MET, is defined by

$$E_T^{\text{miss}} = |\mathbf{p}_T^{\text{miss}}| = \sqrt{(p_x^{\text{miss}})^2 + (p_y^{\text{miss}})^2}$$

Signal Region : discriminant variable

- ▶ n_{jets} : Number of signal jets:
- ▶ $n_{b\text{-jets}}$: Number of b -jets (jet from b quark).



$$\Delta\eta_{ll} = |\eta_1 - \eta_2|$$



$$m_{\text{eff}} = p_T^1 + p_T^2 + E_T^{\text{miss}} + \sum_{\text{signal jets}} p_T$$

- ▶ m_{ll} : The invariant mass of the 4-momentum sum of the two leptons

$$(m_{ll})^2 = (p_1 + p_2)^2$$

Signal Region : discriminant variable

- ▶ m_T : It is designed to reconstruct the mass of the W boson.

$$m_T = \sqrt{2p_T^1 E_T^{\text{miss}} (1 - \cos \Delta\phi)}$$

where $\Delta\phi$ is the azimuthal angle between the leading lepton and the missing transverse momentum.

- ▶ m_{T2} : It is designed to set a lower bound on the masses of the unseen pair of charginos $\tilde{\chi}_1^\pm$ and neutralinos $\tilde{\chi}_2^0$.

$$m_{T2} = \min_{\mathbf{q}_T} \left[\max \left(m_T(\mathbf{p}_T^1, \mathbf{q}_T), m_T(\mathbf{p}_T^2, \mathbf{p}_T^{\text{miss}} - \mathbf{q}_T) \right) \right]$$

$$m_T(\mathbf{p}_T, \mathbf{q}_T) = \sqrt{2p_T q_T (1 - \cos \Delta\phi)}$$

Signal Region : discriminant variable

- ▶ m_{lj} or m_{ljj} : It is designed to reconstruct the mass of the Higgs boson.

$$p_{\text{jet-system}} = \begin{cases} p_{\text{jet1}} & \text{for SRjet1} \\ p_{\text{jet1}} + p_{\text{jet2}} & \text{for SRjet23} \end{cases}$$

$$p_{\text{closest-lepton}} = \begin{cases} p_{\ell 1} & \text{if } \Delta R(p_{\ell 1}, p_{\text{jet-system}}) \leq \Delta R(p_{\ell 2}, p_{\text{jet-system}}) \\ p_{\ell 2} & \text{if } \Delta R(p_{\ell 1}, p_{\text{jet-system}}) > \Delta R(p_{\ell 2}, p_{\text{jet-system}}) \end{cases}$$

$$(m_{lj(j)})^2 = (p_{\text{closest-lepton}} + p_{\text{jet-system}})^2$$





Signal Region : Pre-selection

- ▶ Signal Region(SR) is a set of cuts of the discriminant variables, such that the signal sensitivity is high.
- ▶ By trying different sets of cuts, the signal region with the highest signal sensitivity can be found. This process is called the signal region optimization.
- ▶ In the SR optimization, the MC signal and background are used to study the expected sensitivity.

The following cuts are the pre-selection, before the signal region optimization.

- ▶ Exactly 2 leptons with $p_T > 25$ GeV.
- ▶ Electron: $|\eta| < 2.47$; Muon: $|\eta| < 2.4$
- ▶ The two leptons are same-sign in electric charge (SS).
- ▶ $n_{b\text{-jets}} = 0$, to reduce the SM background from top quark.
- ▶ Two signal regions: SRjet1 with $n_{\text{jets}} = 1$, SRjet23 with $n_{\text{jets}} = 2$ or 3.

Signal Region : Pre-selection plots



Signal Region : cut result

	SRjet1	SRjet23
n_{jets}	1	2 or 3
Leading lepton p_T [GeV]	≥ 25	≥ 25
Sub-leading lepton p_T [GeV]	≥ 25	≥ 25
$ \Delta\eta_{ll} $	< 1.5	-
E_T^{miss} [GeV]	≥ 100	≥ 100
m_{eff} [GeV]	≥ 260	≥ 240
m_T^{l1} [GeV]	≥ 140	≥ 120
m_{lj}/m_{ljj} [GeV]	< 180	< 130
m_{T2} [GeV]	≥ 80	≥ 70

Signal Region : Expected Yields





	Number of events	Significance
Fakes	3.295434 ± 0.819175 (24)	
WZ	2.176292 ± 0.397871 (257)	
charge flip	0.472185 ± 0.052737 (265)	
Rare	0.443536 ± 0.111301 (56)	
WW	0.165504 ± 0.023455 (67)	
$t\bar{t} + V$	0.124898 ± 0.046227 (36)	
ZZ	0.055330 ± 0.027713 (31)	
Total BG	6.733178 ± 0.920855 (736)	
(225.0,75.0)	3.33 ± 0.60 (60)	0.74
(187.5,37.5)	3.77 ± 0.95 (31)	0.86
(175.0,0.0)	4.29 ± 0.73 (36)	0.98

Table: Yields in SRjet1

	Number of events	Significance
WZ	1.848838 ± 0.273025 (319)	
Fakes	1.764693 ± 0.709215 (20)	
Rare	0.731015 ± 0.195114 (48)	
WW	0.514488 ± 0.036878 (235)	
charge flip	0.267485 ± 0.029086 (274)	
$t\bar{t} + V$	0.141679 ± 0.030772 (67)	
ZZ	0.067226 ± 0.024995 (24)	
Total BG	5.335425 ± 0.787005 (987)	
(225.0,75.0)	2.35 ± 0.34 (58)	0.57
(187.5,37.5)	4.72 ± 0.74 (47)	1.26
(175.0,0.0)	8.60 ± 1.51 (58)	2.24

Table: Yields in SRjet23

Signal Region : N-1 plots



Validation Region

Validation Region: Introduction

- ▶ The validation region(VR) is designed to ensure that the background events have a good modelling.
- ▶ Method: The backgrounds are compared with the data, to see whether they agree with each other.
- ▶ One VR is defined for each SR: VRjet1 and VRjet23.

There are three requirements for the validation regions:

- ▶ The signal contribution should be small.
- ▶ The validation region is orthogonal to the corresponding signal region.
- ▶ The background composition is similar to the corresponding signal region.

Validation Region: Method

► VRjet1

- ▶ invert the E_T^{miss} cut, to make it orthogonal to SR.
- ▶ adjust E_T^{miss} lower cut to 70 GeV, to have similar background composition.
- ▶ invert the m_{lj} cut, to reduce the signal.
- ▶ remove the m_{eff} and m_{T2} cuts, to increase statistics.

► VRjet23

- ▶ invert the m_T cut, to make it orthogonal to SR.
- ▶ adjust m_T lower cut to 65 GeV, to have similar background composition.
- ▶ invert the m_{ljj} cut, to reduce the signal.
- ▶ remove the m_{T2} cut, to increase statistics.

Cut	VRjet1	VRjet23
n_{jets}	1	[2, 3]
$\Delta\eta_{ll}$	< 1.5	—
E_T^{miss} [GeV]	[70, 100]	> 100
m_T [GeV]	> 140	[65, 120]
m_{eff} [GeV]	—	> 240
$m_{lj(j)}$ [GeV]	> 130	> 130
m_{T2} [GeV]	—	—

Validation Region: Check for Signal Contribution

The signal contribution is below 12% and 10% for VRjet1 and VRjet23 respectively.




Figure: Percentages of signal contribution are shown for the VRjet1 (left) and VRjet23 (right).


Validation Region: Check for Background Composition

The background compositions for VRjet1 and SRjet1 are similar.

ATLAS Internal
13 TeV, 36.1 fb⁻¹

VRjet1

Rare $t\bar{t}V$
 ZZ WZ
WW Charge flip
Fakes



ATLAS Internal
13 TeV, 36.1 fb⁻¹

SRjet1

Rare $t\bar{t}V$
 ZZ WZ
WW Charge flip
Fakes




Figure: Background composition for the VRjet1 (left) and SRjet1 (right).


Validation Region: Check for Background Composition

The background compositions for VRjet23 and SRjet23 are similar.

ATLAS Internal
13 TeV, 36.1 fb⁻¹

VRjet23

Rare $t\bar{t}V$
 ZZ WZ
 WW Charge flip
Fakes



ATLAS Internal
13 TeV, 36.1 fb⁻¹

SRjet23

Rare $t\bar{t}V$
 ZZ WZ
 WW Charge flip
Fakes




Figure: Background composition for the VRjet23 (left) and SRjet23 (right).

Validation Region: Yield

Process	VRjet1	VRjet23
Rare	$0.775 \pm 0.389^{+0.661}_{-0.362}$	$2.469 \pm 0.674^{+0.998}_{-0.899}$
$t\bar{t}V$	$0.039 \pm 0.013^{+0.018}_{-0.012}$	$0.959 \pm 0.082^{+0.152}_{-0.146}$
ZZ	$0.298 \pm 0.060^{+0.089}_{-0.063}$	$0.247 \pm 0.045^{+0.113}_{-0.047}$
WZ	$4.909 \pm 0.530^{+0.960}_{-0.899}$	$19.325 \pm 0.643^{+4.393}_{-4.346}$
WW	$0.801 \pm 0.051^{+0.123}_{-0.060}$	$10.477 \pm 0.176^{+0.796}_{-0.726}$
Charge flip	$1.997 \pm 0.128^{+0.260}_{-0.260}$	$2.065 \pm 0.085^{+0.166}_{-0.166}$
Fakes	$8.021 \pm 1.390^{+5.806}_{-5.806}$	$19.990 \pm 2.013^{+13.461}_{-13.461}$
Total BG	$16.839 \pm 1.545^{+5.915}_{-5.912}$	$55.534 \pm 2.228^{+14.396}_{-14.332}$
Data	17	54


Table: The uncertainties include the statistical and systematic uncertainties.

The distribution plots for different variables are in the backup slides.

Results


Results: Exclusion Limits

- ▶ Two SR are combined to calculate the exclusion limits.
- ▶ The contour line is in 95% confidence level.



Results: Exclusion Limits

- ▶ The exclusion limits for the masses of $\tilde{\chi}_1^\pm$ and $\tilde{\chi}_2^0$ are extended up to 245 GeV, and the exclusion limits for the mass of $\tilde{\chi}_1^0$ are extended up to 40 GeV.
- ▶ This means that the mass points inside the contour line are rejected with 95% confidence level.



Conclusion

- ▶ The charge flip BG was estimated by the likelihood method.
- ▶ The fake lepton BG was estimated by the matrix method.
- ▶ Two SR were found and their expected sensitivities was also calculated.
- ▶ Two VR were found and different backgrounds were validated.
- ▶ New exclusion limits were set, by combining the two SR.

Thank You

Backup

Charge Flip Background : Likelihood Method

Likelihood Method : Method

- ▶ The probability that the charge of an electron is flipped is denoted by the charge-flip rate ϵ_i , where the index i represents the dependency on the p_T and $|\eta|$ of the electron.
- ▶ The value of index i is defined by the index of the following grids in the table.

Variable	Boundary of the bins
p_T (GeV)	25, 60, 90, 130, 150, 1000
$ \eta $	0, 0.50, 1.00, 1.37, 1.52, 1.80, 2.00, 2.47

Table: Binning in p_T and $|\eta|$ for the charge-flip rate ϵ_i .

- ▶ The probability p_{ij} that an OS data event becomes a SS data event (with the leading lepton in bin i and the subleading lepton in bin j) is

$$p_{ij} = (1 - \epsilon_i)\epsilon_j + (1 - \epsilon_j)\epsilon_i$$

Likelihood Method : Definition of Control Region

A control region is defined.

- ▶ Trigger requirement
- ▶ exactly 2 signal leptons
- ▶ 2 electrons
- ▶ $25 < p_T < 1000$
- ▶ $|\eta| < 2.47$

Likelihood Method : Method

- ▶ The charge flip rate ϵ_i is measured by using the data in the control region, within the Z mass window cut of 80-100 GeV.
- ▶ The events inside the Z mass window is then subtracted by the non-Zee processes, by using the sideband technique.
- ▶ After the subtraction, the total number of event is denoted by N^{ij} , and the number of events with two same-sign leptons is denoted by N_{SS}^{ij} .
- ▶ The expected value of N_{SS}^{ij} is then given by $\lambda = N^{ij} p_{ij}$. And, if N_{SS}^{ij} is described by a Poisson distribution, then

$$P(N_{SS}^{ij}|N^{ij}, \epsilon_i, \epsilon_j) = \frac{(N^{ij} p_{ij})^{N_{SS}^{ij}} e^{-N^{ij} p_{ij}}}{N_{SS}^{ij}!}$$

Likelihood Method : Method

- ▶ Converting it to the likelihood function L and taking the negative natural log yields

$$\begin{aligned}-\ln L &= -\ln \prod_{ij} P(N_{SS}^{ij} | N^{ij}, \epsilon_i, \epsilon_j) \\&= -\sum_{ij} \left[N_{SS}^{ij} \ln(N^{ij}[\epsilon_i(1-\epsilon_j) + (1-\epsilon_i)\epsilon_j]) \right. \\&\quad \left. - N^{ij}[\epsilon_i(1-\epsilon_j) + (1-\epsilon_i)\epsilon_j] \right] + \text{constant}\end{aligned}$$

- ▶ where the summation over i and j is taken over all p_T and $|\eta|$ bins of both electrons.
- ▶ By minimizing this likelihood, the charge-flip rate can be estimated.

Likelihood Method : Background Subtraction

Both for data and MC


- ▶ Central region: 80 - 100 GeV
- ▶ Sideband region: 60 - 80 and 100 - 120
(Sideband width 20 GeV)

$$N_{Central} = N_{Central} - \frac{20}{20 + 20} N_{SB}$$

- ▶ Background subtraction is done both on N^{ij} and N_{SS}^{ij}


Likelihood Method : Result for Charge Flip Rate (statistical error only)

- ▶ Results by likelihood method from data, $\epsilon_{\text{lik,data}}$
- ▶ The error is statistical only, from likelihood method, $\sigma_{\text{lik,data}}$.



Likelihood Method : MC Truth Validation

- ▶ Results by likelihood method from MC (blue points)
- ▶ Results by MC truth (red points)




Likelihood Method : MC Truth Matching

- ▶ It is a complicated part, because we do not know the original charge of the reconstructed electron for some cases.
- ▶ The reconstructed electrons match the truth particle by the smallest ΔR ($\Delta R < 0.1$).
- ▶ No match for reconstructed electrons if no any truth particle is inside $\Delta R < 0.1$. (cannot find the original charge)
- ▶ If the truth particle is matched, the truth particle is not electron. (fake lepton, not charge flip)

Likelihood Method : MC Truth Matching

- ▶ If the truth particle is electron, the (grand)mother particle is not Z Boson. (ignore it)
- ▶ If (grand)mother particle is Z Boson, the daughter of the Z Boson is not electron. (ignore it)
- ▶ If the daughter of the Z Boson is electron, the charge of that electron is the original charge.



Likelihood Method : Systematic uncertainties due to likelihood method

For each bin, the systematic uncertainties due to likelihood method are estimated by the difference between the likelihood method and the MC truth method.

For MC,

$$\sigma_{\text{truth,MC}} = |\epsilon_{\text{lik,MC}} - \epsilon_{\text{MC truth}}|$$

For data,

$$\sigma_{\text{truth,data}} = \epsilon_{\text{lik,data}} \times \frac{\sigma_{\text{truth,MC}}}{\epsilon_{\text{lik,MC}}}$$

Likelihood Method : Systematic uncertainties due to background subtraction

Nominal:

- ▶ Central region: 80 - 100 GeV, Sideband width: 20 GeV

Variations:


- ▶ Central region: 80 - 100 GeV, Sideband width: 15 GeV
- ▶ Central region: 80 - 100 GeV, Sideband width: 25 GeV
- ▶ Central region: 75 - 105 GeV, Sideband width: 20 GeV
- ▶ Central region: 80 - 100 GeV, no background subtraction

For each bin, the largest deviation from the nominal among these variations is the systematic uncertainty due to background subtraction.

$$\sigma_{\text{bgk}} = \max\{|\sigma_{\text{nominal}} - \sigma_{\text{variation}}|\}$$

Likelihood Method : Result for different variation

► Results for different variations



Likelihood Method : Total uncertainties

$$\sigma_{\text{sys}} = \sqrt{\sigma_{\text{truth}}^2 + \sigma_{\text{bgk}}^2}$$

$$\sigma_{\text{tot}} = \sqrt{\sigma_{\text{lik}}^2 + \sigma_{\text{sys}}^2}$$

Method for measuring real efficiencies

Control Region for Real Efficiencies

It uses the Z tag-and-probe method to measure the real efficiencies of electrons and muons.

Pre-selection:

- ▶ Trigger requirement
- ▶ exactly 2 baseline leptons
- ▶ same flavour and opposite charge ($Z \rightarrow ee$, $Z \rightarrow \mu\mu$)
- ▶ $80 < m_{ll} < 100$
- ▶ $p_T > 20$
- ▶ $|\eta| < 2.47$ for electron, $|\eta| < 2.5$ for muon

Requirements for tag lepton

- ▶ $p_T > 25$
- ▶ pass signal requirement

Permutation of tag and probe

Check the probe lepton:

Tight lepton: pass signal requirement

Loose lepton: pass baseline requirement (include tight lepton)

4 possible cases: (slightly simplified)

- ▶ TL: fill loose hist by the subleading lepton
- ▶ LT: fill loose hist by the leading lepton
- ▶ TT: fill both loose and tight hist by the both leading and subleading lepton
- ▶ LL: do nothing

The real eff is calculated by

$$\epsilon = \frac{N_{\text{tight}}}{N_{\text{loose}}}$$

Method for measuring fake efficiencies

Control Region for fake eff for muon

It uses the heavy-flavour fake-enriched region to measure the fake efficiencies of electrons and muons.

For muon fake eff, it uses the muon-muon events.

Pre-selection for muon-muon events:

- ▶ Trigger requirement
- ▶ exactly 2 baseline muons
- ▶ SS
- ▶ at least 1 jet
- ▶ at least 1 b-jet
- ▶ $p_T > 20$
- ▶ $|\eta| < 2.5$

Requirements for the tag lepton

- ▶ $p_T > 40$
- ▶ pass signal requirement

Control Region for fake eff for electron

For electron fake eff, it uses the muon-electron events.

Pre-selection for muon-electron events:

- ▶ Trigger requirement
- ▶ exactly 1 baseline electron and 1 baseline muon
- ▶ SS
- ▶ at least 1 jet
- ▶ at least 1 b-jet
- ▶ $p_T > 20$
- ▶ $|\eta| < 2.5$

Requirements for the tag lepton

- ▶ muon
- ▶ $p_T > 40$
- ▶ pass signal requirement

Calculate the fake eff

The fake eff is calculated by

$$\epsilon = \frac{N_{\text{tight}}^{\text{data}} - N_{\text{tight}}^{\text{prompt MC}}}{N_{\text{loose}}^{\text{data}} - N_{\text{loose}}^{\text{prompt MC}}}$$









The prompt lepton is estimated from MC by using the truth information. The prompt lepton is defined by the MCTruthClassifier, with the ParticleType = IsoElectron or IsoMuon.

Validation plots for fake lepton background

ee channel

	Number of events	
Fakes	$860.738787 \pm 14.359660 \pm 630.863451$ (9663)	
WZ	201.069385 ± 3.094538 (39857)	
charge flip	167.954322 ± 1.110819 (86972)	
WW	45.176809 ± 0.401964 (21140)	
Rare	30.757392 ± 3.403960 (1268)	
ZZ	8.351345 ± 0.345416 (6466)	
$t\bar{t} + V$	8.346022 ± 0.254296 (4158)	
Total BG	$1322.394062 \pm 15.130843 \pm 630.863451$ (169524)	
Data	1201	



ee channel



mumu channel

	Number of events	
WZ	209.219829 ± 2.994737 (42458)	
Fakes	$96.203916 \pm 5.013796 \pm 70.510979$ (1160)	
WW	53.675707 ± 0.409851 (25092)	
Rare	30.460547 ± 4.359663 (1361)	
ZZ	9.474511 ± 0.534342 (7674)	
$t\bar{t} + V$	9.369165 ± 0.284944 (4801)	
charge flip	0.000000 ± 0.000000 (0)	
Total BG	$408.403674 \pm 7.324477 \pm 70.510979$ (82546)	
Data	454	



mumu channel



emu channel

	Number of events	
Fakes	$702.995427 \pm 13.784203 \pm 515.248212$ (8544)	
WZ	516.828375 ± 33.696304 (94571)	
WW	110.280002 ± 0.578575 (51615)	
Rare	58.742111 ± 4.005608 (3037)	
charge flip	36.901490 ± 0.479301 (36998)	
ZZ	23.391749 ± 0.482387 (16698)	
$t\bar{t} + V$	20.739252 ± 0.424107 (10526)	
Total BG	$1469.878407 \pm 36.639692 \pm 515.248212$ (221989)	
Data	1593	

emu channel



Pre-selection plots

Yields in pre-selection



Number of events	
Fakes	$11580.965417 \pm 45.775452$ (107219)
charge flip	3622.756937 ± 2.810481 (3375964)
WZ	691.333566 ± 34.238864 (68626)
ZZ	66.945860 ± 1.049840 (21749)
Rare	47.371873 ± 4.082859 (1662)
WW	38.128141 ± 0.388221 (15666)
$t\bar{t} + V$	1.971267 ± 0.138488 (671)
Total BG	$16049.473060 \pm 57.389305$ (3591557)
(225.0,75.0)	33.08 ± 1.57 (676)
(187.5,37.5)	62.09 ± 3.13 (568)
(175.0,0.0)	74.30 ± 3.57 (556)

Table: Yields in SRjet1



Number of events	
Fakes	$6848.284248 \pm 36.074557$ (66096)
charge flip	2074.427691 ± 2.680118 (1679673)
WZ	776.282836 ± 5.978666 (147413)
WW	153.455294 ± 0.679752 (72175)
Rare	92.181843 ± 6.338615 (3011)
ZZ	62.932966 ± 1.198123 (35945)
$t\bar{t} + V$	16.375388 ± 0.359456 (6082)
Total BG	$10023.940266 \pm 37.235816$ (2010395)
(225.0,75.0)	45.01 ± 1.76 (937)
(187.5,37.5)	86.02 ± 3.84 (754)
(175.0,0.0)	107.96 ± 5.14 (743)

Table: Yields in SRjet23



$$p_T^{l1}$$





p_T^{l2}





$$|\Delta\eta_{ll}|$$







E_T^{miss}





m_{eff} 

$$m_{\mathrm{T}}^{l_1}$$



m_{lj}/m_{ljj}


$$m_{T^2}$$



N-1 plots

Yields in SR



	Number of events	Significance
Fakes	3.295434 ± 0.819175 (24)	
WZ	2.176292 ± 0.397871 (257)	
charge flip	0.472185 ± 0.052737 (265)	
Rare	0.443536 ± 0.111301 (56)	
WW	0.165504 ± 0.023455 (67)	
$t\bar{t} + V$	0.124898 ± 0.046227 (36)	
ZZ	0.055330 ± 0.027713 (31)	
Total BG	6.733178 ± 0.920855 (736)	
(225.0,75.0)	3.33 ± 0.60 (60)	0.74
(187.5,37.5)	3.77 ± 0.95 (31)	0.86
(175.0,0.0)	4.29 ± 0.73 (36)	0.98

Table: Yields in SRjet1



	Number of events	Significance
WZ	1.848838 ± 0.273025 (319)	
Fakes	1.764693 ± 0.709215 (20)	
Rare	0.731015 ± 0.195114 (48)	
WW	0.514488 ± 0.036878 (235)	
charge flip	0.267485 ± 0.029086 (274)	
$t\bar{t} + V$	0.141679 ± 0.030772 (67)	
ZZ	0.067226 ± 0.024995 (24)	
Total BG	5.335425 ± 0.787005 (987)	
(225.0,75.0)	2.35 ± 0.34 (58)	0.57
(187.5,37.5)	4.72 ± 0.74 (47)	1.26
(175.0,0.0)	8.60 ± 1.51 (58)	2.24



Table: Yields in SRjet23

$$p_T^{l1}$$







$$p_T^{l2}$$



$|\Delta\eta_{ll}|$ 



E_T^{miss}





m_{eff} 



m_T^{l1}

Number of events







m_{lj}/m_{ljj}


$$m_{T2}$$







Validation Region : Plots for different variables


VRjet1







VRjet1







VRjet1



VRjet23



VRjet23



VRjet23

