




# Numerical weather prediction

**Numerical weather prediction (NWP)** uses mathematical models of the atmosphere and oceans to predict the weather based on current weather conditions. Though first attempted in the 1920s, it was not until the advent of computer simulation in the 1950s that numerical weather predictions produced realistic results. A number of global and regional forecast models are run in different countries worldwide, using current weather observations relayed from radiosondes, weather satellites and other observing systems as inputs.

Mathematical models based on the same physical principles can be used to generate either short-term weather forecasts or longer-term climate predictions; the latter are widely applied for understanding and projecting climate change. The improvements made to regional models have allowed significant improvements in tropical cyclone track and air quality forecasts; however, atmospheric models perform poorly at handling processes that occur in a relatively constricted area, such as wildfires.

Manipulating the vast datasets and performing the complex calculations necessary to modern numerical weather prediction requires some of the most powerful supercomputers in the world. Even with the increasing power of supercomputers, the forecast skill of numerical weather models extends to only about six days. Factors affecting the accuracy of numerical predictions include the density and quality of observations used as input to the forecasts, along with deficiencies in the numerical models themselves. Post-processing techniques such as model output statistics (MOS) have been developed to improve the handling of errors in numerical predictions.

A more fundamental problem lies in the chaotic nature of the partial differential equations that describe the atmosphere. It is impossible to solve these equations exactly, and small errors grow with time (doubling about every five days). Present understanding is that this chaotic behavior limits accurate forecasts to about 14 days even with accurate input data and a flawless model. In addition, the partial differential equations used in the model need to be supplemented with parameterizations for solar radiation, moist processes (clouds and precipitation), heat exchange, soil, vegetation, surface water, and the effects of terrain. In an effort to quantify the large amount of inherent uncertainty remaining in numerical predictions, ensemble forecasts have been used since the 1990s to help gauge the confidence in the forecast, and to obtain useful results farther into the future than otherwise possible. This approach analyzes multiple forecasts created with an individual forecast model or multiple models.



Weather models use systems of differential equations based on the laws of physics, which are in detail fluid motion, thermodynamics, radiative transfer, and chemistry, and use a coordinate system which divides the planet into a 3D grid. Winds, heat transfer, solar radiation, relative humidity, phase changes of water and surface hydrology are calculated within each grid cell, and the interactions with neighboring cells are used to calculate atmospheric properties in the future.

# History

---

The history of numerical weather prediction began in the 1920s through the efforts of Lewis Fry Richardson, who used procedures originally developed by Vilhelm Bjerknes<sup>[1]</sup> to produce by hand a six-hour forecast for the state of the atmosphere over two points in central Europe, taking at least six weeks to do so.<sup>[2][1][3]</sup> It was not until the advent of the computer and computer simulations that computation time was reduced to less than the forecast period itself. The ENIAC was used to create the first weather forecasts via computer in 1950, based on a highly simplified approximation to the atmospheric governing equations.<sup>[4][5]</sup> In 1954, Carl-Gustav Rossby's group at the Swedish Meteorological and Hydrological Institute used the same model to produce the first operational forecast (i.e., a routine prediction for practical use).<sup>[6]</sup> Operational numerical weather prediction in the United States began in 1955 under the Joint Numerical Weather Prediction Unit (JNWPU), a joint project by the U.S. Air Force, Navy and Weather Bureau.<sup>[7]</sup> In 1956, Norman Phillips developed a mathematical model which could realistically depict monthly and seasonal patterns in the troposphere; this became the first successful climate model.<sup>[8][9]</sup> Following Phillips' work, several groups began working to create general circulation models.<sup>[10]</sup> The first general circulation climate model that combined both oceanic and atmospheric processes was developed in the late 1960s at the NOAA Geophysical Fluid Dynamics Laboratory.<sup>[11]</sup>



The ENIAC main control panel at the Moore School of Electrical Engineering operated by Betty Jennings and Frances Bilas

As computers have become more powerful, the size of the initial data sets has increased and newer atmospheric models have been developed to take advantage of the added available computing power. These newer models include more physical processes in the simplifications of the equations of motion in numerical simulations of the atmosphere.<sup>[6]</sup> In 1966, West Germany and the United States began producing operational forecasts based on primitive-equation models, followed by the United Kingdom in 1972 and Australia in 1977.<sup>[1][12]</sup> The development of limited area (regional) models facilitated advances in forecasting the tracks of tropical cyclones as well as air quality in the 1970s and 1980s.<sup>[13][14]</sup> By the early 1980s models began to include the interactions of soil and vegetation with the atmosphere, which led to more realistic forecasts.<sup>[15]</sup>

The output of forecast models based on atmospheric dynamics is unable to resolve some details of the weather near the Earth's surface. As such, a statistical relationship between the output of a numerical weather model and the ensuing conditions at the ground was developed in the 1970s and 1980s, known as model output statistics (MOS).<sup>[16][17]</sup> Starting in the 1990s, model ensemble forecasts have been used to help define the forecast uncertainty and to extend the window in which numerical weather forecasting is viable farther into the future than otherwise possible.<sup>[18][19][20]</sup>

## Data collection and initialization

---

The atmosphere is a fluid. As such, the idea of numerical weather prediction is to sample the state of the fluid at a given time and use the equations of fluid dynamics and thermodynamics to estimate the state of the fluid at some time in the future. The process of entering observation data into the model to generate initial conditions is called *initialization*. On land, terrain maps available at resolutions down to 1 kilometer (0.6 mi) globally are used to help model atmospheric circulations within regions of rugged topography, in order to better depict features such as downslope winds, mountain waves and related cloudiness that affects incoming solar radiation.<sup>[21]</sup> One main source of input is observations from devices (called radiosondes) in weather balloons which rise through the troposphere and well into the stratosphere that measure various atmospheric parameters and transmits them to a fixed receiver.<sup>[22]</sup> Another main input is data from weather satellites. The World Meteorological Organization acts to standardize the instrumentation, observing practices and timing of these observations worldwide. Stations either report hourly in METAR reports,<sup>[23]</sup> or every six hours in SYNOP reports.<sup>[24]</sup> These observations are irregularly spaced, so they are processed by data assimilation and objective analysis methods, which perform quality control and obtain values at locations usable by the model's mathematical algorithms.<sup>[25]</sup> The data are then used in the model as the starting point for a forecast.<sup>[26]</sup>



Weather reconnaissance aircraft, such as this WP-3D Orion, provide data that is then used in numerical weather forecasts.

Commercial aircraft provide pilot reports along travel routes<sup>[27]</sup> and ship reports along shipping routes.<sup>[28]</sup> Commercial aircraft also submit automatic reports via the WHO's Aircraft Meteorological Data Relay (AMDAR) system, using VHF radio to ground stations or satellites. Research projects use reconnaissance aircraft to fly in and around weather systems of interest, such as tropical cyclones.<sup>[29][30]</sup> Reconnaissance aircraft are also flown over the open oceans during the cold season into systems which cause significant uncertainty in forecast guidance, or are expected to be of high impact from three to seven days into the future over the downstream continent.<sup>[31]</sup> Sea ice began to be initialized in forecast models in 1971.<sup>[32]</sup> Efforts to involve sea surface temperature in model initialization began in 1972 due to its role in modulating weather in higher latitudes of the Pacific.<sup>[33]</sup>


## Computation

---

An atmospheric model is a computer program that produces meteorological information for future times at given locations and altitudes. Within any modern model is a set of equations, known as the primitive equations, used to predict the future state of the atmosphere.<sup>[34]</sup> These equations—along with the ideal gas law—are used to evolve the density, pressure, and potential temperature scalar fields and the air velocity (wind) vector field of the atmosphere through time. Additional transport equations for pollutants and other aerosols are included in some primitive-equation high-resolution models as well.<sup>[35]</sup> The equations used are nonlinear partial differential equations which are impossible to solve exactly through

analytical methods,<sup>[36]</sup> with the exception of a few idealized cases.<sup>[37]</sup> Therefore, numerical methods obtain approximate solutions. Different models use different solution methods: some global models and almost all regional models use finite difference methods for all three spatial dimensions, while other global models and a few regional models use spectral methods for the horizontal dimensions and finite-difference methods in the vertical.<sup>[36]</sup>


These equations are initialized from the analysis data and rates of change are determined. These rates of change predict the state of the atmosphere a short time into the future; the time increment for this prediction is called a *time step*. This future atmospheric state is then used as the starting point for another application of the predictive equations to find new rates of change, and these new rates of change predict the atmosphere at a yet further time step into the future. This time stepping is repeated until the solution reaches the desired forecast time. The length of the time step chosen within the model is related to the distance between the points on the computational grid, and is chosen to maintain numerical stability.<sup>[38]</sup> Time steps for global models are on the order of tens of minutes,<sup>[39]</sup> while time steps for regional models are between one and four minutes.<sup>[40]</sup> The global models are run at varying times into the future. The UK Met Office's Unified Model is run six days into the future,<sup>[41]</sup> while the European Centre for Medium-Range Weather Forecasts' Integrated Forecast System and Environment Canada's Global Environmental Multiscale Model both run out to ten days into the future,<sup>[42]</sup> and the Global Forecast System model run by the Environmental Modeling Center is run sixteen days into the future.<sup>[43]</sup> The visual output produced by a model solution is known as a prognostic chart, or *prog*.<sup>[44]</sup>



A prognostic chart of the 96-hour forecast of 850 mbar geopotential height and temperature from the Global Forecast System

## Parameterization

Some meteorological processes are too small-scale or too complex to be explicitly included in numerical weather prediction models. Parameterization is a procedure for representing these processes by relating them to variables on the scales that the model resolves. For example, the gridboxes in weather and climate models have sides that are between 5 kilometers (3 mi) and 300 kilometers (200 mi) in length. A typical cumulus cloud has a scale of less than 1 kilometer (0.6 mi), and would require a grid even finer than this to be represented physically by the equations of fluid motion. Therefore, the processes that such clouds represent are parameterized, by processes of various sophistication. In the earliest models, if a column of air within a model gridbox was conditionally unstable (essentially, the bottom was warmer and moister than the top) and the water vapor content at any point within the column became saturated then it would be overturned (the warm, moist air would begin



Field of cumulus clouds, which are parameterized since they are too small to be explicitly included within numerical weather prediction




rising), and the air in that vertical column mixed. More sophisticated schemes recognize that only some portions of the box might convect and that entrainment and other processes occur. Weather models that have gridboxes with sizes between 5 and 25 kilometers (3 and 16 mi) can explicitly represent convective clouds, although they need to parameterize cloud microphysics which occur at a smaller scale.<sup>[45]</sup> The formation of large-scale (stratus-type) clouds is more physically based; they form when the relative humidity reaches some prescribed value. The cloud fraction can be related to this critical value of relative humidity.<sup>[46]</sup>

The amount of solar radiation reaching the ground, as well as the formation of cloud droplets occur on the molecular scale, and so they must be parameterized before they can be included in the model. Atmospheric drag produced by mountains must also be parameterized, as the limitations in the resolution of elevation contours produce significant underestimates of the drag.<sup>[47]</sup> This method of parameterization is also done for the surface flux of energy between the ocean and the atmosphere, in order to determine realistic sea surface temperatures and type of sea ice found near the ocean's surface.<sup>[48]</sup> Sun angle as well as the impact of multiple cloud layers is taken into account.<sup>[49]</sup> Soil type, vegetation type, and soil moisture all determine how much radiation goes into warming and how much moisture is drawn up into the adjacent atmosphere, and thus it is important to parameterize their contribution to these processes.<sup>[50]</sup> Within air quality models, parameterizations take into account atmospheric emissions from multiple relatively tiny sources (e.g. roads, fields, factories) within specific grid boxes.<sup>[51]</sup>

## Domains

The horizontal domain of a model is either *global*, covering the entire Earth, or *regional*, covering only part of the Earth. Regional models (also known as *limited-area* models, or LAMs) allow for the use of finer grid spacing than global models because the available computational resources are focused on a specific area instead of being spread over the globe. This allows regional models to resolve explicitly smaller-scale meteorological phenomena that cannot be represented on the coarser grid of a global model. Regional models use a global model to specify conditions at the edge of their domain (boundary conditions) in order to allow systems from outside the regional model domain to move into its area. Uncertainty and errors within regional models are introduced by the global model used for the boundary conditions of the edge of the regional model, as well as errors attributable to the regional model itself.<sup>[52]</sup>



A cross-section of the atmosphere over terrain with a sigma coordinate representation shown. Mesoscale models divide the atmosphere vertically using representations similar to the one shown here.

The vertical coordinate is handled in various ways. Lewis Fry Richardson's 1922 model used geometric height ( $z$ ) as the vertical coordinate. Later models substituted the geometric  $z$  coordinate with a pressure coordinate system, in which the geopotential heights of constant-pressure surfaces become dependent variables, greatly simplifying the primitive equations.<sup>[53]</sup> This correlation between coordinate systems can be made since pressure decreases with height through the Earth's atmosphere.<sup>[54]</sup> The first model used for operational forecasts, the single-layer barotropic model, used a single pressure coordinate at the 500-millibar (about 5,500 m (18,000 ft)) level,<sup>[4]</sup> and thus was essentially two-dimensional. High-





based approach.<sup>[71]</sup> Models within a multi-model ensemble can be adjusted for their various biases, which is a process known as *superensemble forecasting*. This type of forecast significantly reduces errors in model output.<sup>[72]</sup>

## Applications

---

### Air quality modeling

Air quality forecasting attempts to predict when the concentrations of pollutants will attain levels that are hazardous to public health. The concentration of pollutants in the atmosphere is determined by their *transport*, or mean velocity of movement through the atmosphere, their diffusion, chemical transformation, and ground deposition.<sup>[73]</sup> In addition to pollutant source and terrain information, these models require data about the state of the fluid flow in the atmosphere to determine its transport and diffusion.<sup>[74]</sup> Meteorological conditions such as thermal inversions can prevent surface air from rising, trapping pollutants near the surface,<sup>[75]</sup> which makes accurate forecasts of such events crucial for air quality modeling. Urban air quality models require a very fine computational mesh, requiring the use of high-resolution mesoscale weather models; in spite of this, the quality of numerical weather guidance is the main uncertainty in air quality forecasts.<sup>[74]</sup>

### Climate modeling

A General Circulation Model (GCM) is a mathematical model that can be used in computer simulations of the global circulation of a planetary atmosphere or ocean. An atmospheric general circulation model (AGCM) is essentially the same as a global numerical weather prediction model, and some (such as the one used in the UK Unified Model) can be configured for both short-term weather forecasts and longer-term climate predictions. Along with sea ice and land-surface components, AGCMs and oceanic GCMs (OGCM) are key components of global climate models, and are widely applied for understanding the climate and projecting climate change. For aspects of climate change, a range of man-made chemical emission scenarios can be fed into the climate models to see how an enhanced greenhouse effect would modify the Earth's climate.<sup>[76]</sup> Versions designed for climate applications with time scales of decades to centuries were originally created in 1969 by Syukuro Manabe and Kirk Bryan at the Geophysical Fluid Dynamics Laboratory in Princeton, New Jersey.<sup>[77]</sup> When run for multiple decades, computational limitations mean that the models must use a coarse grid that leaves smaller-scale interactions unresolved.<sup>[78]</sup>

### Ocean surface modeling


The transfer of energy between the wind blowing over the surface of an ocean and the ocean's upper layer is an important element in wave dynamics.<sup>[79]</sup> The spectral wave transport equation is used to describe the change in wave spectrum over changing topography. It simulates wave generation, wave movement (propagation within a fluid), wave shoaling, refraction, energy transfer between waves, and wave dissipation.<sup>[80]</sup> Since surface winds are the primary forcing mechanism in the spectral wave transport equation, ocean wave models use information produced by numerical weather prediction models as inputs



to determine how much energy is transferred from the atmosphere into the layer at the surface of the ocean. Along with dissipation of energy through whitecaps and resonance between waves, surface winds from numerical weather models allow for more accurate predictions of the state of the sea surface.<sup>[81]</sup>

## Tropical cyclone forecasting

Tropical cyclone forecasting also relies on data provided by numerical weather models. Three main classes of tropical cyclone guidance models exist: Statistical models are based on an analysis of storm behavior using climatology, and correlate a storm's position and date to produce a forecast that is not based on the physics of the atmosphere at the time. Dynamical models are numerical models that solve the governing equations of fluid flow in the atmosphere; they are based on the same principles as other limited-area numerical weather prediction models but may include special computational techniques such as refined spatial domains that move along with the cyclone. Models that use elements of both approaches are called statistical-dynamical models.<sup>[82]</sup>



NOAA Wavewatch III 120-hour wind and wave forecast for the North Atlantic

In 1978, the first hurricane-tracking model based on atmospheric dynamics—the movable fine-mesh (MFM) model—began operating.<sup>[13]</sup> Within the field of tropical cyclone track forecasting, despite the ever-improving dynamical model guidance which occurred with increased computational power, it was not until the 1980s when numerical weather prediction showed skill, and until the 1990s when it consistently outperformed statistical or simple dynamical models.<sup>[83]</sup> Predictions of the intensity of a tropical cyclone based on numerical weather prediction continue to be a challenge, since statistical methods continue to show higher skill over dynamical guidance.<sup>[84]</sup>

## Weather forecasts

Because weather drifts across the world, producing forecasts a week or more in advance typically involves running a numerical prediction model for the entire planet. Agencies use various software to do this, including:

- North American Ensemble Forecast System, which combines results from:
  - Global Forecast System from the US National Weather Service
  - Global Environmental Multiscale Model from the Canadian Meteorological Centre
- Integrated Forecast System from the European Centre for Medium-Range Weather Forecasts and Météo-France
- Unified Model, produced by a partnership of:
  - UK Met Office
  - Australia Bureau of Meteorology
  - (South) Korea Meteorological Administration
  - India National Centre for Medium Range Weather Forecasting<sup>[85]</sup>
  - New Zealand National Institute of Water and Atmospheric Research

- Icosahedral Nonhydrostatic (ICON) from Deutscher Wetterdienst, the German Meteorological Service
- Navy Global Environmental Model from the US Navy Fleet Numerical Meteorology and Oceanography Center
- Global Spectral Model and Global Ensemble Prediction System from the Japan Meteorological Agency<sup>[86]</sup>
- China Meteorological Administration Global Assimilation Forecasting System<sup>[87]</sup>
- Brazilian Global Atmospheric Model (BAM) from Centro de Previsão do Tempo e Estudos Climáticos (CPTEC)


The global models can be used to supply boundary conditions to higher-resolution models that provide more accurate forecasts for an area of interest, such as the country served by a government agency, or an area where military action or rescue efforts are planned.

- Users of the Unified Model re-run the same system (hence the name) for a specific country or crisis zone at a higher horizontal resolution, feeding it the output of the global Unified Model run. This is given a different name, such as the UKV model or the New Zealand Limited Area Model.<sup>[88]</sup>
- The US National Weather Service runs the Weather Research and Forecasting Model with different parameters to create:
  - North American Mesoscale Model (NAM) every six hours (with an ensemble called Short Range Ensemble Forecast, SREF)
  - Rapid Refresh (RAP) and High Resolution Rapid Refresh (HRRR), every hour<sup>[89][90]</sup>
- The Japan Meteorological Agency runs:<sup>[86]</sup>
  - Meso-Scale Model (MSM) every 3 hours, looking 39 and 78 hours ahead
  - Meso-scale Ensemble Prediction System every 6 hours, looking 39 hours ahead (providing uncertainty estimation)
  - Local Forecast Model every hour, looking 10-18 hours ahead
- The China Meteorological Administration runs the Regional Numerical Forecasting Model (CMA-MESO)<sup>[87]</sup>
- CPTEC runs the Brazilian Regional Atmospheric Modelling System (BRAMS) and ETA Regional Model (ETA) for South America

The output of higher-resolution models may be further modified by model output statistics to take into quirky local phenomena that general models do not handle well, such as mountain waves.

## Wildfire modeling

On a molecular scale, there are two main competing reaction processes involved in the degradation of cellulose, or wood fuels, in wildfires. When there is a low amount of moisture in a cellulose fiber, volatilization of the fuel occurs; this process will generate intermediate gaseous products that will ultimately be the source of combustion. When moisture is present—or when enough heat is being carried away from the fiber, charring occurs. The chemical kinetics of both reactions indicate that there is a point at which the level of moisture is low



A simple wildfire propagation model

enough—and/or heating rates high enough—for combustion processes to become self-sufficient. Consequently, changes in wind speed, direction, moisture, temperature, or lapse rate at different levels of the atmosphere can have a significant impact on the behavior and growth of a wildfire. Since the wildfire acts as a heat source to the atmospheric flow, the wildfire can modify local advection patterns, introducing a feedback loop between the fire and the atmosphere.<sup>[91]</sup>

A simplified two-dimensional model for the spread of wildfires that used convection to represent the effects of wind and terrain, as well as radiative heat transfer as the dominant method of heat transport led to reaction–diffusion systems of partial differential equations.<sup>[92][93]</sup> More complex models join numerical weather models or computational fluid dynamics models with a wildfire component which allow the feedback effects between the fire and the atmosphere to be estimated.<sup>[91]</sup> The additional complexity in the latter class of models translates to a corresponding increase in their computer power requirements. In fact, a full three-dimensional treatment of combustion via direct numerical simulation at scales relevant for atmospheric modeling is not currently practical because of the excessive computational cost such a simulation would require. Numerical weather models have limited forecast skill at spatial resolutions under 1 kilometer (0.6 mi), forcing complex wildfire models to parameterize the fire in order to calculate how the winds will be modified locally by the wildfire, and to use those modified winds to determine the rate at which the fire will spread locally.<sup>[94][95][96]</sup>

## See also

---

- Atmospheric physics
- Atmospheric thermodynamics
- Tropical cyclone forecast model
- Types of atmospheric models

## References

---

1. Lynch, Peter (March 2008). "The origins of computer weather prediction and climate modeling" (<https://web.archive.org/web/20100708191309/http://www.rsmas.miami.edu/personal/miskandarani/Courses/MPO662/Lynch,Peter/OriginsCompWF.JCP227.pdf>) (PDF). *Journal of Computational Physics*. **227** (7): 3431–44. Bibcode:2008JCoPh.227.3431L (<http://ui.adsabs.harvard.edu/abs/2008JCoPh.227.3431L>). doi:10.1016/j.jcp.2007.02.034 (<http://doi.org/10.1016%2Fj.jcp.2007.02.034>). Archived from the original (<http://www.rsmas.miami.edu/personal/miskandarani/Courses/MPO662/Lynch,Peter/OriginsCompWF.JCP227.pdf>) (PDF) on 2010-07-08. Retrieved 2010-12-23.
2. Simmons, A. J.; Hollingsworth, A. (2002). "Some aspects of the improvement in skill of numerical weather prediction" (<https://doi.org/10.1256/003590002321042135>). *Quarterly Journal of the Royal Meteorological Society*. **128** (580): 647–677. Bibcode:2002QJRMS.128..647S (<http://ui.adsabs.harvard.edu/abs/2002QJRMS.128..647S>). doi:10.1256/003590002321042135 (<https://doi.org/10.1256%2F003590002321042135>). S2CID 121625425 (<https://api.semanticscholar.org/CorpusID:121625425>).
3. Lynch, Peter (2006). "Weather Prediction by Numerical Process". *The Emergence of Numerical Weather Prediction* (<https://archive.org/details/emergencenumeric00lync>). Cambridge University Press. pp. 1 (<https://archive.org/details/emergencenumeric00lync/page/n11>)–27. ISBN 978-0-521-85729-1.

4. Charney, Jule; Fjørtoft, Ragnar; von Neumann, John (November 1950). "Numerical Integration of the Barotropic Vorticity Equation" (<https://doi.org/10.3402%2Ftellusa.v2i4.8607>). *Tellus*. **2** (4): 237. Bibcode:1950Tell....2..237C (<https://ui.adsabs.harvard.edu/abs/1950Tell....2..237C>). doi:10.3402/tellusa.v2i4.8607 (<https://doi.org/10.3402%2Ftellusa.v2i4.8607>).
5. Cox, John D. (2002). *Storm Watchers* ([https://archive.org/details/stormwatcherstur00cox\\_df1/page/208](https://archive.org/details/stormwatcherstur00cox_df1/page/208)). John Wiley & Sons, Inc. p. 208 ([https://archive.org/details/stormwatcherstur00cox\\_df1/page/208](https://archive.org/details/stormwatcherstur00cox_df1/page/208)). ISBN 978-0-471-38108-2.
6. Harper, Kristine; Uccellini, Louis W.; Kalnay, Eugenia; Carey, Kenneth; Morone, Lauren (May 2007). "2007: 50th Anniversary of Operational Numerical Weather Prediction" (<https://doi.org/10.1175%2FBAMS-88-5-639>). *Bulletin of the American Meteorological Society*. **88** (5): 639–650. Bibcode:2007BAMS...88..639H (<https://ui.adsabs.harvard.edu/abs/2007BAMS...88..639H>). doi:10.1175/BAMS-88-5-639 (<https://doi.org/10.1175%2FBAMS-88-5-639>).
7. American Institute of Physics (2008-03-25). "Atmospheric General Circulation Modeling" (<http://web.archive.org/web/20080325084036/http://www.aip.org/history/sloan/gcm/>). Archived from the original (<http://www.aip.org/history/sloan/gcm/>) on 2008-03-25. Retrieved 2008-01-13.
8. Phillips, Norman A. (April 1956). "The general circulation of the atmosphere: a numerical experiment". *Quarterly Journal of the Royal Meteorological Society*. **82** (352): 123–154. Bibcode:1956QJRMS..82..123P (<https://ui.adsabs.harvard.edu/abs/1956QJRMS..82..123P>). doi:10.1002/qj.49708235202 (<https://doi.org/10.1002%2Fqj.49708235202>).
9. Cox, John D. (2002). *Storm Watchers* ([https://archive.org/details/stormwatcherstur00cox\\_df1/page/210](https://archive.org/details/stormwatcherstur00cox_df1/page/210)). John Wiley & Sons, Inc. p. 210 ([https://archive.org/details/stormwatcherstur00cox\\_df1/page/210](https://archive.org/details/stormwatcherstur00cox_df1/page/210)). ISBN 978-0-471-38108-2.
10. Lynch, Peter (2006). "The ENIAC Integrations". *The Emergence of Numerical Weather Prediction* (<https://archive.org/details/emergencenumeric00lync>). Cambridge University Press. pp. 206 (<https://archive.org/details/emergencenumeric00lync/page/n216>)–208. ISBN 978-0-521-85729-1.
11. National Oceanic and Atmospheric Administration (2008-05-22). "The First Climate Model" ([http://celebrating200years.noaa.gov/breakthroughs/climate\\_model/welcome.html](http://celebrating200years.noaa.gov/breakthroughs/climate_model/welcome.html)). Retrieved 2011-01-08.
12. Leslie, L.M.; Dietachmeyer, G.S. (December 1992). "Real-time limited area numerical weather prediction in Australia: a historical perspective" (<http://www.bom.gov.au/amoj/docs/1992/leslie2.pdf>) (PDF). *Australian Meteorological Magazine*. **41** (SP): 61–77. Retrieved 2011-01-03.
13. Shuman, Frederick G. (September 1989). "History of Numerical Weather Prediction at the National Meteorological Center" (<https://doi.org/10.1175%2F1520-0434%281989%29004%3C0286%3AHONWPA%3E2.0.CO%3B2>). *Weather and Forecasting*. **4** (3): 286–296. Bibcode:1989WtFor...4..286S (<https://ui.adsabs.harvard.edu/abs/1989WtFor...4..286S>). doi:10.1175/1520-0434(1989)004<0286:HONWPA>2.0.CO;2 (<https://doi.org/10.1175%2F1520-0434%281989%29004%3C0286%3AHONWPA%3E2.0.CO%3B2>).
14. Steyn, D. G. (1991). *Air pollution modeling and its application VIII, Volume 8*. Birkhäuser. pp. 241–242. ISBN 978-0-306-43828-8.
15. Xue, Yongkang; Fennessey, Michael J. (1996-03-20). "Impact of vegetation properties on U. S. summer weather prediction" (<https://web.archive.org/web/20100710080304/http://www.geog.ucla.edu/~yxue/pdf/1996jgr.pdf>) (PDF). *Journal of Geophysical Research*. **101** (D3): 7419. Bibcode:1996JGR...101.7419X (<https://ui.adsabs.harvard.edu/abs/1996JGR...101.7419X>). CiteSeerX 10.1.1.453.551 (<https://citeseerx.ist.psu.edu/viewdoc/summary?doi=10.1.1.453.551>). doi:10.1029/95JD02169 (<https://doi.org/10.1029%2F95JD02169>). Archived from the original (<http://www.geog.ucla.edu/~yxue/pdf/1996jgr.pdf>) (PDF) on 2010-07-10. Retrieved 2011-01-06.



16. Hughes, Harry (1976). *Model output statistics forecast guidance* (<https://apps.dtic.mil/dtic/tr/fulltext/u2/a037148.pdf>) (PDF). United States Air Force Environmental Technical Applications Center. pp. 1–16. Archived (<https://web.archive.org/web/20190617044348/https://apps.dtic.mil/dtic/tr/fulltext/u2/a037148.pdf>) (PDF) from the original on June 17, 2019.
17. Best, D. L.; Pryor, S. P. (1983). *Air Weather Service Model Output Statistics Systems*. Air Force Global Weather Central. pp. 1–90.
18. Toth, Zoltan; Kalnay, Eugenia (December 1997). "Ensemble Forecasting at NCEP and the Breeding Method". *Monthly Weather Review*. **125** (12): 3297–3319. Bibcode:1997MWRv..125.3297T (<https://ui.adsabs.harvard.edu/abs/1997MWRv..125.3297T>). CiteSeerX 10.1.1.324.3941 (<https://citeseerx.ist.psu.edu/viewdoc/summary?doi=10.1.1.324.3941>). doi:10.1175/1520-0493(1997)125<3297:EFANAT>2.0.CO;2 (<https://doi.org/10.1175%2F1520-0493%281997%29125%3C3297%3AEFANAT%3E2.0.CO%3B2>). S2CID 14668576 (<https://api.semanticscholar.org/CorpusID:14668576>).
19. "The Ensemble Prediction System (EPS)" ([https://web.archive.org/web/20101030055238/http://ecmwf.int/products/forecasts/guide/The\\_Ensemble\\_Prediction\\_System\\_EPS\\_1.html](https://web.archive.org/web/20101030055238/http://ecmwf.int/products/forecasts/guide/The_Ensemble_Prediction_System_EPS_1.html)). ECMWF. Archived from the original ([http://ecmwf.int/products/forecasts/guide/The\\_Ensemble\\_Prediction\\_System\\_EPS\\_1.html](http://ecmwf.int/products/forecasts/guide/The_Ensemble_Prediction_System_EPS_1.html)) on 2010-10-30. Retrieved 2011-01-05.
20. Molteni, F.; Buizza, R.; Palmer, T.N.; Petroliagis, T. (January 1996). "The ECMWF Ensemble Prediction System: Methodology and validation". *Quarterly Journal of the Royal Meteorological Society*. **122** (529): 73–119. Bibcode:1996QJRMS.122...73M (<https://ui.adsabs.harvard.edu/abs/1996QJRMS.122...73M>). doi:10.1002/qj.49712252905 (<https://doi.org/10.1002/qj.49712252905>).
21. Stensrud, David J. (2007). *Parameterization schemes: keys to understanding numerical weather prediction models* (<https://books.google.com/books?id=IMXSpRwKNO8C&pg=PA56>). Cambridge University Press. p. 56. ISBN 978-0-521-86540-1.
22. Gaffen, Dian J. (2007-06-07). "Radiosonde Observations and Their Use in SPARC-Related Investigations" (<https://web.archive.org/web/20070607142822/http://www.aero.jussieu.fr/~sparc/News12/Radiosondes.html>). Archived from the original (<http://www.aero.jussieu.fr/~sparc/News12/Radiosondes.html>) on 2007-06-07.
23. National Climatic Data Center (2008-08-20). "Key to METAR Surface Weather Observations" (<https://web.archive.org/web/20021101221848/http://www0.ncdc.noaa.gov/oa/climate/conversion/swometardecoder.html>). National Oceanic and Atmospheric Administration. Archived from the original (<http://www.ncdc.noaa.gov/oa/climate/conversion/swometardecoder.html>) on 2002-11-01. Retrieved 2011-02-11.
24. "SYNOP Data Format (FM-12): Surface Synoptic Observations" (<https://web.archive.org/web/20071230100059/http://weather.unisys.com/wxp/Appendices/Formats/SYNOP.html>). UNISYS. 2008-05-25. Archived from the original (<http://weather.unisys.com/wxp/Appendices/Formats/SYNOP.html>) on 2007-12-30.
25. Krishnamurti, T. N. (January 1995). "Numerical Weather Prediction". *Annual Review of Fluid Mechanics*. **27** (1): 195–225. Bibcode:1995AnRFM..27..195K (<https://ui.adsabs.harvard.edu/abs/1995AnRFM..27..195K>). doi:10.1146/annurev.fl.27.010195.001211 (<https://doi.org/10.1146%2Fannurev.fl.27.010195.001211>). S2CID 122230747 (<https://api.semanticscholar.org/CorpusID:122230747>).
26. "The WRF Variational Data Assimilation System (WRF-Var)" (<https://web.archive.org/web/20070814044336/http://www.mmm.ucar.edu/wrf/WG4/wrfvar/wrfvar-tutorial.htm>). University Corporation for Atmospheric Research. 2007-08-14. Archived from the original (<http://www.mmm.ucar.edu/wrf/WG4/wrfvar/wrfvar-tutorial.htm>) on 2007-08-14.
27. Ballish, Bradley A.; V. Krishna Kumar (November 2008). "Systematic Differences in Aircraft and Radiosonde Temperatures" ([http://amdar.noaa.gov/docs/bams\\_ballish\\_kumar.pdf](http://amdar.noaa.gov/docs/bams_ballish_kumar.pdf)) (PDF). *Bulletin of the American Meteorological Society*. **89** (11): 1689–1708. Bibcode:2008BAMS...89.1689B (<https://ui.adsabs.harvard.edu/abs/2008BAMS...89.1689B>). doi:10.1175/2008BAMS2332.1 (<https://doi.org/10.1175%2F2008BAMS2332.1>). Retrieved 2011-02-16.

28. National Data Buoy Center (2009-01-28). "The WMO Voluntary Observing Ships (VOS) Scheme" ([http://www.vos.noaa.gov/vos\\_scheme.shtml](http://www.vos.noaa.gov/vos_scheme.shtml)). National Oceanic and Atmospheric Administration. Retrieved 2011-02-15.
29. 403rd Wing (2011). "The Hurricane Hunters" (<https://web.archive.org/web/20120530232904/http://www.hurricanehunters.com/>). 53rd Weather Reconnaissance Squadron. Archived from the original (<http://www.hurricanehunters.com>) on 2012-05-30. Retrieved 2006-03-30.
30. Lee, Christopher (2007-10-08). "Drone, Sensors May Open Path Into Eye of Storm" ([https://www.washingtonpost.com/wp-dyn/content/article/2007/10/07/AR2007100700971\\_pf.html](https://www.washingtonpost.com/wp-dyn/content/article/2007/10/07/AR2007100700971_pf.html)). *The Washington Post*. Retrieved 2008-02-22.
31. National Oceanic and Atmospheric Administration (2010-11-12). "NOAA Dispatches High-Tech Research Plane to Improve Winter Storm Forecasts" ([http://www.noaanews.noaa.gov/stories2010/20100112\\_plane.html](http://www.noaanews.noaa.gov/stories2010/20100112_plane.html)). Retrieved 2010-12-22.
32. Stensrud, David J. (2007). *Parameterization schemes: keys to understanding numerical weather prediction models* (<https://books.google.com/books?id=IMXSpRwKNO8C&pg=PA137>). Cambridge University Press. p. 137. ISBN 978-0-521-86540-1.
33. Houghton, John Theodore (1985). *The Global Climate* (<https://books.google.com/books?id=SV04AAAAIAAJ&pg=PA38>). Cambridge University Press archive. pp. 49–50. ISBN 978-0-521-31256-1.
34. Pielke, Roger A. (2002). *Mesoscale Meteorological Modeling* (<https://archive.org/details/mesoscalemeteo00srro>). Academic Press. pp. 48 (<https://archive.org/details/mesoscalemeteo00srro/page/n64>)–49. ISBN 978-0-12-554766-6.
35. Pielke, Roger A. (2002). *Mesoscale Meteorological Modeling* (<https://archive.org/details/mesoscalemeteo00srro>). Academic Press. pp. 18 (<https://archive.org/details/mesoscalemeteo00srro/page/n34>)–19. ISBN 978-0-12-554766-6.
36. Strikwerda, John C. (2004). *Finite difference schemes and partial differential equations* ([https://books.google.com/books?id=SH8R\\_fIZBGIC&pg=PA165](https://books.google.com/books?id=SH8R_fIZBGIC&pg=PA165)). SIAM. pp. 165–170. ISBN 978-0-89871-567-5.
37. Pielke, Roger A. (2002). *Mesoscale Meteorological Modeling* (<https://archive.org/details/mesoscalemeteo00srro>). Academic Press. p. 65 (<https://archive.org/details/mesoscalemeteo00srro/page/n81>). ISBN 978-0-12-554766-6.
38. Pielke, Roger A. (2002). *Mesoscale Meteorological Modeling* (<https://archive.org/details/mesoscalemeteo00srro>). Academic Press. pp. 285 (<https://archive.org/details/mesoscalemeteo00srro/page/n301>)–287. ISBN 978-0-12-554766-6.
39. Sunderam, V. S.; van Albada, G. Dick; Peter, M. A.; Sloot, J. J. Dongarra (2005). *Computational Science – ICCS 2005: 5th International Conference, Atlanta, GA, USA, May 22–25, 2005, Proceedings, Part 1* (<https://books.google.com/books?id=JZiklbXzipwC&pg=PA131>). Springer. p. 132. ISBN 978-3-540-26032-5.
40. Zwiefelhofer, Walter; Kreitz, Norbert; European Centre for Medium Range Weather Forecasts (2001). *Developments in teracomputing: proceedings of the ninth ECMWF Workshop on the Use of High Performance Computing in Meteorology* ([https://books.google.com/books?id=UV6PnF2z5\\_wC&pg=PA276](https://books.google.com/books?id=UV6PnF2z5_wC&pg=PA276)). World Scientific. p. 276. ISBN 978-981-02-4761-4.
41. Chan, Johnny C. L. & Jeffrey D. Kepert (2010). *Global Perspectives on Tropical Cyclones: From Science to Mitigation* (<https://books.google.com/books?id=6gFiunmKWWAC&pg=PA297>). World Scientific. pp. 295–296. ISBN 978-981-4293-47-1. Retrieved 2011-02-24.
42. Holton, James R. (2004). *An introduction to dynamic meteorology, Volume 1* (<https://books.google.com/books?id=fhW5oDv3EPsC&pg=PA474>). Academic Press. p. 480. ISBN 978-0-12-354015-7. Retrieved 2011-02-24.
43. Brown, Molly E. (2008). *Famine early warning systems and remote sensing data* (<https://books.google.com/books?id=mTZvR3R6YdkC&pg=PA121>). Springer. p. 121. Bibcode:2008fews.book.....B (<https://ui.adsabs.harvard.edu/abs/2008fews.book.....B>). ISBN 978-3-540-75367-4. Retrieved 2011-02-24.

44. Ahrens, C. Donald (2008). *Essentials of meteorology: an invitation to the atmosphere* (<http://books.google.com/books?id=2Yn29IFukbgC&pg=PA244>). Cengage Learning. p. 244. ISBN 978-0-495-11558-8.
45. Narita, Masami & Shiro Ohmori (2007-08-06). "3.7 Improving Precipitation Forecasts by the Operational Nonhydrostatic Mesoscale Model with the Kain-Fritsch Convective Parameterization and Cloud Microphysics" (<http://ams.confex.com/ams/pdfpapers/126017.pdf>) (PDF). *12th Conference on Mesoscale Processes*. Retrieved 2011-02-15.
46. Frierson, Dargan (2000-09-14). "The Diagnostic Cloud Parameterization Scheme" ([https://web.archive.org/web/20110401013742/http://www.atmos.washington.edu/~dargan/591/diag\\_cloud.tech.pdf](https://web.archive.org/web/20110401013742/http://www.atmos.washington.edu/~dargan/591/diag_cloud.tech.pdf)) (PDF). University of Washington. pp. 4–5. Archived from the original ([http://www.atmos.washington.edu/~dargan/591/diag\\_cloud.tech.pdf](http://www.atmos.washington.edu/~dargan/591/diag_cloud.tech.pdf)) (PDF) on 2011-04-01. Retrieved 2011-02-15.
47. Stensrud, David J. (2007). *Parameterization schemes: keys to understanding numerical weather prediction models* (<https://books.google.com/books?id=IMXSpRwKNO8C&pg=PA56>). Cambridge University Press. p. 6. ISBN 978-0-521-86540-1.
48. McGuffie, K. & A. Henderson-Sellers (2005). *A climate modelling primer*. John Wiley and Sons. p. 188. ISBN 978-0-470-85751-9.
49. Mel'nikova, Irina N. & Alexander V. Vasilyev (2005). *Short-wave solar radiation in the earth's atmosphere: calculation, observation, interpretation* (<https://books.google.com/books?id=vdg5BgBmMkQC&pg=PA226>). Springer. pp. 226–228. ISBN 978-3-540-21452-6.
50. Stensrud, David J. (2007). *Parameterization schemes: keys to understanding numerical weather prediction models* (<https://books.google.com/books?id=IMXSpRwKNO8C&pg=PA56>). Cambridge University Press. pp. 12–14. ISBN 978-0-521-86540-1.
51. Baklanov, Alexander, Sue Grimmond, Alexander Mahura (2009). *Meteorological and Air Quality Models for Urban Areas* (<https://books.google.com/books?id=wh-Xf0WZQIMC&pg=PA11>). Springer. pp. 11–12. ISBN 978-3-642-00297-7. Retrieved 2011-02-24.
52. Warner, Thomas Tomkins (2010). *Numerical Weather and Climate Prediction* (<https://books.google.com/books?id=6RQ3dnjE8lgC&pg=PA261>). Cambridge University Press. p. 259. ISBN 978-0-521-51389-0.
53. Lynch, Peter (2006). "The Fundamental Equations". *The Emergence of Numerical Weather Prediction* (<https://archive.org/details/emergencenumeric00lync>). Cambridge University Press. pp. 45 (<https://archive.org/details/emergencenumeric00lync/page/n55>)–46. ISBN 978-0-521-85729-1.
54. Ahrens, C. Donald (2008). *Essentials of meteorology: an invitation to the atmosphere* (<http://books.google.com/books?id=2Yn29IFukbgC&pg=PA244>). Cengage Learning. p. 10. ISBN 978-0-495-11558-8.
55. Janjic, Zavis; Gall, Robert; Pyle, Matthew E. (February 2010). "Scientific Documentation for the NMM Solver" (<https://web.archive.org/web/20110823082059/http://nldr.library.ucar.edu/collections/technotes/asset-000-000-000-845.pdf>) (PDF). National Center for Atmospheric Research. pp. 12–13. Archived from the original (<http://nldr.library.ucar.edu/collections/technotes/asset-000-000-000-845.pdf>) (PDF) on 2011-08-23. Retrieved 2011-01-03.
56. Pielke, Roger A. (2002). *Mesoscale Meteorological Modeling* (<https://archive.org/details/mesoscalemeteoro00srro>). Academic Press. pp. 131 (<https://archive.org/details/mesoscalemeteoro00srro/page/n147>)–132. ISBN 978-0-12-554766-6.
57. Baum, Marsha L. (2007). *When nature strikes: weather disasters and the law* ([https://books.google.com/books?id=bIEMoIKX\\_0IC&pg=PA188](https://books.google.com/books?id=bIEMoIKX_0IC&pg=PA188)). Greenwood Publishing Group. p. 189. ISBN 978-0-275-22129-4.

58. Glahn, Harry R.; Lowry, Dale A. (December 1972). "The Use of Model Output Statistics (MOS) in Objective Weather Forecasting" (<https://doi.org/10.1175%2F1520-0450%281972%29011%3C1203%3ATUOMOS%3E2.0.CO%3B2>). *Journal of Applied Meteorology*. **11** (8): 1203–1211. Bibcode:1972JApMe..11.1203G (<https://ui.adsabs.harvard.edu/abs/1972JApMe..11.1203G>). doi:10.1175/1520-0450(1972)011<1203:TUOMOS>2.0.CO;2 (<https://doi.org/10.1175%2F1520-0450%281972%29011%3C1203%3ATUOMOS%3E2.0.CO%3B2>).
59. Gultepe, Ismail (2007). *Fog and boundary layer clouds: fog visibility and forecasting* (<https://books.google.com/books?id=QwzHZ-wV-BAC&pg=PA1144>). Springer. p. 1144. ISBN 978-3-7643-8418-0. Retrieved 2011-02-11.
60. Barry, Roger Graham; Chorley, Richard J. (2003). *Atmosphere, weather, and climate* (<http://books.google.com/books?id=Xs9LiGpNX-AC&pg=PA171>). Psychology Press. p. 172. ISBN 978-0-415-27171-4. Retrieved 2011-02-11.
61. Cox, John D. (2002). *Storm Watchers* ([https://archive.org/details/stormwatcherstur00cox\\_df1/page/222](https://archive.org/details/stormwatcherstur00cox_df1/page/222)). John Wiley & Sons, Inc. pp. 222–224 ([https://archive.org/details/stormwatcherstur00cox\\_df1/page/222](https://archive.org/details/stormwatcherstur00cox_df1/page/222)). ISBN 978-0-471-38108-2.
62. Manousos, Peter (2006-07-19). "Ensemble Prediction Systems" (<http://www.wpc.ncep.noaa.gov/ensembletraining>). Hydrometeorological Prediction Center. Retrieved 2010-12-31.
63. Weickmann, Klaus; Jeff Whitaker; Andres Roubicek; Catherine Smith (2001-12-01). "The Use of Ensemble Forecasts to Produce Improved Medium Range (3–15 days) Weather Forecasts" (<https://web.archive.org/web/20100528082602/http://www.esrl.noaa.gov/psd/spotlight/12012001/>). Climate Diagnostics Center. Archived from the original (<http://www.esrl.noaa.gov/psd/spotlight/12012001/>) on 2010-05-28. Retrieved 2007-02-16.
64. Chakraborty, Arindam (October 2010). "The Skill of ECMWF Medium-Range Forecasts during the Year of Tropical Convection 2008" (<https://doi.org/10.1175%2F2010MWR3217.1>). *Monthly Weather Review*. **138** (10): 3787–3805. Bibcode:2010MWRv..138.3787C (<https://ui.adsabs.harvard.edu/abs/2010MWRv..138.3787C>). doi:10.1175/2010MWR3217.1 (<https://doi.org/10.1175%2F2010MWR3217.1>).
65. Epstein, E.S. (December 1969). "Stochastic dynamic prediction". *Tellus A*. **21** (6): 739–759. Bibcode:1969Tell...21..739E (<https://ui.adsabs.harvard.edu/abs/1969Tell...21..739E>). doi:10.1111/j.2153-3490.1969.tb00483.x (<https://doi.org/10.1111%2Fj.2153-3490.1969.tb00483.x>).
66. Leith, C.E. (June 1974). "Theoretical Skill of Monte Carlo Forecasts" (<https://doi.org/10.1175%2F1520-0493%281974%29102%3C0409%3ATSOMCF%3E2.0.CO%3B2>). *Monthly Weather Review*. **102** (6): 409–418. Bibcode:1974MWRv..102..409L (<https://ui.adsabs.harvard.edu/abs/1974MWRv..102..409L>). doi:10.1175/1520-0493(1974)102<0409:TSOMCF>2.0.CO;2 (<https://doi.org/10.1175%2F1520-0493%281974%29102%3C0409%3ATSOMCF%3E2.0.CO%3B2>).
67. "MOGREPS" (<https://web.archive.org/web/20121022215636/http://www.metoffice.gov.uk/research/areas/data-assimilation-and-ensembles/ensemble-forecasting/MOGREPS>). Met Office. Archived from the original (<http://www.metoffice.gov.uk/research/areas/data-assimilation-and-ensembles/ensemble-forecasting/MOGREPS>) on 2012-10-22. Retrieved 2012-11-01.
68. Warner, Thomas Tomkins (2010). *Numerical Weather and Climate Prediction* (<https://books.google.com/books?id=6RQ3dnjE8lgC&pg=PA261>). Cambridge University Press. pp. 266–275. ISBN 978-0-521-51389-0.
69. Palmer, T.N.; Shutts, G.J.; Hagedorn, R.; Doblas-Reyes, F.J.; Jung, T.; Leutbecher, M. (May 2005). "Representing Model Uncertainty in Weather and Climate Prediction". *Annual Review of Earth and Planetary Sciences*. **33**: 163–193. Bibcode:2005AREPS..33..163P (<https://ui.adsabs.harvard.edu/abs/2005AREPS..33..163P>). doi:10.1146/annurev.earth.33.092203.122552 (<https://doi.org/10.1146%2Fannurev.earth.33.092203.122552>).



70. Gritmit, Eric P.; Mass, Clifford F. (October 2004). "Redefining the Ensemble Spread-Skill Relationship from a Probabilistic Perspective" ([https://web.archive.org/web/20110926204842/http://www.atmos.washington.edu/~ens/pdf/WEM\\_WKSHP\\_2004.epgrimit.pdf](https://web.archive.org/web/20110926204842/http://www.atmos.washington.edu/~ens/pdf/WEM_WKSHP_2004.epgrimit.pdf)) (PDF). University of Washington. Archived from the original ([http://www.atmos.washington.edu/~ens/pdf/WEM\\_WKSHP\\_2004.epgrimit.pdf](http://www.atmos.washington.edu/~ens/pdf/WEM_WKSHP_2004.epgrimit.pdf)) (PDF) on 2011-09-26. Retrieved 2010-01-02.
71. Zhou, Binbin; Du, Jun (February 2010). "Fog Prediction From a Multimodel Mesoscale Ensemble Prediction System" ([http://www.emc.ncep.noaa.gov/mmb/SREF/2222289\\_WAF\\_Feb-2010.official.PDF](http://www.emc.ncep.noaa.gov/mmb/SREF/2222289_WAF_Feb-2010.official.PDF)) (PDF). *Weather and Forecasting*. **25** (1): 303. Bibcode:2010WtFor..25..303Z (<https://ui.adsabs.harvard.edu/abs/2010WtFor..25..303Z>). doi:10.1175/2009WAF2222289.1 (<https://doi.org/10.1175%2F2009WAF2222289.1>). S2CID 4947206 (<https://api.semanticscholar.org/CorpusID:4947206>). Retrieved 2011-01-02.
72. Cane, D.; Milelli, M. (2010-02-12). "Multimodel SuperEnsemble technique for quantitative precipitation forecasts in Piemonte region" (<https://www.nat-hazards-earth-syst-sci.net/10/265/2010/nhess-10-265-2010.pdf>) (PDF). *Natural Hazards and Earth System Sciences*. **10** (2): 265. Bibcode:2010NHES..10..265C (<https://ui.adsabs.harvard.edu/abs/2010NHES..10..265C>). doi:10.5194/nhess-10-265-2010 (<https://doi.org/10.5194%2Fnhess-10-265-2010>). Retrieved 2011-01-02.
73. Daly, Aaron & Paolo Zannetti (2007). *Ambient Air Pollution* (<http://www.envirocomp.org/books/chapters/2aap.pdf>) (PDF). The Arab School for Science and Technology and The EnviroComp Institute. p. 16. Retrieved 2011-02-24.
74. Baklanov, Alexander; Rasmussen, Alix; Fay, Barbara; Berge, Erik; Finardi, Sandro (September 2002). "Potential and Shortcomings of Numerical Weather Prediction Models in Providing Meteorological Data for Urban Air Pollution Forecasting". *Water, Air, & Soil Pollution: Focus*. **2** (5): 43–60. doi:10.1023/A:1021394126149 (<https://doi.org/10.1023%2FA%3A1021394126149>). S2CID 94747027 (<https://api.semanticscholar.org/CorpusID:94747027>).
75. Marshall, John; Plumb, R. Alan (2008). *Atmosphere, ocean, and climate dynamics: an introductory text* (<https://archive.org/details/atmosphereoceanc00mars>). Amsterdam: Elsevier Academic Press. pp. 44 (<https://archive.org/details/atmosphereoceanc00mars/page/n64>)–46. ISBN 978-0-12-558691-7.
76. Australian Bureau of Statistics (2005). *Year book, Australia, Issue 87* ([https://books.google.com/books?id=Pzx\\_Nz1qgd8C&pg=PA40](https://books.google.com/books?id=Pzx_Nz1qgd8C&pg=PA40)). p. 40. Retrieved 2011-02-18.
77. National Oceanic and Atmospheric Administration 200th Celebration (2008-05-22). "The First Climate Model" ([http://celebrating200years.noaa.gov/breakthroughs/climate\\_model/welcome.html](http://celebrating200years.noaa.gov/breakthroughs/climate_model/welcome.html)). National Oceanic and Atmospheric Administration. Retrieved 2010-04-20.
78. Bridgman, Howard A., John E. Oliver, Michael H. Glantz (2006). *The global climate system: patterns, processes, and teleconnections* (<https://books.google.com/books?id=bV3C5VCC-0EC&pg=PA282>). Cambridge University Press. pp. 284–289. ISBN 978-0-521-82642-6. Retrieved 2011-02-18.
79. Chalikov, D. V. (August 1978). "The numerical simulation of wind-wave interaction". *Journal of Fluid Mechanics*. **87** (3): 561–82. Bibcode:1978JFM....87..561C (<https://ui.adsabs.harvard.edu/abs/1978JFM....87..561C>). doi:10.1017/S0022112078001767 (<https://doi.org/10.1017%2FS0022112078001767>). S2CID 122742282 (<https://api.semanticscholar.org/CorpusID:122742282>).
80. Lin, Pengzhi (2008). *Numerical modeling of water waves* (<https://books.google.com/books?id=yBtOwfUG6cgC>). Psychology Press. p. 270. ISBN 978-0-415-41578-1.
81. Bender, Leslie C. (January 1996). "Modification of the Physics and Numerics in a Third-Generation Ocean Wave Model" (<https://doi.org/10.1175%2F1520-0426%281996%29013%3C0726%3AMOTPAN%3E2.0.CO%3B2>). *Journal of Atmospheric and Oceanic Technology*. **13** (3): 726–750. Bibcode:1996JAOT..13..726B (<https://ui.adsabs.harvard.edu/abs/1996JAOT..13..726B>). doi:10.1175/1520-0426(1996)013<0726:MOTPAN>2.0.CO;2 (<https://doi.org/10.1175%2F1520-0426%281996%29013%3C0726%3AMOTPAN%3E2.0.CO%3B2>).

82. National Hurricane Center (July 2009). "Technical Summary of the National Hurricane Center Track and Intensity Models" ([http://www.nhc.noaa.gov/pdf/model\\_summary\\_20090724.pdf](http://www.nhc.noaa.gov/pdf/model_summary_20090724.pdf)) (PDF). National Oceanic and Atmospheric Administration. Retrieved 2011-02-19.
83. Franklin, James (2010-04-20). "National Hurricane Center Forecast Verification" (<http://www.nhc.noaa.gov/verification/verify6.shtml>). National Hurricane Center. Retrieved 2011-01-02.
84. Rappaport, Edward N.; Franklin, James L.; Avila, Lixion A.; Baig, Stephen R.; Beven II, John L.; Blake, Eric S.; Burr, Christopher A.; Jiing, Jiann-Gwo; Juckins, Christopher A.; Knabb, Richard D.; Landsea, Christopher W.; Mainelli, Michelle; Mayfield, Max; McAdie, Colin J.; Pasch, Richard J.; Sisko, Christopher; Stewart, Stacy R.; Tribble, Ahsha N. (April 2009). "Advances and Challenges at the National Hurricane Center". *Weather and Forecasting*. **24** (2): 395–419. Bibcode:2009WtFor..24..395R (<https://ui.adsabs.harvard.edu/abs/2009WtFor..24..395R>). CiteSeerX 10.1.1.207.4667 (<https://citeseerx.ist.psu.edu/viewdoc/summary?doi=10.1.1.207.4667>). doi:10.1175/2008WAF222128.1 (<https://doi.org/10.1175%2F2008WAF222128.1>). S2CID 14845745 (<https://api.semanticscholar.org/CorpusID:14845745>).
85. Global NCMRWF Unified Model (NCUM-G) System (<https://ncmrwf.gov.in/ncmrwf/NCUMG-Writeup-for-WEB-June2020.pdf>)
86. Numerical Weather Prediction Activities (<https://www.jma.go.jp/jma/en/Activities/nwp.html>)
87. Numerical Weather Prediction ([https://www.cma.gov.cn/en/forecast/highlight/202311/t20231117\\_5892086.html](https://www.cma.gov.cn/en/forecast/highlight/202311/t20231117_5892086.html))
88. A 36 hour forecast by NZCSM takes 130 minutes to complete using 810 processors spread across 13 nodes of FitzRoy (<https://www.nesi.org.nz/case-studies/improving-new-zealands-weather-forecasting-ability>)
89. Rapid Refresh (RAP) (<https://rapidrefresh.noaa.gov/>)
90. The High-Resolution Rapid Refresh (HRRR) (<https://rapidrefresh.noaa.gov/hrrr/>)
91. Sullivan, Andrew L. (June 2009). "Wildland surface fire spread modelling, 1990–2007. 1: Physical and quasi-physical models". *International Journal of Wildland Fire*. **18** (4): 349. arXiv:0706.3074 (<https://arxiv.org/abs/0706.3074>). doi:10.1071/WF06143 (<https://doi.org/10.1071%2FWF06143>). S2CID 16173400 (<https://api.semanticscholar.org/CorpusID:16173400>).
92. Asensio, M. I. & L. Ferragut (2002). "On a wildland fire model with radiation". *International Journal for Numerical Methods in Engineering*. **54** (1): 137–157. Bibcode:2002IJNME..54..137A (<https://ui.adsabs.harvard.edu/abs/2002IJNME..54..137A>). doi:10.1002/nme.420 (<https://doi.org/10.1002%2Fnme.420>). S2CID 122302719 (<https://api.semanticscholar.org/CorpusID:122302719>).
93. Mandel, Jan, Lynn S. Bennethum, Jonathan D. Beezley, Janice L. Coen, Craig C. Douglas, Minjeong Kim, and Anthony Vodacek (2008). "A wildfire model with data assimilation". *Mathematics and Computers in Simulation*. **79** (3): 584–606. arXiv:0709.0086 (<https://arxiv.org/abs/0709.0086>). Bibcode:2007arXiv0709.0086M (<https://ui.adsabs.harvard.edu/abs/2007arXiv0709.0086M>). doi:10.1016/j.matcom.2008.03.015 (<https://doi.org/10.1016%2Fj.matcom.2008.03.015>). S2CID 839881 (<https://api.semanticscholar.org/CorpusID:839881>).
94. Clark, T. L., M. A. Jenkins, J. Coen, and David Packham (1996). "A coupled atmospheric-fire model: Convective Froude number and dynamic fingering" (<https://zenodo.org/record/1236052>). *International Journal of Wildland Fire*. **6** (4): 177–190. doi:10.1071/WF9960177 (<https://doi.org/10.1071%2FWF9960177>).
95. Clark, Terry L., Marry Ann Jenkins, Janice Coen, and David Packham (1996). "A coupled atmospheric-fire model: Convective feedback on fire line dynamics" (<https://doi.org/10.1175%2F1520-0450%281996%29035%3C0875%3AACAMCF%3E2.0.CO%3B2>). *Journal of Applied Meteorology*. **35** (6): 875–901. Bibcode:1996JApMe..35..875C (<https://ui.adsabs.harvard.edu/abs/1996JApMe..35..875C>). doi:10.1175/1520-0450(1996)035<0875:ACAMCF>2.0.CO;2 (<https://doi.org/10.1175%2F1520-0450%281996%29035%3C0875%3AACAMCF%3E2.0.CO%3B2>).

96. Rothermel, Richard C. (January 1972). "A mathematical model for predicting fire spread in wildland fires" ([https://web.archive.org/web/20220324005215/https://www.fs.fed.us/rm/pubs\\_int/int\\_rp115.pdf](https://web.archive.org/web/20220324005215/https://www.fs.fed.us/rm/pubs_int/int_rp115.pdf)) (PDF). United States Forest Service. Archived from the original ([http://www.fs.fed.us/rm/pubs\\_int/int\\_rp115.pdf](http://www.fs.fed.us/rm/pubs_int/int_rp115.pdf)) (PDF) on March 24, 2022. Retrieved 2011-02-28.

## Further reading

---

- Beniston, Martin (1998). *From Turbulence to Climate: Numerical Investigations of the Atmosphere with a Hierarchy of Models*. Berlin: Springer. ISBN 978-3-540-63495-9.
- Blum, Andrew (2019). *The Weather Machine: A Journey Inside the Forecast*. New York: HarperCollins. ISBN 978-0-062-36861-4.
- Kalnay, Eugenia (2003). *Atmospheric Modeling, Data Assimilation and Predictability*. New York: Cambridge University Press. ISBN 978-0-521-79629-3.
- Roulstone, Ian & Norbury, John (2013). *Invisible in the Storm: the role of mathematics in understanding weather* (<https://books.google.com/books?id=qnMrFEHMrWwC>). Princeton University Press. ISBN 978-0691152721.
- Thompson, Philip (1961). *Numerical Weather Analysis and Prediction* (<https://archive.org/details/numericalweather00thom>). New York: The Macmillan Company.
- U.S. Department of Commerce; National Oceanic; Atmospheric Administration; National Weather Service, eds. (1979). *National Weather Service Handbook No. 1 – Facsimile Products*. Washington, DC: Department of Commerce. From Turbulence to sCI

## External links

---

- NOAA Supercomputer upgrade (<https://web.archive.org/web/20191012133855/https://www.noaa.gov/media-release/noaa-kicks-off-2018-with-massive-supercomputer-upgrade>)
- Air Resources Laboratory (<https://www.arl.noaa.gov/>)
- Fleet Numerical Meteorology and Oceanography Center (<https://web.archive.org/web/20180111155953/http://www.metoc.navy.mil/fnmoc/fnmoc.html>)
- European Centre for Medium-Range Weather Forecasts (<https://www.ecmwf.int/>)
- UK Met Office (<https://www.metoffice.gov.uk/research/approach/modelling-systems/unified-model/weather-forecasting>)

---

Retrieved from "[https://en.wikipedia.org/w/index.php?title=Numerical\\_weather\\_prediction&oldid=1286480147](https://en.wikipedia.org/w/index.php?title=Numerical_weather_prediction&oldid=1286480147)"