

Notes from *Functional Analysis, Sobolev Spaces, and  
PDEs*

by Haim Brezis

taken by Samuel T. Wallace

## **Publisher's Description**

Uniquely, this book presents a coherent, concise and unified way of combining elements from two distinct worlds, functional analysis (FA) and partial differential equations (PDEs), and is intended for students who have a good background in real analysis. This text presents a smooth transition from FA to PDEs by analyzing in great detail the simple case of one-dimensional PDEs (i.e., ODEs), a more manageable approach for the beginner. Although there are many books on functional analysis and many on PDEs, this is the first to cover both of these closely connected topics. Moreover, the wealth of exercises and additional material presented, leads the reader to the frontier of research. This book has its roots in a celebrated course taught by the author for many years and is a completely revised, updated, and expanded English edition of the important *Analyse Fonctionnelle* (1983). Since the French book was first published, it has been translated into Spanish, Italian, Japanese, Korean, Romanian, Greek and Chinese. The English version is a welcome addition to this list. The first part of the text deals with abstract results in FA and operator theory. The second part is concerned with the study of spaces of functions (of one or more real variables) having specific differentiability properties, e.g., the celebrated Sobolev spaces, which lie at the heart of the modern theory of PDEs. The Sobolev spaces occur in a wide range of questions, both in pure and applied mathematics, appearing in linear and nonlinear PDEs which arise, for example, in differential geometry, harmonic analysis, engineering, mechanics, physics etc. and belong in the toolbox of any graduate student studying analysis.

## **Transcription Notes**

placeholder

# Contents

0.1	The Hahn-Banach Theorems. . . . .	3
0.1.1	The Analytic Form of the Hahn-Banach Theorem: Extensions of Linear Functionals . . . . .	3
0.1.2	The Geometric Forms of the Hahn-Banach Theorem: Separation of Convex Sets . . . . .	3
0.1.3	The Bidual $E^{**}$ . Orthogonality Relations . . . . .	4
0.1.4	A Quick Introduction to the Theory of Conjugate Convex Functions . . . . .	5

## 0.1 The Hahn-Banach Theorems.

### 0.1.1 The Analytic Form of the Hahn-Banach Theorem: Extensions of Linear Functionals

Let  $E$  be a vector space over  $\mathbb{R}$ . We recall that a *functional* is a function defined on  $E$ , or a subspace of  $E$ , with values in  $\mathbb{R}$ .

**Theorem 1** (Helly, Hahn-Banach analytic form). *Let  $p : E \rightarrow \mathbb{R}$  be a function satisfying*

1.  $p(\lambda x) = \lambda p(x)$
2.  $p(x + y) \leq p(x) + p(y)$

*Let  $G \subset E$  be a linear subspace and let  $g : G \rightarrow \mathbb{R}$  be a linear functional such that*

$$g(x) \leq p(x) \quad \forall x \in G$$

*Then there exists a linear functional  $f : E \rightarrow \mathbb{R}$  such that  $f|_G = g$ .*

### 0.1.2 The Geometric Forms of the Hahn-Banach Theorem: Separation of Convex Sets

**Definition 1.** *An affine hyperplane is a subset  $H \subset E$  of the form*

$$H = \{x \in E : f(x) = \alpha\}$$

where  $f$  is a linear functional that does not vanish identically and  $\alpha \in \mathbb{R}$ . We write  $H = [f = \alpha]$  and say that  $f = \alpha$  is the equation of  $H$ .

**Proposition 1.** *The hyperplane  $H = [f = \alpha]$  is closed if and only if  $f$  is continuous.*

**Definition 2.** *Let  $A$  and  $B$  be two subsets of  $E$ . We say the hyperplane  $H = [f = \alpha]$  separates  $A$  and  $B$  if*

$$f(a) \leq \alpha \quad \forall a \in A \text{ and } f(b) \geq \alpha \quad \forall b \in B$$

*We say that  $H$  strictly separates  $A$  and  $B$  if there exists an  $\epsilon > 0$  such that*

$$f(a) \leq \alpha - \epsilon \quad \forall a \in A \text{ and } f(b) \geq \alpha + \epsilon \quad \forall b \in B$$

*A subset  $A \subset E$  is convex if*

$$tx + (1 - t)y \in A \quad \forall x, y \in A \quad \forall t \in [0, 1]$$

**Theorem 2** (Hahn-Banach, first geometric form). *Let  $A \subset E$  and  $B \subset E$  be two nonempty convex disjoint subsets, one of which is open. Then there exists a closed hyperplane separating them.*

**Theorem 3** (Hahn-Banach, second geometric form). *Let  $A \subset E$  and  $B \subset E$  be two empty convex disjoint subsets. If  $A$  is closed and  $B$  is compact, then there exists a closed hyperplane separating  $A$  and  $B$ .*

**Corollary 1.** *Let  $F \subset E$  be a linear subspace such that  $\overline{F} \neq E$ . Then there exists some  $f \in E^*$  not identically zero such that  $f(F) = 0$*

### 0.1.3 The Bidual $E^{**}$ . Orthogonality Relations

Let  $E$  be a normed vector space and let  $E^*$  be the dual space with norm

$$\|f\|_{E^*} = \sup_{\|x\| \leq 1} |\langle f, x \rangle|$$

The bidual  $E^{**}$  is the dual of  $E^*$  with norm

$$\|\xi\|_{E^{**}} = \sup_{\|f\| \leq 1} |\langle \xi, f \rangle|$$

There is a canonical injection  $J : E \rightarrow E^{**}$  defined as

$$\langle Jx, f \rangle_{E^{**}, E^*} = \langle f, x \rangle_{E^*, E}$$

which is an *isometry*.  $J$  may not be surjective, but if it is, we say  $E$  is reflexive.

**Definition 3.** *If  $M \subset E$  is a linear subspace, let*

$$M^\perp = \{f \in E^* : \langle f, x \rangle = 0 \quad \forall x \in M\}$$

*If  $N \subset E^*$  is a linear subspace we set*

$$N^\perp = \{x \in E : \langle f, x \rangle = 0 \quad \forall f \in N\}$$

**Proposition 2.** *Let  $M \subset E$  be a linear subspace. Then*

$$(M^\perp)^\perp = \overline{M}$$

*Let  $N \subset E^*$  be a linear subspace. Then*

$$\overline{N} \subset (N^\perp)^\perp$$

#### 0.1.4 A Quick Introduction to the Theory of Conjugate Convex Functions

**Definition 4.** *Let  $E$  be a set, and  $\phi : E \rightarrow (-\infty, +\infty]$  a function. Let*

$$D(\phi) = \{x \in E : \phi(x) < +\infty\}$$

*be the domain of  $\phi$ . We define the epigraph of  $\phi$*

$$\text{epi}\phi = \{[x, \lambda] \in E \times \mathbb{R} ; \phi(x) \leq \lambda\}$$

*If  $E$  is a topological space, we say  $\phi$  is lower semicontinuous if  $\lambda \in \mathbb{R}$  the set*

$$[\phi \leq \lambda] = \{x \in E : \phi(x) \leq \lambda\}$$

*is closed.*

**Proposition 3.** *If  $\phi$  is lower-semicontinuous, then*

1.  *$\text{epi}\phi$  is closed in  $E \times \mathbb{R}$  and conversely,*
2. *for every  $x \in E$  and  $\epsilon > 0$  there is a neighborhood  $V$  of  $x$  such that*

$$\phi(y) \geq \phi(x) - \epsilon \quad \forall y \in V$$

*and conversely.*

3. *If  $\phi_1$  and  $\phi_2$  are lower semicontinuous, then so is  $\phi_1 + \phi_2$*
4. *If  $(\phi_i)_{i \in I}$  is a family of lsc functions then so is*

$$\phi(x) = \sup_{i \in I} \phi_i(x)$$

*called the superior envelope.*

5. *If  $E$  is compact and  $\phi$  is lsc, then  $\inf_E \phi$  is achieved.*

**Definition 5.** *A function  $\phi : E \rightarrow (-\infty, +\infty]$  is convex if*

$$\phi(tx + (1-t)y) \leq t\phi(x) + (1-t)\phi(y) \quad \forall x, y \in E, \quad \forall t \in (0, 1)$$

**Proposition 4.** *If  $\phi$  is a convex function, then*

1.  $\text{epi}\phi$  is a convex set in  $E \times \mathbb{R}$  and conversely
2.  $\forall \lambda \in \mathbb{R}$  the set  $[\phi \leq \lambda]$  is convex, but not the converse
3. a sum of convex functions is again convex
4. the superior envelope of a family of convex functions is again convex.

Let  $E$  be a normed vector space.

**Definition 6.** Let  $\phi : E \rightarrow (-\infty, +\infty]$  be a function with nonempty domain. We define the conjugate function  $\phi^* : E^* \rightarrow (-\infty, +\infty]$  by

$$\phi^*(f) = \sup_{x \in E} \{\langle f, x \rangle - \phi(x)\}$$

**Proposition 5.** Assume that  $\phi : E \rightarrow (-\infty, +\infty]$  is convex lsc with nonempty domain. Then  $\phi^*$  has nonempty domain and is bounded below by an affine continuous function.

**Definition 7.** Instead of defining  $\phi^{**}$  on  $E^{**}$ , we can define it on  $E$  by

$$\phi^{**}(x) = \sup_{f \in E^*} \{\langle f, x \rangle - \phi^*(f)\}$$

**Theorem 4** (Fenchel-Moreau). Let  $\phi : E \rightarrow (-\infty, +\infty]$  is convex lsc with nonempty domain. Then  $\phi^{**} = \phi$ .

**Theorem 5** (Fenchel-Rockafeller). Let  $\phi, \psi$  be two convex functions. Assume there is some  $x_0 \in D(\phi) \cap D(\psi)$  such that  $\phi$  is continuous at  $x_0$ . Then

$$\begin{aligned} \inf_{x \in E} \{\phi(x) + \psi(x)\} &= \sup_{f \in E^*} \{-\phi^*(-f) - \psi^*(f)\} \\ &= \max_{f \in E^*} \{-\phi^*(-f) - \psi^*(f)\} = - \min_{f \in E^*} \{\phi^*(-f) + \psi^*(f)\} \end{aligned}$$