Oral Health Risk Assessment Tool



The American Academy of Pediatrics (AAP) has developed this tool to aid in the implementation of oral health risk assessment during health supervision visits. This tool has been reviewed and endorsed by the National Interprofessional Initiative on Oral Health.

Instructions for Use: This tool is intended for documenting caries risk of the child, though some risk factors are based on the caregiver's oral health. All other factors and findings should be documented for the child.

Child Information

| Child Name | Alice Smith |
|--------------------------|----------------------------------------------|
| Case Number | 67890 |
| Assessment Date | 2024-03-12 |
| Date of Birth | 2019-09-01 |
| Dental Appointments | next Monday |
| Nursing Note | Child appears cooperative. Routine check-up. |
| Required Comment | Follow up on dietary habits. |
| RN Completing Assessment | N/A |

| Risk Factors | |
|---------------------------|--|
| Mother Active Decay | |
| Mother No Dentist | |
| Bottle Usage | |
| Frequent Snacking | |
| Special Health Care Needs | |
| Medicaid Eligible | |

| Protective Factors | |
|----------------------|-----------------------------------------------|
| Existing Dental Home | |
| Fluoridated Water | |
| Fluoride Varnish | |
| Brushing Twice Daily | |
| | , <u>, </u> |
| | |

Clinical Findings White Spots Obvious Decay Fillings Present Plaque Accumulation Gingivitis Teeth Present Healthy Teeth Assessment Plan Caries Risk Anticipatory Guidance Fluoride Varnish Completed Dental Referral

Self Management Goals

| Regular Dental Visits | |
|---------------------------------|--|
| Dental Treatment for Caregivers | |
| Brush Twice Daily | |
| Use Fluoride Toothpaste | |
| Wean Bottle | |
| Less or No Juice | |
| Water in Sippy | |
| Drink Tap Water | |
| Healthy Snacks | |
| Less or No Junk Food | |
| No Soda | |
| Xylitol | |

Nursing Recommendations

Maintain current routine and schedule next check-up in 6 months.

Treatment of High Risk Children

If appropriate, high-risk children should receive professionally applied fluoride varnish and have their teeth brushed twice daily with an age-appropriate amount of fluoridated toothpaste. Referral to a pediatric dentist or a dentist comfortable caring for children should be made to ensure proper follow-up.