


# Methods of Differentiation

## One Shot

Hello NVians!!





# Nishant Vora

**B.Tech - IIT Patna**

- ❑ 7+ years Teaching experience
- ❑ Mentored 5 lac+ students
- ❑ Teaching Excellence Award




# Join with us in Telegram

## TELEGRAM CHANNEL

- [t.me/unacademyatoms](https://t.me/unacademyatoms) ★

## COMPLETE NOTES AND LECTURES

- [livedaily.me/atoms](https://livedaily.me/atoms)



# + India's BEST Educators

# Unacademy Subscription

**BATCH**

HINDI

Igneous Batch for JEE Advanced & Olympiads 2021

Started on Jul 7

Nishant Vora and 1 more

**BATCH**

HINDI

Evolve Batch Course for Class 12th JEE Main and Advanced 2022

Starts on Apr 7

Anupam Gupta and 2 more

**BATCH**

HINDI

Mega Batch Course for Class 12th JEE Main and Advanced 2022

Starts on Apr 6

Narendra Avasthi and 1 more

**BATCH**

HINDI

Enthuse: Class 12th for JEE Main and Advanced 2022

Starts on Apr 14

Amarnath Anand and 2 more

**BATCH**

HINDI

Final Rapid Revision Batch for JEE Main 2021

Starts on Apr 6

Manoj Chauhan and 2 more

**LIVE**

HINDI PHYSICS

Course of 12th syllabus Physics for JEE Aspirants 2022: Part - I






Lesson 1 • Apr 2, 2021 12:30 PM

D C Pandey

If you want to be the **BEST**  
“Learn” from the **BEST**



Last min  
JEE Adv  
RBC 10  
20  
RRB





×

IIT JEE subscription

ICONIC <sup>+</sup> PLUS

✓ India's Best Educators

✓ Interactive Live Classes

✓ Structured Courses & PDFs

✓ Live Tests & Quizzes

✗ 1:1 Live Mentorship

✗ Live Doubt Solving

NVLIVE

30 months

₹1,980/mo

No cost EMI

+10% OFF ₹59,399

>

24 months

₹2,310/mo

No cost EMI

+10% OFF ₹55,440

>

12 months

₹3,176/mo

No cost EMI

+10% OFF ₹38,115

>

6 months

₹4,620/mo

No cost EMI

+10% OFF ₹27,720

>

3 months

₹5,775/mo

+10% OFF ₹17,325

>

To be paid as a one-time payment

NVLIVE

×

×

IIT JEE subscription

PLUS ICONIC <sup>+</sup>

✓ India's Best Educators

✓ Interactive Live Classes

✓ Structured Courses & PDFs

✓ Live Tests & Quizzes

✓ Personal Coach

✓ Study Planner

24 months

₹3,750/mo

No cost EMI

+10% OFF ₹90,000

>

18 months

₹4,000/mo

No cost EMI

+10% OFF ₹72,000

>

12 months

₹4,875/mo

No cost EMI

+10% OFF ₹58,500

>

6 months

₹5,700/mo

No cost EMI

+10% OFF ₹34,200

>

To be paid as a one-time payment

NVLIVE

×



# Standard Derivatives



# Standard Derivatives



$$\sqrt{x} \rightarrow \frac{1}{2\sqrt{x}}$$

$$\frac{d}{dx}(a^x) = a^x \ln a$$

$$\frac{d}{dx}(\ln x) = \frac{1}{x}$$

y or f(x)	$\frac{dy}{dx}$ or f'(x)
$K$ , K is a constant	$\frac{d}{dx}(K) = 0$
$x^n$	$\frac{d}{dx}(x^n) = nx^{n-1}$
$\sqrt{x}$	$\frac{d}{dx}(\sqrt{x}) = \frac{1}{2\sqrt{x}}$
$e^x$	$\frac{d}{dx}(e^x) = e^x$
$a^x$ , $a > 0$	$\frac{d}{dx}(a^x) = \underline{\underline{a^x \log a}}$



# Standard Derivatives



y or f(x)	$\frac{dy}{dx}$ or f'(x)
$\sin x$	$\frac{d}{dx} (\sin x) = \cos x$
$\cos x$	$\frac{d}{dx} (\cos x) = -\sin x$
$\tan x$	$\frac{d}{dx} (\tan x) = \sec^2 x$
$\cot x$	$\frac{d}{dx} (\cot x) = -\operatorname{cosec}^2 x$
$\operatorname{cosec} x$	$\frac{d}{dx} (\operatorname{cosec} x) = -\operatorname{cosec} x \cot x$
$\sec x$	$\frac{d}{dx} (\sec x) = \sec x \tan x$





# Standard Derivatives



$$\frac{d}{dx}(\sin^{-1}x) = \frac{1}{\sqrt{1-x^2}}$$

$$\frac{d}{dx}(\cos^{-1}x) = \frac{-1}{\sqrt{1-x^2}}$$

$$\frac{d}{dx}(\tan^{-1}x) = \frac{1}{1+x^2}$$

$$\frac{d}{dx}(\cot^{-1}x) = \frac{-1}{1+x^2}$$

$$\frac{d}{dx}(\sec^{-1}x) = \frac{1}{x\sqrt{x^2-1}}$$

$$\frac{d}{dx}(\operatorname{cosec}^{-1}x) = \frac{-1}{x\sqrt{x^2-1}}$$



# # Algebra of Differentiation



# Algebra of Differentiation



①

$$\frac{d}{dx}(f(x) \pm g(x)) = \frac{d}{dx}f(x) \pm \frac{d}{dx}g(x)$$

Eg.

$$\frac{d}{dx}(x^3 + x^2 + 5)$$

$$= \frac{d}{dx}(x^3) + \frac{d}{dx}(x^2) + \cancel{\frac{d}{dx}(5)}$$

$$= \underline{3x^2 + 2x'}$$



# Algebra of Differentiation

eg.

$$\frac{d}{dx} (x^3 + 2 + \ln x + \sqrt{x} + \operatorname{cosec} x)$$

$$\underline{3x^2 + 0 + \frac{1}{x} + \frac{1}{2\sqrt{x}} - \operatorname{cosec} x \cot x}$$



# Algebra of Differentiation

2.

$$\frac{d}{dx}(\underline{k} \underline{f(x)}) = k \frac{d}{dx}(f(x))$$

Eg.  $\frac{d}{dx}(5x^4 + \underline{3}e^x + 2 \sin x)$

$$= \frac{d}{dx}(x^4) + 3 \frac{d}{dx}(e^x) + 2 \frac{d}{dx}(\sin x)$$

$$\underline{5(4x^3) + 3e^x + 2 \cos x}$$



# # Product Rule



## Product Rule

$$\frac{d}{dx} (\overset{\text{I}}{f(x)} \cdot \overset{\text{II}}{g(x)}) = f(x) \frac{d}{dx} g(x) + g(x) \frac{d}{dx} f(x)$$

$$(I \cdot II)' = I \cdot II' + II \cdot I'$$

Eg.  $\frac{d}{dx} (x^3 \sin x)$

$$= x^3 \frac{d}{dx} (\sin x) + \sin x \frac{d}{dx} (x^3)$$

$$= \underline{x^3 \cos x + \sin x (3x^2)}$$



## Product Rule

$$\frac{d}{dx}(\underline{x}^2 e^x) = x^2 e^x + e^x (2x)$$

$$\frac{d}{dx}(\overset{\text{I}}{\underline{x}} \overset{\text{II}}{\ln x}) = x \times \frac{1}{x} + \ln x (1)$$

$$= \boxed{1 + \ln x}$$





If  $(a + \sqrt{2}b \cos x)(a - \sqrt{2}b \cos y) = a^2 - b^2$ , where  $a > b > 0$ , then  $dx/dy$  at  $(\pi/4, \pi/4)$  is :


A.  $a - 2b / a + 2b$

B.  $a - b / a + b$

☒ C.  $a + b / a - b$

D.  $2a + b / 2a - b$

JEE Main 2020



$$\underbrace{(a + \sqrt{2}b \cos x)}_{\text{I}} \underbrace{(a - \sqrt{2}b \cos y)}_{\text{II}} = \underline{a^2 - b^2}$$

$$(a + \cancel{\sqrt{2}b \cos x}) \left( +\cancel{\sqrt{2}b \sin y} \frac{dy}{dx} \right) - (a - \cancel{\sqrt{2}b \cos y}) (\cancel{\sqrt{2}b \sin x}) = 0$$

$$x = \frac{\pi}{4} \quad y = \frac{\pi}{4}$$

$$(a+b) \left( \cancel{1} \frac{dy}{dx} \right) - (a-b) \cancel{1} = 0$$

$$\cdot \frac{dy}{dx} = \frac{a-b}{a+b}$$

$$\boxed{\frac{dx}{dy} = \frac{a+b}{a-b}}$$



# # Extended Product Rule



## Extended Product Rule



$$(I \cdot II \cdot III)' = I' \cdot II \cdot III + I \cdot II' \cdot III + I \cdot II \cdot III'$$



If  $f(x) = \underbrace{(1+x)}_I \underbrace{(3+x^2)^{1/2}}_II \underbrace{(9+x^3)^{1/3}}_III$  then  $f'(-1)$  is equal to

A. 0

B.  $2\sqrt{2}$

☒ C. 4

D. 6

$$f'(x) = (1) (3+x^2)^{1/2} (9+x^3)^{1/3} + \cancel{(1+x) II' III} + \cancel{(1+x) II III'}$$

$$f'(-1) = (1) (4)^{1/2} (8)^{1/3}$$

$$= 2 \times 2$$

$$= \textcircled{4}$$



#

# Quotient Rule



## Quotient Rule

$\frac{u}{v}$

$$\frac{d}{dx} \left( \frac{f(x)}{g(x)} \right) = \frac{f'(x)g(x) - g'(x)f(x)}{g^2(x)}$$

$$\frac{d}{dx} \left( \frac{u}{v} \right) = \frac{u'v - uv'}{v^2}$$



# # Chain Rule





## Chain Rule # NVStyle

$$\frac{d}{dx} f(g(x)) = \underline{f'(g(x))} \cdot \underline{g'(x)}$$

$$\square^3 \rightarrow 3 \square^2$$

Eg.

$$\frac{d}{dx} (\sin^3(2x+5))$$

$$= 3 \sin^2(2x+5) \times \cos(2x+5) \times 2$$

**P**



**Power**

**B**



**Base**

**A**



**Angle**



## Chain Rule



1.  $\frac{d}{dx}(\tan^5(\underline{7x+2})) = 5 \tan^4(7x+2) \sec^2(7x+2) \times 7$

2.  $\frac{d}{dx}(\cos^2(\underline{x^3})) = 2 \cos(x^3) (-\sin(x^3)) (3x^2)$

3.  $\frac{d}{dx}(\ln^5(\underline{2x+3})) = 5 \ln^4(2x+3) \times \frac{1}{2x+3} \times 2$



If  $f(1) = 1$ ,  $f'(1) = 3$ , then the derivative of  $f(f(f(x))) + (f(x))^2$  at  $x = 1$  is :

- ☒ A. 33
- ☐ B. 12
- ☐ C. 15
- ☐ D. 9
- $$= f'(f(f(1))) f'(f(1)) f'(1) + 2 f(1) f'(1)$$
- $$= f'(f(1)) f'(1) f'(1) + 2(1)(3)$$
- $$= f'(1) f'(1) f'(1) + 6$$
- $$= 3^3 + 6$$
- $$= 33$$

JEE Main 2019



"NV Tip"

**First**  
**Simplify**  
**then**  
**Differentiate**





i. If  $y = \frac{x^4 + x^2 + 1}{x^2 + x + 1}$  then  $\frac{dy}{dx} = ax + b$  find a and b

# class 8<sup>th</sup>

\*\*\*

$$x^4 + x^2 + 1 = (x^2 + x + 1)(x^2 - x + 1)$$

$$y = \frac{(x^2 + x + 1)(x^2 - x + 1)}{(x^2 + x + 1)}$$

$$y = x^2 - x + 1$$

$$\frac{dy}{dx} = 2x - 1$$

$$ax + b$$

$$\begin{aligned} a &= 2 \\ b &= -1 \end{aligned}$$

$$\begin{aligned} &x^4 + x^2 + 1 \\ &= x^4 + 2x^2 + 1 - x^2 \\ &= (x^2 + 1)^2 - (x)^2 \\ &= (x^2 + 1 + x)(x^2 + 1 - x) \end{aligned}$$



If  $y(x) = \cot^{-1} \left( \frac{\sqrt{1+\sin x} + \sqrt{1-\sin x}}{\sqrt{1+\sin x} - \sqrt{1-\sin x}} \right)$ ,  $x \in \left( \frac{\pi}{2}, \pi \right)$ ,

then  $dy/dx$  at  $x = 5\pi/6$  is:

$$x \in \left( \frac{\pi}{2}, \pi \right)$$

$$\frac{x}{2} \in \left( \frac{\pi}{4}, \frac{\pi}{2} \right)$$

☒ A.  $-1/2$

☐ B.  $-1$

☐ C.  $1/2$

☐ D.  $0$

$$y(x) = \cot^{-1} \left( \frac{\cancel{\cos \frac{x}{2}} + \cancel{\sin \frac{x}{2}} + \sin \frac{x}{2} - \cancel{\cos \frac{x}{2}}}{\cos \frac{x}{2} + \cancel{\sin \frac{x}{2}} - \cancel{\sin \frac{x}{2}} + \cos \frac{x}{2}} \right)$$

$$= \cot^{-1} \left( \frac{\cancel{2} \sin \frac{x}{2}}{\cancel{2} \cos \frac{x}{2}} \right)$$


$$= \cot^{-1} \left( \tan \frac{x}{2} \right)$$

$$= \frac{\pi}{2} - \tan^{-1} \left( \tan \frac{x}{2} \right) = \frac{\pi}{2} - \frac{x}{2}$$

JEE Main 2021

$$y(x) = \frac{\pi}{2} - \frac{x}{2}$$

$$y'(x) = -\frac{1}{2}$$







If  $f(x) = \sin\left(\cos^{-1}\left(\frac{1-2^{2x}}{1+2^{2x}}\right)\right)$  and its first derivative with respect to x is  $-b/a \log_e 2$  when  $x = 1$ , where a and b are integers, then the minimum value of  $|a^2 - b^2|$  is \_

$$f'(1) = -\frac{b}{a} \ln 2$$

$$f(x) = \sin \cos^{-1} \left( \frac{1 - (2^x)^2}{1 + (2^x)^2} \right)$$

$$2^x = \tan \theta$$

$$= \sin \cos^{-1} \left( \frac{1 - \tan^2 \theta}{1 + \tan^2 \theta} \right)$$

$$= \sin \cos^{-1} \cos 2\theta$$

$$= \sin 2\theta$$

$$= \frac{2 \tan \theta}{1 + \tan^2 \theta} = \frac{2 \cdot 2^x}{1 + (2^x)^2}$$

JEE Main 2021

$$f(x) = \frac{2^{x+1}}{1+4^x}$$

$$f'(x) = \frac{2^{x+1} \ln 2 (1+4^x) - (4^x \ln 4) (2^{x+1})}{(1+4^x)^2}$$

$$f'(1) = \frac{4 \ln 2 (5) - (4 \times 2 \ln 2) (4)}{5^2}$$

$$f'(1) = \frac{-12 \ln 2}{25} = -\frac{b}{a} \ln 2$$

$$\begin{aligned} b &= 12 \\ a &= 25 \end{aligned}$$

$$\begin{array}{r} a^2 - b^2 \\ 5 \overline{) 25} \\ \underline{-144} \\ 481 \end{array}$$





If  $y(\alpha) = \sqrt{2 \left( \frac{\tan \alpha + \cot \alpha}{1 + \tan^2 \alpha} \right) + \frac{1}{\sin^2 \alpha}}, \alpha \in \left( \frac{3\pi}{4}, \pi \right),$

then  $dy/d\alpha$  at  $\alpha = 5\pi/6$  is :

☒ A. 4

☐ B.  $4/3$

☒ C. -4

☐ D.  $-1/4$

$$y(\alpha) = \sqrt{2 \left( \frac{\frac{\sin \alpha}{\cos \alpha} + \frac{\cos \alpha}{\sin \alpha}}{\frac{1}{\cos^2 \alpha}} \right) + \frac{1}{\sin^2 \alpha}}$$

$$= \sqrt{2 \cot \alpha + 1 + \cot^2 \alpha}$$

$$= \sqrt{(\cot \alpha + 1)^2}$$

$$= |\cot \alpha + 1| = -\cot \alpha - 1$$

$$\alpha > \frac{3\pi}{4}$$
$$\cot \alpha < \cot \frac{3\pi}{4}$$

$$\cot \alpha < -1$$

$$\cot \alpha + 1 < 0$$


**JEE Main 2020**

$$\frac{dy}{d\alpha} = \operatorname{cosec}^2 \alpha$$

$$= \operatorname{cosec}^2 \left( \frac{5\pi}{6} \right)$$

$$= 4$$

↓ funca



$$\left(\frac{3\pi}{4}, \pi\right)$$

$$\cot \alpha < -1$$

$$\underline{1 + \cot \alpha < 0}$$



Let  $y = y(x)$  be a function of  $x$  satisfying  $y\sqrt{1-x^2} = k - x\sqrt{1-y^2}$  where  $k$  is a constant and  $y(1/2) = -1/4$ . Then  $dy/dx$  at  $x = 1/2$ , is equal to :

A.  $-\sqrt{5}/4$

☒ B.  $-\sqrt{5}/2$  ✓

C.  $2/\sqrt{5}$

D.  $\sqrt{5}/2$  ✓

$$y\sqrt{1-x^2} + x\sqrt{1-y^2} = k$$

$$x = \sin(A) \quad y = \sin(B)$$

$$\sin B \cos A + \sin A \cos B = k$$

$$\sin(A+B) = k$$

$$A + (B) = \underbrace{\sin^{-1}(k)}$$

$$\sin^{-1}x + \sin^{-1}y = c$$

JEE Main 2020



$$\sin^{-1}x + \sin^{-1}y = c$$

$$\frac{1}{\sqrt{1-x^2}} + \frac{1}{\sqrt{1-y^2}} \frac{dy}{dx} = 0$$

$$-\frac{dy}{dx} = \frac{-\sqrt{1-y^2}}{\sqrt{1-x^2}} = \frac{-\sqrt{1-\frac{1}{16}}}{\sqrt{1-\frac{1}{4}}} = \frac{-\sqrt{15}}{4} \times \frac{2}{\sqrt{3}} = \frac{-\sqrt{5}}{2}$$



# # Logarithmic Differentiation





# Logarithmic Differentiation



$$y = \underline{f_1(x)} \underline{f_2(x)} \underline{f_3(x)} \dots$$

OR

$$y = (\underline{f(x)})^{\underline{g(x)}}$$

$$\begin{aligned} &x^x \\ &x^{\ln x} \\ &x^{\sin x} \end{aligned}$$



$$f(0) = 1 \times 2 \times 3 \times \dots \times n = n!$$

If  $f(x) = (x+1)(x+2)(x+3) \dots (x+n)$  then  $f'(0)$  is

A.  $n!$

B.

$$\frac{n(n+1)}{2}$$

C.

$$(n!)(\ln n!)$$

☒ D.

$$n! \left( 1 + \frac{1}{2} + \frac{1}{3} + \frac{1}{4} + \dots + \frac{1}{n} \right)$$

$$\ln f(x) = \ln(x+1) + \ln(x+2) + \dots + \ln(x+n)$$

$$\frac{f'(x)}{f(x)} = \frac{1}{x+1} + \frac{1}{x+2} + \dots + \frac{1}{x+n}$$

$$\Rightarrow f'(x) = f(x) \left( \frac{1}{x+1} + \frac{1}{x+2} + \dots + \frac{1}{x+n} \right)$$

$$f'(0) = \underbrace{f(0)}_{n!} \left( \frac{1}{1} + \frac{1}{2} + \frac{1}{3} + \dots + \frac{1}{n} \right)$$



Find  $\frac{dy}{dx}$  for  $y = x^x$ .

$$y = x^x$$

$$\ln y = \ln x^x$$

$$\ln y = \underline{x} \underline{\ln x}$$

$$\frac{1}{y} y' = x \times \frac{1}{x} + \ln x \times 1$$

$$y' = y(1 + \ln x)$$

$$y' = x^x(1 + \ln x)$$





★

if  $y = \underline{2^{\log_2 x^{2x}}} + \left(\tan \frac{\pi x}{4}\right)^{\frac{4}{\pi x}}$  then  $\left.\frac{dy}{dx}\right|_{x=1}$  is

✓ A. 4

B. 5/2

C. 3

D. Not defined

$$y = \underbrace{x^{2x}}_u + \underbrace{\left(\tan \frac{\pi x}{4}\right)^{\frac{4}{\pi x}}}_v$$

$$y = u + v$$

$$\left.\frac{dy}{dx}\right|_{x=1} = \left.\frac{du}{dx}\right|_{x=1} + \left.\frac{dv}{dx}\right|_{x=1}$$

$$= 2 + 2$$

$$= (4)$$



$$u = x^{2x}$$

$$\ln u = 2(x \ln x)$$

$$\frac{1}{u} \frac{du}{dx} = 2(1 + \ln x)$$

$$\boxed{\frac{du}{dx} = 2x^{2x}(1 + \ln x)}$$

$$\left. \frac{du}{dx} \right|_{x=1} = 2 \times 1 \times 1$$

$$= (2)$$

$$v = \left( \tan \frac{\pi x}{4} \right)^{\frac{4}{\pi x}}$$

$$\ln v = \left( \frac{4}{\pi x} \right) \ln \left( \tan \left( \frac{\pi x}{4} \right) \right)$$

$$\frac{1}{v} \frac{dv}{dx} = \frac{4}{\pi} \left\{ \frac{1}{x} \frac{\sec^2 \left( \frac{\pi x}{4} \right)}{\tan \left( \frac{\pi x}{4} \right)} \times \frac{\pi}{4} + \underbrace{\ln \left( \tan \frac{\pi x}{4} \right)}_{\left( \frac{-1}{x^2} \right)} \right\}$$

$$\frac{dv}{dx} = \frac{4}{\pi} \boxed{\left( \tan \frac{\pi x}{4} \right)^{\frac{4}{\pi x}}} \left\{ \frac{1}{\tan \frac{\pi}{4}} \times \frac{\pi}{4} + 0 \right\}$$

$$= \frac{4}{\cancel{\pi}} \times 1 \times \frac{2}{1} \times \frac{\cancel{\pi}}{4}$$

$$= (2)$$





$$y = x^3 + 2 \sin x + 1$$

$$\underline{y = f(x)}$$

# Derivative of Implicit Functions



# Implicit functions

Example

i.  $x^2 + 2xy + y^3 = 4$

ii.  $x + y + \sin(xy) = 1$

$y \neq f(x)$





If  $x^2 + 2xy + y^3 = 4$ , find  $\frac{dy}{dx}$

2 x K  
2K (1)

X  $2x + 2(xy' + y) + 3y^2 y' = 0$

X  $2x + \underline{2xy'} + \underline{2y} + \underline{3y^2 y'} = 0$

✓  $y' = \frac{-(2x+2y)}{(2x+3y^2)}$

# NVstyle

$$\frac{dy}{dx} = \frac{-\frac{\partial f}{\partial x}}{\left(\frac{\partial f}{\partial y}\right)} = \frac{-(2x + 2y(1) + 0 + 0)}{(0 + 2x(1) + 3y^2 + 0)}$$



If  $y^5 + xy^2 + x^3 = 4x + 3$ , then find

$\frac{dy}{dx}$  at (2, 1)

$$f = y^5 + xy^2 + x^3 - 4x - 3$$

$$\begin{aligned}\frac{dy}{dx} &= - \frac{\frac{\partial f}{\partial x} \big|_{y=k}}{\frac{\partial f}{\partial y} \big|_{x=k}} = - \frac{(0 + y^2 + 3x^2 - 4)}{(5y^4 + x(2y))} \\ &= - \frac{(1 + 12 - 4)}{(5 + 4)} = \textcircled{-1}\end{aligned}$$





If  $y = \sqrt{\sin x + \sqrt{\sin x + \sqrt{\sin x + \dots \infty}}}$  find  $\frac{dy}{dx}$

$$y = \sqrt{\sin x + y}$$

$$y^2 = \sin x + y$$

$$2y \frac{dy}{dx} = \cos x + \frac{dy}{dx}$$

$$\frac{dy}{dx} = \frac{\cos x}{2y - 1}$$



If  $ax^2 + 2hxy + by^2 = 0$  then prove that

$$\frac{dy}{dx} = \frac{ax + hy}{hx + by} = \frac{y}{x}$$



# NVStyle

$$2x^3 + 55x^2y' + 2022xy^2 + y^3 = 0$$

→  $\boxed{\frac{dy}{dx} = \frac{y}{x}}$



If  $y = y(x)$  is an implicit function of  $x$  such that  $\log_e (x + y) = 4xy$ , then  $d^2y/dx^2$  at  $x = 0$  is equal to.

$$x + y = e^{4xy} \quad \text{--- (1)}$$

$$x=0 \quad y=1$$

$$x=0 \quad y=1 \quad y'=3$$

$$1 + y' = 1 \times 4 \times 1$$

$$\boxed{y' = 3}$$

$$1 + y' = e^{4xy} \times 4 \underbrace{(xy' + y)}_{\text{II}} \quad \text{--- (2)}$$



JEE Main 2021

$$y'' = 4 \left\{ e^{4xy} (\cancel{xy''} + y' + y') + (\cancel{xy' + y})^2 e^{4xy} \times 4 \right\}$$

$$y'' = 4 \left\{ 1 \times (6) + 1 \times 1 \times 4 \right\}$$

$$= \boxed{40}$$





If  $x \log_e (\log_e x) - x^2 + y^2 = 4$  ( $y > 0$ ), then  $dy/dx$  at  $x = e$  is equal to :

$$x=e \quad -e^2 + y^2 = 4 \Rightarrow y^2 = 4 + e^2 \Rightarrow \boxed{y = \sqrt{4+e^2}}$$

A.  $\frac{(1+2e)}{2\sqrt{4+e^2}}$

$$x \left( \frac{1}{\ln x} \times \frac{1}{x} \right) + \ln(\ln x) \times 1 - 2x + 2y y' = 0$$

☒ B.  $\frac{(2e-1)}{2\sqrt{4+e^2}}$

$$e \left( \frac{1}{1} \times \frac{1}{e} \right) + 0 - 2(e) + 2\sqrt{4+e^2} y' = 0$$

JEE Main 2019

C.  $\frac{(1+2e)}{\sqrt{4+e^2}}$

$$1 - 2e + 2\sqrt{4+e^2} y' = 0$$

D.  $\frac{e}{\sqrt{4+e^2}}$

$$y' = \frac{2e-1}{2\sqrt{4+e^2}}$$







For  $x > 1$ , if  $(2x)^{2y} = 4e^{2x-2y}$ , then  $(1 + \log_e 2x)^2 dy/dx$  is equal to :

Skip

A.  $\frac{x \log_e 2x - \log_e 2}{x}$

$$(2x)^{2y} = 2^2 e^{2x-2y}$$

B.  $\log_e 2x$

$$2y \ln(2x) = 2 \ln 2 + 2x - 2y$$

C.  $\frac{x \log_e 2x + \log_e 2}{x}$

$$x \left( y \left( \frac{2}{2x} \right) + \ln(2x) y' \right) - 1 + y' = 0$$

D.  $x \log_e 2x$

$$\frac{y}{x} + y' (\ln 2x) - 1 + y' = 0$$

$$x y (\ln 2x + 1) = x \ln 2 + x$$

$$y = \frac{\ln 2 + x}{\ln 2x + 1}$$

JEE Main 2019


$$y'(\ln 2x+1) = 1 - \frac{1}{x} \left( \frac{\ln 2+x}{\ln 2x+1} \right)$$

$$y'(\ln 2x+1)^2 = \frac{x(\ln 2x+1) - \ln 2 - x}{x}$$

$$= \frac{x \ln 2x - \ln 2}{x}$$



If  $e^y + \cancel{xy} = e$ , the ordered pair ( $dy/dx$ ,  $d^2y/dx^2$ ) at  $x = 0$  is equal to :

$x=0 \quad y=1$

☒ A.  $\left(\frac{1}{e}, -\frac{1}{e^2}\right)$

☒ B.  $\left(-\frac{1}{e}, \frac{1}{e^2}\right)$

☒ C.  $\left(\frac{1}{e}, \frac{1}{e^2}\right)$

☐ D.  $\left(-\frac{1}{e}, -\frac{1}{e^2}\right)$

JEE Main 2019

$$e^y y' + \cancel{xy'} + y = 0$$

$$e^1 y' + 0 + 1 = 0 \implies \boxed{y' = -\frac{1}{e}}$$

$$e^y y'' + (y')^2 e^y + \cancel{xy''} + y' + y' = 0$$

$$e^1 y'' + \frac{1}{e^2} \times e^1 + 2\left(-\frac{1}{e}\right) = 0$$

$$e y'' + \frac{1}{e} - \frac{2}{e} = 0$$

$$\boxed{y'' = \frac{1}{e^2}}$$





# # ☆☆ Derivatives Of Inverse Function



# Derivative of inverse function

$$f(x) = \text{given}$$

$$g(x) = f^{-1}(x)$$

$$\frac{d}{dx}(g(x)) = \frac{d}{dx}(f^{-1}(x))$$

Property

$$g(f(x)) = f(g(x)) = x$$

$$\downarrow \quad \downarrow$$
$$g(f(x)) = x$$

$$f(g(x)) = x$$

$$g'(f(x)) f'(x) = 1$$

$$f'(g(x)) g'(x) = 1$$

$$g'(f(x)) = \frac{1}{f'(x)}$$

$$f'(g(x)) = \frac{1}{g'(x)}$$



If  $f(x) = x^3 + x^5$  and  $g$  is the inverse of  $f$  find  $g'(2)$

$$f(1) = 1 + 1 = 2$$

$$g'(f(x)) = \frac{1}{f'(x)}$$

Put  $x=1$

$$g'(f(1)) = \frac{1}{f'(1)}$$

$$g'(2) = \frac{1}{8}$$

$$f(x) = x^3 + x^5$$

$$f'(x) = 3x^2 + 5x^4$$

$$f'(1) = 3 + 5 \\ = 8$$



Let  $f(x) = \exp(x^3 + x^2 + x)$  for any real number  $x$ , and let  $g$  be the inverse for  $f$ . The value of  $g'(e^3)$  is

☒ A.  $\frac{1}{6e^3}$

☐ B.  $\frac{1}{6}$

☐ C.  $\frac{1}{34e^{39}}$

☐ D. 6

$$f(x) = e^{x^3 + x^2 + x}$$

$$g'(e^3) = ?$$

$$g'(f(x)) = \frac{1}{f'(x)}$$

$$\boxed{x=1}$$

$$g'(f(1)) = \frac{1}{f'(1)}$$

$$\boxed{g'(e^3) = \frac{1}{6e^3}}$$

$$f(x) = e^{x^3 + x^2 + x}$$

$$f'(x) = e^{x^3 + x^2 + x} \times (3x^2 + 2x + 1)$$

$$f'(1) = e^3 \times 6 = \boxed{6e^3}$$







Let f and g be differentiable functions on  $\mathbb{R}$  such that fog is the identity function. If for some  $a, b \in \mathbb{R}$ ,  $g'(a) = 5$  and  $g(a) = b$ , then  $f'(b)$  is equal to :

✓ A.  $1/5$

B. 1

C. 5

D.  $2/5$

$$f(g(x)) = x$$

$$f'(g(x)) = \frac{1}{g'(x)}$$

$$x = a$$

$$f'(g(a)) = \frac{1}{g'(a)}$$

$$\boxed{f'(b) = \frac{1}{5}}$$

JEE Main 2020

If the function  $f(x) = x^3 + e^{\frac{x}{2}}$  and  $g(x) = f^{-1}(x)$ , then the value of  $g'(1)$  is  $f(0) = 0 + e^0 = 1$

$$g'(f(x)) = \frac{1}{f'(x)}$$

(JEE Adv. 2009)

Put  $x = 0$

$$g'(f(0)) = \frac{1}{f'(0)}$$

$$g'(1) = \frac{1}{(1/2)} = 2$$

$$f'(x) = 3x^2 + \frac{1}{2} e^{x/2}$$

$$f'(0) = 0 + \frac{1}{2} e^0$$

$$= 1/2$$



Let  $f: \mathbb{R} \rightarrow \mathbb{R}$ ,  $g: \mathbb{R} \rightarrow \mathbb{R}$  and  $h: \mathbb{R} \rightarrow \mathbb{R}$  be differentiable functions such that  $f(x) = x^3 + 3x + 2$ ,  $g(f(x)) = x$  and  $h(g(g(x))) = x$  for all  $x \in \mathbb{R}$ . Then

(JEE Adv. 2016)

~~(a)~~  $g'(2) = \frac{1}{15}$

$\checkmark$  (c)  $h(0) = 16$

~~(b)~~  $h'(1) = 666$

~~(d)~~  $h(g(3)) = 36$

$$g(f(x)) = x$$

$$g'(f(0)) = \frac{1}{f'(0)}$$

$$g'(2) = \frac{1}{3}$$

$$f'(x) = 3x^2 + 3$$

$$f'(0) = 3$$

$$f'(1) = 6$$

$$f'(6) = 3(6)^2 + 3$$

$$= \underline{\underline{111}}$$

$$h(g(g(x))) = x$$

$$x \rightarrow f(x)$$

$$h(g(g(f(x)))) = f(x)$$

$$h(g(x)) = f(x)$$

$$x \rightarrow f(x)$$

$$h(g(f(x))) = f(f(x))$$

$$h(x) = f(f(x))$$

$$g(f(x)) = x$$

$$h(0) = f(f(0))$$

$$= f(2)$$

$$= 2^3 + 3(2) + 2$$

$$= 16$$

$$\textcircled{b} \quad h'(x) = f'(f(x)) f'(x)$$

$$h'(1) = f'(f(1)) f'(1)$$

$$= f'(6) f'(1)$$

$$= 111 \times 6$$

$$= \underline{666}$$



$$h(g(x)) = f(x)$$

$$h(g(3)) = f(3) = 3^3 + 3(3) + 2$$

$$= 27 + 11$$

$$= \textcircled{38}$$



# Deducing new identities by using Differentiation





## Deduction of new identities by differentiating a given identity

# If  $\cos \frac{x}{2} \cdot \cos \frac{x}{2^2} \cdot \cos \frac{x}{2^3} \dots \cos \frac{x}{2^n} = \frac{\sin x}{2^n \sin \left( \frac{x}{2^n} \right)}$ , then prove that

$$\sum_{r=1}^n \frac{1}{2^r} \tan \frac{x}{2^r} = \frac{1}{2^n} \cot \frac{x}{2^n} - \cot x$$

$$\ln \cos \frac{x}{2} + \ln \cos \left( \frac{x}{2^2} \right) + \dots + \ln \cos \left( \frac{x}{2^n} \right) = \ln \sin x - \ln(2^n) - \ln \sin \left( \frac{x}{2^n} \right)$$

$$-\frac{\frac{1}{2} \sin \frac{x}{2}}{\cos \frac{x}{2}} - \frac{\frac{1}{2^2} \sin \left( \frac{x}{2^2} \right)}{\cos \left( \frac{x}{2^2} \right)} + \dots + -\frac{\frac{1}{2^n} \sin \left( \frac{x}{2^n} \right)}{\cos \left( \frac{x}{2^n} \right)} = \frac{\cos x}{\sin x} - \frac{\frac{1}{2^n} \cos \left( \frac{x}{2^n} \right)}{\sin \left( \frac{x}{2^n} \right)}$$

$$- \left\{ \frac{1}{2^1} \tan \frac{x}{2} + \frac{1}{2^2} \tan \frac{x}{2^2} + \dots + \frac{1}{2^n} \tan \left( \frac{x}{2^n} \right) \right\} = \cot x - \frac{1}{2^n} \cot \left( \frac{x}{2^n} \right)$$

$$\star \star \quad \boxed{\sum_{r=1}^n \frac{1}{2^r} \tan \left( \frac{x}{2^r} \right) = \frac{1}{2^n} \cot \left( \frac{x}{2^n} \right) - \cot x}$$





# # Standard Substitution



## Some standard substitutions

Expressions	Substitution
$\sqrt{a^2 - x^2}$	$x = a \sin \theta$ or $a \cos \theta$
$\sqrt{a^2 + x^2}$	$x = a \tan \theta$ or $a \cot \theta$
$\sqrt{x^2 - a^2}$	$x = a \sec \theta$ or $a \operatorname{cosec} \theta$
$\sqrt{\frac{a+x}{a-x}}$ or $\sqrt{\frac{a-x}{a+x}}$	$x = a \cos \theta$ or $a \cos 2\theta$

Root  
Hatao!

$$\begin{aligned}\sqrt{\frac{a+a\cos 2\theta}{a-a\cos 2\theta}} &= \sqrt{\frac{1+\cos 2\theta}{1-\cos 2\theta}} \\ &= \sqrt{\frac{2\cos^2 \theta}{2\sin^2 \theta}} \\ &= \cot \theta\end{aligned}$$



Find  $\frac{dy}{dx}$  for  $y = \tan^{-1} \left\{ \frac{1 - \cos x}{\sin x} \right\}$ , where  $-\pi < x < \pi$ .

$$y = \tan^{-1} \left\{ \frac{\cancel{x} \sin^{\cancel{2}} \frac{x}{2}}{\cancel{x} \sin \frac{x}{2} \cos \frac{x}{2}} \right\}$$


$$= \tan^{-1} \left( \tan \left[ \frac{x}{2} \right] \right)$$

$$y = \left[ \frac{x}{2} \right]$$

$$\frac{dy}{dx} = \frac{1}{2}$$



$$-\frac{\pi}{2} < \left( \frac{x}{2} \right) < \frac{\pi}{2}$$







$$y = \tan^{-1} \left( \frac{\sqrt{1+x^2}-1}{x} \right) \text{ Find } \frac{dy}{dx} \text{ at } x=1$$

A.  $1/2$

B.  $\sqrt{3}/2$

C.  $1$

☒ D.  $1/4$

$$x = \tan \theta$$

$$y = \tan^{-1} \left( \frac{\sec \theta - 1}{\tan \theta} \right)$$

$$= \tan^{-1} \left( \frac{1 - \cos \theta}{\sin \theta} \right)$$

$$= \tan^{-1} \left( \frac{2 \sin^2 \frac{\theta}{2}}{2 \sin \frac{\theta}{2} \cos \frac{\theta}{2}} \right)$$

$$y = \frac{\theta}{2}$$

MCM

$$\begin{aligned} \underline{M-1} \quad y &= \frac{1}{2} \tan^{-1} x \\ \frac{dy}{dx} &= \frac{1}{2} \times \frac{1}{1+x^2} = \left( \frac{1}{4} \right) \end{aligned}$$

$$\underline{M-2} \quad \frac{dy}{d\theta} = \frac{1}{2}$$

$$\frac{dx}{d\theta} = \sec^2 \theta$$

$$\begin{aligned} \frac{dy}{dx} &= \frac{1}{2(\sec^2 \theta)} \\ &= \frac{1}{2(1+x^2)} \end{aligned}$$







If  $f(x) = \sin^{-1} \left( \frac{2^{x+1}}{1 + 4^x} \right)$ , find  $f'(0)$ .

- ☒ A.  $\ln 2$
- ☐ B.  $\ln 4$
- ☐ C.  $-\ln 2$
- ☐ D.  $-\ln 4$

$$f(x) = \sin^{-1} \left( \frac{2 \cdot 2^x}{1 + (2^x)^2} \right)$$


$$= \sin^{-1} \left( \frac{2 \tan \theta}{1 + \tan^2 \theta} \right)$$

$$= \sin^{-1} (\sin 2\theta)$$

$$= 2\theta$$
$$f(x) = 2 \tan^{-1}(2^x)$$

$$2^x = \tan \theta$$

$$\tan^{-1}(2^x) = \theta$$


$$f'(x) = 2 \frac{1}{1+(2^x)^2} \times 2^x \ln 2$$

$$f'(0) = \cancel{2} \times \frac{1}{\cancel{2}} \times 1 \ln 2$$

$$= \ln 2$$

$$y = f(t)$$

# # Parametric Differentiation



# Parametric differentiation



M C M

$$x = at^2 \quad y = 2at \quad \frac{dy}{dx} = ?$$

$$\frac{dy}{dx} = \frac{\left(\frac{dy}{dt}\right)}{\left(\frac{dx}{dt}\right)} = \frac{2a}{2at} = \frac{1}{t}$$

↗  $\frac{d^2y}{dx^2} \neq \frac{-1}{t^2}$

$$\begin{aligned} \frac{d^2y}{dx^2} &= \frac{d}{dx} \left( \frac{dy}{dx} \right) = \boxed{\frac{d}{dt} \left( \frac{1}{t} \right)} \frac{dt}{dx} \quad (\# \text{ chالaki}) \\ &= \frac{-1}{t^2} \times \frac{1}{2at} = \frac{-1}{2at^3} \end{aligned}$$



Find  $\frac{dy}{dx}$  if  $x = a(\theta - \sin \theta)$  and  $y = a(1 - \cos \theta)$ .

☒ A.  $\cot\left(\frac{\theta}{2}\right)$

☐ B.  $\tan\left(\frac{\theta}{2}\right)$

☐ C.  $\sin\left(\frac{\theta}{2}\right)$

☐ D.  $\cos\left(\frac{\theta}{2}\right)$

$$\begin{aligned}\frac{dy}{dx} &= \frac{\frac{dy}{d\theta}}{\frac{dx}{d\theta}} = \frac{+a(\sin \theta)}{a(1 - \cos \theta)} = \frac{\cancel{2} \sin \frac{\theta}{2} \cos \frac{\theta}{2}}{\cancel{2} \sin^2 \frac{\theta}{2}} \\ &= \cot\left(\frac{\theta}{2}\right)\end{aligned}$$



If  $x = 3 \tan t$  and  $y = 3 \sec t$ , then the value of  $d^2y/dx^2$  at  $t = \pi/4$ , is :

eye 3)

A.  $1 / 3\sqrt{2}$

✓ B.  $1 / 6\sqrt{2}$

C.  $3 / 2\sqrt{2}$

D.  $1 / 6$

$$\frac{dy}{dx} = \frac{\frac{dy}{dt}}{\frac{dx}{dt}} = \frac{3 \sec t \tan t}{3 \sec^2 t} = \sin t$$

$$\frac{d^2y}{dx^2} = \frac{d}{dx} \left( \frac{dy}{dx} \right) = \frac{d}{dt} (\sin t) \left( \frac{dt}{dx} \right)$$

$$= (\cos t) \times \frac{1}{3 \sec^2 t}$$

$$= \frac{\cos^3 t}{3} = \frac{\left(\frac{1}{\sqrt{2}}\right)^3}{3} = \frac{1}{6\sqrt{2}}$$

JEE Main 2019







If  $x = 2\sin\theta - \sin 2\theta$  and  $y = 2\cos\theta - \cos 2\theta$ ,  $\theta \in [0, 2\pi]$ , then  $d^2y/dx^2$  at  $\theta = \pi$  is :

JEE Main 2020


A.  $3/4$

☒ B.  $3/8$

C.  $3/2$

D.  $-3/4$

$$\begin{aligned}\frac{dy}{dx} &= \frac{\left(\frac{dy}{d\theta}\right)}{\left(\frac{dx}{d\theta}\right)} = \frac{-\cancel{2}\sin\theta + \cancel{2}\sin 2\theta}{\cancel{2}\cos\theta - \cancel{2}\cos 2\theta} \\&= \frac{\sin 2\theta - \sin\theta}{\cos\theta - \cos 2\theta} \\&= \frac{\cancel{2}\cos\left(\frac{3\theta}{2}\right)\cancel{\sin\left(\frac{\theta}{2}\right)}}{\cancel{+2}\sin\left(\frac{3\theta}{2}\right)\cancel{\sin\left(\frac{\theta}{2}\right)}} \\&= \cot\left(\frac{3\theta}{2}\right)\end{aligned}$$


$$\frac{d^2y}{dx^2} = \frac{d}{dx} \left( \cot \frac{3\theta}{2} \right)$$

$$= \frac{d \left( \cot \frac{3\theta}{2} \right)}{d\theta} \frac{d\theta}{dx}$$

$$\left. \frac{d^2y}{dx^2} \right|_{\theta=\pi} = -\frac{3}{2} \underbrace{\operatorname{cosec}^2 \left( \frac{3\theta}{2} \right)}_{=1} \times \frac{1}{2\cos\theta - 2\cos 2\theta}$$

$$= -\frac{3}{2} \times 1 \times \frac{1}{2(-1) - 2(1)}$$

$$= \left( \frac{3}{8} \right)$$



# Differentiation  
of one function  
w.r.t other



## Differentiation of one function w.r.t. other function

$$\frac{dy}{dx}$$

$$\frac{d f(x)}{d g(x)} = \frac{\frac{d(f(x))}{dx}}{\frac{d(g(x))}{dx}} = \frac{f'(x)}{g'(x)}$$

$$\ln x \text{ w.r.t } \sin x \Rightarrow \frac{\frac{1}{x}}{\cos x}$$



Differentiate log sin x w.r.t  $\sqrt{\cos x}$

$$\Rightarrow \frac{\frac{1}{\sin x} \times \cos x}{\frac{1}{2\sqrt{\cos x}} \times -\sin x}$$





The derivative of  $\tan^{-1} \left( \frac{\sqrt{1+x^2}-1}{x} \right)$  with respect to  $\tan^{-1} \left( \frac{2x\sqrt{1-x^2}}{1-2x^2} \right)$  at  $x = \frac{1}{2}$  is :

A.  $2\sqrt{3} / 5$

B.  $\sqrt{3} / 12$

C.  $2\sqrt{3} / 3$

☒ D.  $\sqrt{3} / 10$

$x = \tan \theta$

$\Rightarrow \tan^{-1} \left( \frac{\sec \theta - 1}{\tan \theta} \right)$

$\Rightarrow \tan^{-1} \left( \frac{1 - \cos \theta}{\sin \theta} \right)$

$\Rightarrow \tan^{-1} \left( \tan \frac{\theta}{2} \right)$

$\Rightarrow \frac{\theta}{2}$

$\Rightarrow \boxed{\frac{\tan^{-1} x}{2}}$

$x = \sin \theta$

$\Rightarrow \tan^{-1} \left( \frac{2 \sin \theta \cos \theta}{\cos 2\theta} \right)$

$\Rightarrow \tan^{-1} (\tan 2\theta)$

$\Rightarrow 2\theta$

$\Rightarrow \underline{2 \sin^{-1} x}$

JEE Main 2020

$$\frac{\frac{1}{2} \times \frac{1}{1+x^2}}{2 \times \frac{1}{\sqrt{1-x^2}}} \quad \bigg| \quad x = \frac{1}{2}$$

$$\frac{\frac{1}{4} \times \frac{\sqrt{1-\frac{1}{4}}}{1+\frac{1}{4}}}{}$$

$$\frac{1}{4} \times \frac{\sqrt{3}}{2} \times \frac{1}{5} = \left( \frac{\sqrt{3}}{10} \right)$$





The derivative of  $\tan^{-1} \left( \frac{\sin x - \cos x}{\sin x + \cos x} \right)$ ,

with respect to  $x/2$ , where  $\left( x \in \left( 0, \frac{\pi}{2} \right) \right)$  is :

- A. 1  $\Rightarrow \tan^{-1} \left( \frac{\tan x - 1}{\tan x + 1} \right) \Rightarrow \left( \frac{x}{2} \right)$
- B.  $2/3 \Rightarrow \tan^{-1} \left( \tan \left( x - \frac{\pi}{4} \right) \right)$
- C.  $1/2 \Rightarrow \left( x - \frac{\pi}{4} \right)$
- ☒ D. 2  $\Rightarrow \frac{1}{\left( \frac{1}{2} \right)} = 2$

JEE Main 2019





Let  $f(\theta) = \sin\left(\tan^{-1}\left(\frac{\sin\theta}{\sqrt{\cos 2\theta}}\right)\right)$ , where  $-\frac{\pi}{4} < \theta < \frac{\pi}{4}$ .

$$\sqrt{\sin^2\theta + 1 - 2\sin^2\theta} = \sqrt{1 - \sin^2\theta} = \cos\theta$$

(JEE Adv. 2011)


Then the value of  $\frac{d}{d(\tan\theta)}(f(\theta))$  is

$f(\theta)$  wrt  $\tan\theta$

$$f(\theta) = \sin \sin^{-1}\left(\frac{\sin\theta}{\cos\theta}\right)$$

$$= \tan\theta$$

1





$$y' = \frac{dy}{dx} = y_1 = D_1(y) = f'(x)$$

$$y'' = \frac{d^2y}{dx^2} = y_2 = D_2(y) = f''(x)$$

# # Successive Differentiation





# Successive Differentiation

---





# Successive Differentiation

1.

$$\frac{dx}{dy} = \frac{+1}{\left(\frac{dy}{dx}\right)}$$

$$\frac{dx}{dy} = \frac{1}{\left(\frac{dy}{dx}\right)}$$

$$\frac{\Delta y}{\Delta x} = \frac{1}{\left(\frac{\Delta x}{\Delta y}\right)}$$

2.

$$\frac{d^2x}{dy^2} \neq \frac{1}{\frac{d^2y}{dx^2}}$$

$$\frac{d^2x}{dy^2} = \frac{d}{dy} \left( \frac{dx}{dy} \right)$$



## Successive Differentiation



3.

$$\frac{d^2y}{dx^2} = \frac{d}{dx} \left( \frac{dy}{dx} \right)$$

4.

$$\frac{d^2x}{dy^2} = \frac{d}{dy} \left( \frac{dx}{dy} \right)$$



$\frac{d^2x}{dy^2}$  equals:

$$y = f(x) \quad y' = f'(x)$$

A.  $-\left(\frac{d^2y}{dx^2}\right)^{-1} \left(\frac{dy}{dx}\right)^{-3}$

B.  $\left(\frac{d^2y}{dx^2}\right) \left(\frac{dy}{dx}\right)^{-2}$

☒ C.  $-\left(\frac{d^2y}{dx^2}\right) \left(\frac{dy}{dx}\right)^{-3}$

D.  $\left(\frac{d^2y}{dx^2}\right)^{-1}$

$$\frac{d^2x}{dy^2}$$

$$= \frac{d}{dy} \left( \frac{dx}{dy} \right)$$

$$= \boxed{\frac{d}{dx} \left( \frac{dy}{dx} \right)^{-1}} \frac{dx}{dy}$$

$$= -1 \left( \frac{dy}{dx} \right)^{-2} \times \frac{d^2y}{dx^2} \times \left( \frac{dy}{dx} \right)^{-1}$$

$$= - \left( \frac{dy}{dx} \right)^{-3} \left( \frac{d^2y}{dx^2} \right)$$

(JEE Adv. 2007)  
(JEE Main)





If  $y = \sin(\sin x)$  then prove that  $\underline{y_2} + (\tan x) \underline{y_1} + y \cos^2 x = 0$

$$y_1 = \boxed{\cos(\sin x)} \cos x$$

$$y_2 = \boxed{\cos(\sin x)} (-\sin x) + (\cos x)^2 (-\sin(\sin x))$$

$$y_2 = \left( \frac{y_1}{\cos x} \right) (-\sin x) + \cos^2 x (-y)$$

$$\boxed{y_2 + \tan x y_1 + \cos^2 x y = 0}$$





If  $y^2 + 2 \log_e (\cos x) = y$ ,  $x \in (-\pi/2, \pi/2)$ , then :

$$x=0 \quad y=0, 1$$

\*\*\*

☒ A.  $y''(0) = 0$

☒ B.  $|y'(0)| + |y''(0)| = 1$

☒ C.  $|y''(0)| = 2$

☒ D.  $|y'(0)| + |y''(0)| = 3$

$$y^2 + 0 = y$$

$$y^2 - y = 0$$

$$y = 0, 1$$

$$2yy' + 2 \frac{(-\sin x)}{\cos x} = y'$$

$x=0$

$$2yy' = y'$$

$$(2y-1)y' = 0$$

$\neq 0$

$$y' = 0$$

$$2(y y'' + (y')^2) - 2 \sec^2 x = y''$$

$$2(y y'' + 0) - 2 = y''$$

$$y'' = \frac{2}{2y-1}$$

$y=0 \rightarrow -2$

$y=1 \rightarrow +2$

$$|y''| = 2$$

JEE Main 2020





If  $f : S \rightarrow S$  where  $S = (0, \infty)$  be a twice differentiable function such that  $f(x+1) = xf(x)$  If  $g : S \rightarrow \mathbb{R}$  be defined as  $g(x) = \log_e f(x)$ , then the value of  $|g''(5) - g''(1)|$  is equal to :

A. 205/144

B. 197/144

C. 187/144

D. 1

JEE Main 2021



Let  $g(x) = \log f(x)$  where  $f(x)$  is twice differentiable positive function on  $(0, \infty)$  such that  $f(x+1) = x f(x)$ . Then, for  $N=1, 2, 3, \dots$

\*\*\*\* MOD

$$g''\left(N + \frac{1}{2}\right) - g''\left(\frac{1}{2}\right) =$$

(a)  $-4 \left\{ 1 + \frac{1}{9} + \frac{1}{25} + \dots + \frac{1}{(2N-1)^2} \right\}$

(b)  $4 \left\{ 1 + \frac{1}{9} + \frac{1}{25} + \dots + \frac{1}{(2N-1)^2} \right\}$

(c)  $-4 \left\{ 1 + \frac{1}{9} + \frac{1}{25} + \dots + \frac{1}{(2N+1)^2} \right\}$

(d)  $4 \left\{ 1 + \frac{1}{9} + \frac{1}{25} + \dots + \frac{1}{(2N+1)^2} \right\}$

$$\frac{f(x+1)}{f(x)} = x$$

$$\ln f(x+1) - \ln f(x) = \ln x$$

(JEE Adv. 2008)

$$g(x+1) - g(x) = \ln x$$

$$g'(x+1) - g'(x) = \frac{1}{x}$$

$$g''(x+1) - g''(x) = -\frac{1}{x^2}$$

$$x \rightarrow x - \frac{1}{2}$$

$$g''\left(x - \frac{1}{2} + 1\right) - g''\left(x - \frac{1}{2}\right) = \frac{-1}{\left(x - \frac{1}{2}\right)^2}$$

$$g''\left(x + \frac{1}{2}\right) - g''\left(x - \frac{1}{2}\right) = \frac{-4}{(2x-1)^2}$$

$$x=1 \quad \cancel{g''\left(\frac{3}{2}\right) - g''\left(\frac{1}{2}\right) = \frac{-4}{1^2}}$$

$$x=2 \quad \cancel{g''\left(\frac{5}{2}\right) - g''\left(\frac{3}{2}\right) = \frac{-4}{3^2}}$$

$$x=3 \quad \cancel{g''\left(\frac{7}{2}\right) - g''\left(\frac{5}{2}\right) = \frac{-4}{5^2}}$$

$$x=N \quad \cancel{g''\left(N + \frac{1}{2}\right) - g''\left(N - \frac{1}{2}\right) = \frac{-4}{(2N-1)^2}}$$

---


$$g''\left(N + \frac{1}{2}\right) - g''\left(\frac{1}{2}\right) = -4 \left( \frac{1}{1^2} + \frac{1}{3^2} + \frac{1}{5^2} \cdots + \frac{1}{(2N-1)^2} \right)$$







## Bonus Concept

(Odd)' = Even ✓

Even  $f(-x) = f(x)$

odd  $f(-x) = -f(x)$

$$f \rightarrow \text{Even}$$

$$f' \rightarrow \text{Odd}$$

(Even)' = Odd ✓

$$f(x) = x^4 \quad (\text{Even})$$

$$f'(x) = 4x^3 \quad (\text{odd})$$

$$f''(x) = 12x^2 \quad (\text{Even})$$

$$f'''(x) = 24x \quad (\text{odd})$$

$$f^{(4)}(x) = 24 \quad (\text{Even})$$

$$f(-x) = f(x)$$

$$f'(-x) \times (-1) = f'(x)$$

$$\underline{\underline{f'(-x) = -f'(x)}}$$


# Join with us in Telegram

## TELEGRAM CHANNEL

- [t.me/unacademyatoms](https://t.me/unacademyatoms) <sup>SN</sup> <sub>PDF</sub>

## COMPLETE NOTES AND LECTURES

- [livedaily.me/atoms](https://livedaily.me/atoms)



# + India's BEST Educators

# Unacademy Subscription

**BATCH**

HINDI

Igneous Batch for JEE Advanced & Olympiads 2021

Started on Jul 7

Nishant Vora and 1 more

**BATCH**

HINDI

Evolve Batch Course for Class 12th JEE Main and Advanced 2022

Starts on Apr 7

Anupam Gupta and 2 more

**BATCH**

HINDI

Mega Batch Course for Class 12th JEE Main and Advanced 2022

Starts on Apr 6

Narendra Avasthi and 1 more

**BATCH**

HINDI

Enthuse: Class 12th for JEE Main and Advanced 2022

Starts on Apr 14

Amarnath Anand and 2 more

**BATCH**

HINDI

Final Rapid Revision Batch for JEE Main 2021

Starts on Apr 6

Manoj Chauhan and 2 more

**LIVE**

HINDI PHYSICS

Course of 12th syllabus Physics for JEE Aspirants 2022: Part - I






Lesson 1 • Apr 2, 2021 12:30 PM

D C Pandey

If you want to be the **BEST**  
“Learn” from the **BEST**



A O D  
Last Min\*  
JEE Adv  
R B.C  
R R.B





NVLIVE

×

IIT JEE subscription

ICONIC <sup>+</sup> PLUS

✓ India's Best Educators

✓ Interactive Live Classes

✓ Structured Courses & PDFs

✓ Live Tests & Quizzes

✗ 1:1 Live Mentorship

✗ Live Doubt Solving

30 months

₹1,980/mo

No cost EMI

+10% OFF ₹59,399

>

24 months

₹2,310/mo

No cost EMI

+10% OFF ₹55,440

>

12 months

₹3,176/mo

No cost EMI

+10% OFF ₹38,115

>

6 months

₹4,620/mo

No cost EMI

+10% OFF ₹27,720

>

3 months

₹5,775/mo

+10% OFF ₹17,325

>

To be paid as a one-time payment

NVLIVE

×



×

IIT JEE subscription

PLUS ICONIC <sup>+</sup>

✓ India's Best Educators

✓ Interactive Live Classes

✓ Structured Courses & PDFs

✓ Live Tests & Quizzes

✓ Personal Coach

✓ Study Planner

24 months

₹3,750/mo

No cost EMI

+10% OFF ₹90,000

>

18 months

₹4,000/mo

No cost EMI

+10% OFF ₹72,000

>

12 months

₹4,875/mo

No cost EMI

+10% OFF ₹58,500

>

6 months

₹5,700/mo

+10% OFF ₹34,200

>

To be paid as a one-time payment

NVLIVE

×







# ICONIC





# PLUS


 **Personal Guidance**  
Get one on one guidance from top exam experts

 **Study Planner**  
Customized study plan with bi-weekly reviews


 Rec  
**Live Classes**


  
**Weekly Tests**

 (IT)  
**Structured Courses**

  
**Unlimited Access**

 **Test Analysis**  
Get one on one guidance from top exam experts

 **Study Material**  
Specialised Notes & Practice Sets

 **Experts' Guidelines**  
Study booster workshops by exam experts

V.C  
DCS  
M.T

# Upcoming Batches in December



**All Stars Crash Course 3.0** Batch for Class 11th JEE Main and Advanced 2023



Starts on **1st Dec. 2021**

**Rapid revision** for JEE Main and Advanced 2022



Starts on **8th Dec. 2021**

**Edge for JEE Main and Advanced 2023** - Early class 12th



Starts on **15th Dec. 2021**

**Rapid revision** for JEE Main and Advanced 2022 - 2.0



Starts on **15th Dec. 2021**

**Striker Batch: JEE Main 2021**



Starts on **15th Dec. 2021**

**Excellence** for JEE Main and Advanced 2023



Starts on **22nd Dec. 2021**

**Striker Batch: JEE Main 2021 - 2.0**



Starts on **22nd Dec. 2021**

**Edge for JEE Main and Advanced 2023** - Early class 12th - 2.0




Starts on **29th Dec. 2021**


**Excellence** for JEE Main and Advanced 2023 - 2.0



Starts on **5th Jan. 2022**







ALL INDIA MOCK TESTS  
FOR JEE 2022

# All India Mock Test Series for JEE 2022

Starting Dec 25 | 2-5 PM



All India IIT JEE Rank Predictor



Curated questions by top educators & Video Solution to each Question



Detailed test report to identify areas of strength and improvement

**NVLIVE**

Enroll Now

USE CODE -





# Unacademy Atom Test Series



## Weekly Mock Test

- **28<sup>th</sup> December**  
**8:00 PM**

**Enroll Now**

Use code:





UNACADEMY  
CHAMPIONS  
LEAGUE

# Ace UCL for JEE

Stand a chance to meet and get inspired by  
**Sachin Tendulkar**

Aptitude Test on Dec 18, 19, 26 | 1 PM Onwards

**Enroll Now**

Use Invite  
Code





UNACADEMY  
CHAMPIONS  
LEAGUE

# Unacademy Champions League

Is Back, once again for You !!

## Participate and Win!! | Mega Prizes



Apple MacBook Air



Apple watch



Airpod



13 Participants win  
Amazon echo dot



60 Participants to  
get 1000 AGV

Winners will be decided based on lucky draw of all  
participants of December 26th test

[Enroll Now](#)

\*Plus subscribers are not eligible





# IIT-JEE COMBAT

ANDROID  
APP AND WEB

LIVE Test

21 Questions

60 Minutes

26 DECEMBER

11 : 00 AM

- ★ Practice IIT-JEE Quality Questions
- ★ Based on the Recent Syllabus of IIT JEE Mains & Advance
- ★ Questions & Video Solutions by Top Educators
- ★ Separate test for Class 11, 12 and 13
- ★ Top 25 Rankers to get Merit Certificate

Rewards from a pool of ₹4 CRORE +



RANK 1 : GOLD  
MEDAL + MERIT  
CERTIFICATE+  
100% SCHOLARSHIP



RANK 2 : SILVER  
MEDAL + MERIT  
CERTIFICATE+  
100% SCHOLARSHIP



RANK 3 : BRONZE  
MEDAL + MERIT  
CERTIFICATE+  
100% SCHOLARSHIP


AND MANY MORE...



I-PAD



JBL Headphones



JBL Speakers



Amazon Vouchers

Enroll Now

250 lucky participants to win Exciting Rewards!

\*T&C applied

# Thank You!

+ SUBSCRIBE

