

Commonly found disease in Kanchanpur

POTATO DISEASE :- LATE BLIGHT

(Pachaute Dhadhuwa) RAPID



(Soil borne fungus type organism) bata lagne disease ho "Phytophthora infestans".
Can survive in soil under adverse conditions for several years. (Spreads through Spores)

GROWING / SPREADING CONDITIONS:-
wet, foggy, frosty jado/ pani parne mahina poorly drained fields where standing water

persists for several days.



SYMPTOMS :- distinctive odor (spots in leaves stems and tubers)

DIRECT CONTROLS :- Once symptoms are seen, nothing much can be done to stop without chemicals. Stopping blight is all about precautions.

KUN PLANTS Asar garxata ?
Commonly in Tomatoes cucumber and

soybeans.

HUMAN LAI ASAR?? :- Direct disease le asar pardaina but tissue damgele aru problem create garna sakxa.

Imp POINTS :- damaged potatoes should scritcly NOT be used in making compost.

KANCHANPUR CASE:-

KAILALI, Jan 8: Winter rains and cold waves, as well as the outbreak of Dadhuwa disease (blight), have damaged potato crop in Kailali and Kanchanpur district.

Farmers in Gauriganga Municipality-7 are facing significant losses due to a blight outbreak affecting their potato and tomato crops, worsened by cold waves and winter rains. Chandra Dev Joshi and Khushi Ram Chaudhary reported that even the resilient Tharu variety has been severely impacted,

forcing them to consider buying vegetables for their daily needs.

SOURCE :- <https://www.google.com/amp/myrepublica.nagariknetwork.com/amp/blight-cold-wave-damage-potato-plants-in-kailali-kanchanpur/>