

Daniel **Fitz**  
(43961229)



University Of Queensland

**STAT2203** – Probability Models and Data Analysis for Engineering

STAT2203 Assignment 2

## Question 1

Let  $X$  be the event that the ball falls in box  $x$ . Where  $X \sim \text{Bin}(4, 0.5)$

### SubQuestion A

$$\begin{aligned}\mathbb{P}(X = x) &= \binom{4}{x} \left(\frac{1}{2}\right)^x \times \left(1 - \frac{1}{2}\right)^{4-x} \\ &= \binom{4}{x} \left(\frac{1}{2}\right)^4 \\ &= \frac{1}{16} \binom{4}{x} \\ \mathbb{P}(X = 3) &= \frac{1}{16} \binom{4}{3} \\ &= \frac{4}{16} = \frac{1}{4}\end{aligned}$$

### SubQuestion B

$$\begin{aligned}\mathbb{P}(X = x) &= \binom{4}{x} \left(\frac{1}{2}\right)^x \times \left(1 - \frac{1}{2}\right)^{4-x} \\ &= \binom{4}{x} \frac{1}{2^x} \times \frac{1}{2^{4-x}} \\ &= \binom{4}{x} \frac{2^x}{2^4 2^x} \\ &= \frac{1}{16} \binom{4}{x}\end{aligned}$$

## Question 2

### Part 1

We are given:

$$\begin{aligned}\mathbb{P}(X = 1) &= r \\ \mathbb{P}(X = 0) &= 1 - r \\ \mathbb{P}(Y = 1) &= s \\ \mathbb{P}(Y = 0) &= 1 - s\end{aligned}$$

Therefore the following table can be generated:

$Y \backslash X$	0	1
0	$(1-r)(1-s)$	$r(1-s)$
1	$s(1-r)$	$rs$

Table 1: Combinations of  $P(X=x)P(Y=y)$

Using the above table, we can see that  $U$  and  $V$  will both take on values  $\{0, 1\}$

$U, V$	$X, Y$	Solution
0,0	0,0	$(1-r)(1-s)$
0,1	1,0 + 0,1	$r(1-s) + s(1-r)$
1,0	Not possible	0
1,1	1,1	$rs$

Therefore the following joint pmf table can be calculated

$U \backslash V$	0	1
------------------	---	---

0	$(1-r)(1-s)$	$r(1-s) + s(1-r)$
1	0	$rs$

## Part 2

$$\mathbb{P}(V = 1, U = 1) = \mathbb{P}(V = 1)\mathbb{P}(U = 1)$$

$$rs = (rs)(r + s - rs)$$

$$1 = r + s - rs$$

$$1 - s = r - rs$$

$$= r(1 - s)$$

$$1 = r$$

$$\mathbb{P}(V = 0, U = 0) = \mathbb{P}(V = 0)\mathbb{P}(U = 0)$$

$$(1 - r)(1 - s) = (1 - rs)(1 - s - r + rs)$$

$$1 - r - s + rs = 1 - s - r + rs - rs + rs^2 + sr^2 + (rs)^2$$

$$1 + rs = 1 + rs^2 + sr^2 - (rs)^2$$

$$1 + s = 1 + s^2 + s - s^2 \quad (\text{When } r \text{ is } 1)$$

$$1 + s = 1 + s$$

$$\mathbb{P}(V = 1, U = 0) = \mathbb{P}(U = 0)\mathbb{P}(V = 1)$$

$$r + s - 2rs = (1 - rs)(r + s - rs)$$

$$r + s - 2rs = r + s - rs - r^2s - rs^2 + r^2s^2$$

$$rs = -r^2s - rs^2 + r^2s^2$$

$$s = -s - s^2 + s^2 \quad (\text{When } r \text{ is } 1)$$

$$s = -s$$

$$s = 0$$

$$\mathbb{P}(U = 1, V = 0) = \mathbb{P}(U = 1)\mathbb{P}(V = 0)$$

$$0 = rs(1 - s - r + rs) \quad (\text{When } r \text{ is } 1, \text{ When } s \text{ is } 0)$$

$$0 = 0$$

---


## Question 3

$$\begin{aligned}L(\theta; p) &= \prod_{i=1}^5 (1-p)^{x_i} p \\l(\theta; p) &= \sum_{i=1}^5 \log((1-p)^{x_i-1}) \log(p) \\&= n \log(p) + \sum_{i=1}^5 (x_i - 1) \log(1-p) \\&= n \log(p) + \log(1-p) \sum_{i=1}^5 (x_i - 1) \\\frac{d}{dp} \frac{n}{p} - \frac{\sum_{i=1}^5 (x_i - 1)}{1-p} \\0 &= \frac{n}{p} - \frac{\sum_{i=1}^5 (x_i - 1)}{1-p} \\\frac{n}{p} &= \frac{\sum_{i=1}^5 (x_i - 1)}{1-p} \\\frac{n}{p} - n &= -n \sum_{i=1}^5 (x_i) \\\frac{n}{p} &= \sum_{i=1}^5 (x_i) \\p &= \frac{n}{\sum_{i=1}^5 (x_i)} = \frac{1}{\bar{x}}\end{aligned}$$

---

## Question 4

```
1 function result = ass2q1
2     N=1e4;
3     results = 1:N;
4     for i = 1:N;
5         results(i) = drop_ball(0, 0);
6     endfor
7     hist(results, 0:4);
8     xlabel("Ball Fall Position");
9     ylabel("Number of balls in position");
10 endfunction
11
12 function result = drop_ball(level, position)
13     if (level >= 4)
14         result = position;
15     else
16         direction = rand > 0.5;
17         result = drop_ball(level + 1, position + direction);
18     endif
19 endfunction
```



*Figure 1: Result for Question 4*

---

## Question 5

---

```
1 function result = ass2q2
2     N = 1e2;
3     totals = zeros(N, N);
4     for i = 1:N
5         for j = 1:N
6             totals(i, j) = calcRS(i/N, j/N);
7         endfor
8     endfor
9     mesh(1:N, 1:N, totals);
10    title("Question 5");
11    xlabel("Value for r");
12    ylabel("Value for s");
13    zlabel("Resulting probability");
14 endfunction

15
16 function result = calcRS(r, s)
17     result = sumRS(0, 0, r, s) + sumRS(0, 1, r, s) + sumRS(1, 0, r, s) +
18         sumRS(1, 1, r, s);
19 endfunction

20
21 function result = sumRS(u, v, r, s)
22     result = abs(probUV(u, v, r, s) - (probU(u, r, s) * probV(v, r, s)));
23 endfunction

24
25 function result = probUV(u, v, r, s)
26     if (u == 1 && v == 1)
27         result = r * s;
28     elseif (u == 0 && v == 0)
29         result = (1 - r) * (1 - s);
30     elseif (u == 0 && v == 1)
31         result = r(1-s) + s(1-r);
32     elseif (u == 1 && v == 0)
33         result = 0;
34     else
35         # Should not reach this
36         result = -1;
37     endif;
38 endfunction

39
40 function result = probU(u, r, s)
41     result = probUV(u, 0, r, s) + probUV(u, 1, r, s);
42 endfunction

43
44 function result = probV(v, r, s)
45     result = probUV(0, v, r, s) + probUV(1, v, r, s);
46 endfunction
```

### Question 5





Figure 2: Result for Question 5

### Question 6

```

1 function result = ass2q6
2     N=5e4;
3     totals=1:N;
4     for i = 1:N
5         totals(i) = pmf(randi(5));
6     endfor
7     hist(totals, 0:0.1:0.4);
8     xlabel("pmf result");
9     ylabel("Number of results");
10 endfunction
11
12 function result = pmf(x)
13     result = (1/3) * (2/3)^(x - 1);
14 endfunction

```



*Figure 3: Result for Question 6*