

Ideas Parked — Not Yet Merged (v9)

Staging document; each entry marks its eventual destination

This document holds ideas discussed against Pattern Arithmetic v15_19, Inference, and Assembly, none of which have been written into any of those three files yet. Each entry states its status and its intended destination. Nothing here is settled; the point of this file is to let that be true without leaving Paper 1 mid-edit. v5 added one scribble: an identity handle on an atom as a meta field, for an honest “uncollapse” that looks up a kept bowl rather than inverting Mix. v6 added two: a sphere-packing bound relevant to the path-reduction item’s distance requirement, and a new theoretical problem — merging every atom in a unit space to one representative — which splits the same way the τ_{weather} correction already did. v7 added one scribble: a scoping refinement to Route 1 of Gap 1. v8 added a proposed terminology realignment — Collapse, UnCollapse, Bundle, MixMerge, UnMix — together with a precise cost assessment of merging it into Paper 1, flagging $\ominus$ as a direct symbol collision with the existing Axiom 4. v9 resolves that collision and one more: a checked, collision-free symbol for Bundle as a verb ($\oplus$) and for UnCollapse ($\ominus$), plus a precise account of exactly when ids are needed for UnCollapse to be identity-exact rather than merely value-exact. Nothing in this section is written into v15_19.

1 Destined for Pattern Arithmetic (still formally open — but a leading resolution candidate has emerged, with a citation)

Mix across differing directions. The paragraph’s own text still reads “left open.” Nothing here changes that text. What changed is that the paper already contains a worked, concrete instance of this exact question, in a section not previously connected to it.

[**Found: the paper already answers this question once, for one sort, and points toward a general answer.**] The `chladni` inclusion states, word for word: “Across modes the map is not available, and not for want of effort. Mode space is not closed under addition... Cross-mode coexistence is therefore $\oplus$ — a bowl of modes — and the resulting figure is a reading of that bowl which is not $\boxplus$. This is the open question of [the differing-directions open item] met in a concrete domain, and it suggests the question may be closed negatively for this sort rather than left waiting.” This is a citation, not a new argument — the paper’s own words, in a section that was not previously read as bearing on Gap 1 directly.

[**Why this converges with everything else worked through in this project, not just `chladni`.**] Once Mix is read as “resolve several reports on one shared question into a verdict” (argued in v2/v3, carried forward below unchanged), refusing across-direction combination stops looking like a gap and starts looking like the correct behavior of an operator that was never asked that question in the first place. And every practical workaround that actually worked in this project was already living on the $\oplus$ side of precisely this line, independently: bowls holding separate weather channels, per-channel similarity scoring, the candlestick summary for reference-atom building, Assembly’s own camp-to-system Hill bijection. None of them ever needed Mix to

cross a direction. All four arrived, separately, at “keep it in a bowl, read it there” — which the chladni section had already stated as a formal result.

[Status: a leading candidate, not yet a decision.] This supports Route 1 (refuse outright, from the three already on file) over Route 2 (rotate first) or Route 3 (two-stage operator) as the likely resolution, with a real citation behind it rather than preference. Writing this into Paper 1 would mean editing the “Open: Mix across differing directions” paragraph to state the refusal explicitly, citing the chladni section as precedent. Not done here, on request — parked, ready, pending a decision to merge.

[Carried forward unchanged: the reading argued in v2/v3.] Mix is not “multiple vectors becoming one vector.” It takes several reports already agreed to answer one shared question and returns a verdict about that question. The shared $\hat{n}$ is the precondition that makes “these answer the same thing” true, not a boundary condition on a more general operator. Supporting evidence: same- direction opposite-phase cancellation to black (principled contradiction-handling, not a fabricated compromise); unmixing’s proven ill-posedness, with metamerism as independent real-world precedent; the paper’s own opening line, “two notes yield a chord, not a larger integer... the word ‘add’ names a different rule.”

[Carried forward unchanged: the “we mix all the time” challenge, resolved.] Sugar in tea is one dial, never needed differing directions. Red plus green additively sits “at the same place” in the paper’s own words — already inside Mix’s settled domain. Paint-layer flattening is confirmed, across every source checked, as destructive and irreversible, matching unmixing exactly; compositing unflattened layers is $\oplus$, not $\boxplus$, which is why “unmerge them later” was always possible. Representing a whole painting as one vector remains Gap 1 itself, not a counterexample to it.

[Carried forward unchanged: the two-part split, now read as one part likely answered.] Embedding validity — does geometric closeness on a shared sphere track real semantic agreement at all — was the harder, prior question, true for hue by deliberate construction, unstated for an arbitrary sphere of concepts. Given the chladni precedent above, the likely answer for sorts where this was never stated is that no valid embedding needs constructing, because cross-direction combination is refused before the embedding question would even matter.

[Carried forward unchanged: still confirmed not to help.] Magnitude self-bounding near opposition does not license the direction that comes with it. The Hill form and Assembly’s Hill bijection both fix magnitude only, verified twice, never direction.

2 A correction, recorded so it is not repeated

τ_{weather} **names two different problems, and only one of them is Gap 1.** Raised once as a proposed admissible sort for combining different weather concepts (rain, drought) across directions — that question is genuinely Gap 1, and the discussion above bears on it. Raised a second time, later, meaning something narrower: given several already closely-aligned vectors within one bounded region, how to arrive at a single representative atom, using the Hill form. That second question does not invoke $\boxplus$ across differing directions at all — it is Assembly’s already-verified system-promotion mechanism (16 camps to 13 systems under the Hill bijection), recognized in a new setting, not a Paper 1 admissibility question. The two were briefly conflated in this project; recorded here so the same conflation is not repeated. Assembly’s version needs nothing further; Gap 1’s version is addressed in Section 1 above.

3 Destined for Pattern Arithmetic (advances an existing open item)

Location as a sort — a third candidate. The existing paragraph names two candidates and adopts neither. A third was tested here: dedicate separate, disjoint dimensions to content, to time, and to location, rather than letting any of them share an axis.

[**Verified.**] Two shifts acting through disjoint content dimensions commute exactly (checked numerically, not approximately: time-then-location and location-then-time landed on the identical point to full floating-point precision). Two shifts forced to share a single content dimension do not commute, confirming the disjointness is what matters, not rotation itself.

[**Verified.**] Content stays recoverable after both shifts: projecting back onto the content dimensions alone reproduced the original direction closely, while the time-axis and location-axis each carried a clean, separately-readable amount of shift, with nothing blended between them.

[**Open, unchanged.**] The encoding itself — what specific rotation a given time value or location corresponds to — is not stated anywhere. The geometry now supports it cleanly; the mapping does not exist yet.

This is real, checkable content, not yet written as an addition to the existing paragraph.

4 Destined for Pattern Arithmetic (open item stays open, unchanged)

Reducing a path by merging close actors. The existing paragraph asks for three things: a stated distance, a merge rule, and a bound on how far a reduced path’s endpoint may drift from the unreduced one. Discussion here supplied a distance (see below) and partially addressed the merge rule, but only for magnitude, not for direction — and direction is the harder half this item is actually about.

[**Verified.**] Switching the strength calculation from the companion paper’s clip to Inference’s already-established Hill form, $r' = |z|/(K + |z|)$, changes nothing about the resulting direction $\chi' = \arg(z)$. Checked directly on two constructed sequences with identical content and opposite order: both gave the same angle under either formula. The Hill substitution fixes evidence-count blindness in strength; it does not touch the order-blindness this open item is actually asking about.

[**New: a candidate grounding for the stated distance, item (1) of the three things required.**] How many actors can be mutually distinguishable — separated by at least some angle $\theta_{\min}$ — on $\mathcal{S}$ at all is a real, named problem (spherical codes, the Tammes problem), not an arbitrary choice to make. Exact optimal counts are known only for small N ; a standard approximate bound, treating each point as owning a non-overlapping cap of half-angle $\theta_{\min}/2$, gives

$$N_{\max} \approx \frac{4\pi}{2\pi(1 - \cos(\theta_{\min}/2))} = \frac{2}{1 - \cos(\theta_{\min}/2)}.$$

Computed for illustration:

$\theta_{\min}$	approx. max. distinct points
1	52,525
5	2,101
10	526
30	59
60	15
90	7
120	4

This is an approximation, not an exact extremal result, but it is a real, checkable mathematical object rather than a guess: naming a $\theta_{\min}$ for “close enough to merge” directly bounds how many genuinely distinct actors can exist near each other at all, which bears on item (3), the drift bound, as well as item (1).

[**Open, unchanged.**] No merge rule for direction exists. No drift bound exists for either magnitude or direction. This item remains exactly as open as it was before this discussion, now with the magnitude half more precisely separated from the direction half, and a real candidate supplied for the distance half specifically.

5 Destined for Inference (new mechanism)

Candlestick summary: a richer merge rule for reference-atom building. Not a resolution of the Paper 1 item above — a different question. Paper 1’s open item is about *walking*: can a shortened path still be walked to roughly the same endpoint. This is about *reading*: how to summarize a group of atoms for comparison against a reference atom, which is Inference’s own established business (its reference-atom section), not *T*’s.

[**Verified.**] Open/close/high/low, computed the way a trading candle is, distinguishes exactly the case flat Mix could not. Two sequences, identical content, opposite order — naive Mix gave the identical angle (47.5) for both. The candle’s open and close swapped between them ($20 \rightarrow 160$ versus $160 \rightarrow 20$), correctly reading one as ending badly and the other as ending well. High and low stayed identical for both, correctly capturing that only the *range* was shared, not the *direction of travel*.

[**Settled, matches existing machinery.**] The scope rule — every atom belongs to exactly one time unit, no overlap between buckets — is not new; it is precisely the discrete-time-bucket metadata already defined on a walk’s own steps in the companion paper.

[**Open.**] This gives Inference a genuinely better reference-atom merge rule than flat Mix. It is not proposed as a fix for Paper 1’s walking-specific open item, which remains untouched above. Whether open/close/high/low itself needs a coherence check analogous to $\bar{R}_c$ has not been considered.

6 Destined for Inference (naming caution, not yet a mechanism)

“Inference Mandelbrot” is not Assembly’s Mandelbrot, and should not share its name. Assembly’s multi-zoom promotes an unordered thing (a settled camp) and is fully handled by the Hill bijection already verified there. Whatever would promote an *ordered* thing (a walked path, a movie, a time series) into one higher-level atom is a different, harder problem — it is the direction half of the path-reduction item above, restated from a different angle. [**Naming flag only.**] If this is ever built, it wants its own name; reusing “Mandelbrot” for it would be the same kind of collision already caught and corrected for “collapse,” “singularity,” and the quantum swap test.

7 Destined across three papers (new, split by piece)

Meta vectors: a second kind of vector alongside information vectors. Proposed: an information atom may carry two distinct kinds of vector. Information vectors are Axiom 7’s bundle v_i ’s, capped at $r_i \leq 1$, potentially several sorts τ across a whole record. Meta vectors describe the atom rather than being part of its content — three named so far: V_{derived} , link vectors (pointers between atoms, one- or two-directional), and a timestamp vector. Explicitly left open whether more meta-vector types are needed. Meta vectors are not capped at 1, and grouping several atoms’ meta vectors together on like terms is proposed as a distinct mechanism for building higher structures. This is one idea that lands in three different places once traced to where each piece actually belongs.

[**Destined for Pattern Arithmetic.**] The taxonomy itself — information vectors versus meta vectors as a formal, named distinction sitting on top of Axiom 7 — is structural, the same kind of addition Axiom 7 itself was. V_{derived} already lives there. A timestamp vector, if adopted, would be a third, genuinely different treatment of time from the two already on file above: distinct from the discrete-bucket metadata already defined on a walk’s own step, and distinct from the dedicated-axis encoding in the location-as-a-sort entry. Not yet clear whether more than one of the three should coexist, or which.

[**Destined for Assembly.**] Link vectors specifically. This is the mechanism the shelf’s own neighbour-graph note was already naming without a concrete form: “a data structure on seats after rest, not T .” Grouping link vectors on like terms is collecting edges into a graph, not merging them — gathering, in the $\oplus$ sense — so it carries none of the open questions below it.

[**Destined for Inference.**] The grouping mechanism itself: querying several atoms’ meta vectors, confirming they sit close together, and promoting one shared meta-vector up to the group as a whole. This is the space-time gating idea from earlier discussion, generalized into a standing principle. Timestamp and location get *used* here; they are only *defined* in Pattern Arithmetic above.

[**Verified, carried forward as a caveat rather than a new finding.**] Combining several atoms’ V_{derived} ’s stays safe only under the constraint the original already carries: auxiliary, never promoted to a trusted, walkable atom on its own. Checked directly that neither Inference’s Hill form nor Assembly’s Hill bijection touches this constraint — both fix magnitude only, never direction — so grouping V_{derived} ’s inherits that same limit rather than escaping it.

[**Note added here.**] Link vectors were separately discussed as one possible mechanism for a limited, honest form of “uncollapse”: not inverting Mix (still impossible, unaffected by anything here), but keeping an explicit, external pointer from a collapsed atom back to the bowl that produced it, created at collapse time, before the bowl itself might otherwise be discarded. This is navigation via a kept record, not reconstruction from the collapsed result — the distinction the white-light/prism discussion turned on, confirmed there to matter.

8 Destined across three papers (scribble, v5)

Identity handle as a meta field, for honest uncollapse. An information atom may carry one more meta quantity: an opaque identity id , assigned before any gather or Mix, and kept on the atom afterwards. The scribble that prompted this note called it a “meta vector” with τ_{id} equal to that identity. Fine-tuning below refuses τ_{id} as a sort and keeps id as a *field*, in the same shelf as timestamp, mint-location, and link pointers already named in the meta-vector entry.

[**What it is for.**] Mix remains irreversible (Paper 1 v1.1). “Uncollapse” here does not reconstruct ingredients from a yellow. It means: at collapse time, write a pointer from the resulting

atom (or from the register occupant after a walk) back to the bowl that produced it, and keep that bowl under the same *id* (or a pair *result_id* $\leftrightarrow$ *bowl_id*). Later, given the result, look the bowl up. Navigation by a kept record, not inversion. This is the same distinction already recorded under link vectors and the white-light / prism discussion.

[Viability — what works.] An opaque handle is the right shape. It does not need to live on $\mathcal{S}$, does not need $r \leq 1$, and does not need a direction. A string or integer key is enough. Creating the pointer *before* the bowl is discarded is the whole trick; after discard there is nothing to uncollapse. Links already named that graph; *id* is the node name those edges need. Assembly can own the edge table (neighbour / parent-bowl). Paper 1 only needs the taxonomy: *id* is meta, not a unit of truth. Inference may use *id* to re-open a bowl and run a second reading.

[Viability — what to refuse now.] Do not mint τ_{id} . τ is the legality tag of an information vector. An identity is not a pattern and must not appear in the type table. Do not encode *id* as an angle on the sphere; colliding two ids geometrically would be a false Mix. Do not claim that $\{A\} \ominus A$ or T^{-1} plus *id* inverts $\boxplus$. Gather still satisfies $A \oplus A = A$: two tokens of the same pin share one address and, if they are the same *instance*, one *id*; two distinct instances of the same pin (two physical chairs) need two ids and are not collapsed by $\oplus$ unless the lexicon says they are the same state. That instance-versus-kind cut is the fine-tuning this scribble does not finish.

[Tiny picture.] Bowl $B = \{\text{red}, \text{green}\}$, $id_B = \#17$. Mix reads yellow, $id_Y = \#18$, meta link $\#18 \rightarrow \#17$. Uncollapse($\#18$) returns B , still red and green, still two atoms. The yellow is not pulled apart. The prism was kept on the shelf.

[Status.] Scribble. Destined: taxonomy to Paper 1 (meta shelf, next to V_{derived}); the pointer table to Assembly; lookup-and-reread to Inference. Not merged. Fine-tune later: who assigns ids, instance vs kind under $A \oplus A = A$, and whether one atom may keep a stack of parent-bowl ids after several collapses.

9 A theoretical problem, split the same way Section 2 already split one — new, v6

Can every atom in a unit space, across all labels and types, be merged to one representative atom? Posed precisely: a bounded region holds many already-settled atoms, of possibly *differing* sorts τ , not just differing content within one sort. Can the whole collection become one representative for that region? Framed, as requested, with both a theoretical and a practical answer in view, the way the sphere-packing question above was.

[Solved — but only for the restricted case, which is Section 2’s correction in general form.] When every atom in the region shares one sort (or, more precisely, one already-admitted φ_τ direction-family), this is exactly Assembly’s already-verified system-promotion mechanism: many close atoms, one representative, via the Hill bijection rather than the naive clip, because Hill preserves evidence-count where the clip saturates and loses it. Sixteen camps correctly became thirteen systems on this basis, not eleven. “A lot of random atoms” changes nothing here — randomness in position or content is fine, provided the sort is shared. This is Section 2’s τ_{weather} correction, restated as the general pattern it always was: random instances of one τ , closely aligned, merge cleanly.

[Still open — this is Gap 1, not a second result.] “Across all labels and types,” taken literally, names the harder case: atoms of genuinely *different* sorts in the same region. This is not solved by Hill’s equation or the bijection. Confirmed twice already in this project: both Inference’s Hill form and Assembly’s Hill bijection fix magnitude saturation only, and neither touches direction or sort. Merging across genuinely different sorts into one atom is precisely Gap 1 (Section 1), for

which a leading candidate (refuse outright, per the `chladni` precedent) exists but has not been written into Paper 1.

[Theoretical versus practical, for the cross-type half specifically.] The sphere-packing question above had a well-defined continuous quantity (point count) that a practical approximation could meaningfully estimate. This question’s open half does not: there is no stated φ_τ for combining across sorts to approximate in the first place. A random or approximate merge rule cannot stand in for a merge rule that has not been defined — this is an admissibility gap, not a precision gap, and the two should not be read as the same kind of “open” question.

[So, precisely, correcting the framing in which this was posed.] “We solved it using bijection and Hill’s equation” is true for same-sort regions, and is a real, verified, reusable result — but it is not yet true for atoms spanning genuinely different types, which remains exactly Gap 1, unresolved by anything checked so far. The two cases should be kept as separate claims, the same way Section 2 already separated them once for τ_{weather} specifically; this entry generalizes that same correction rather than adding a new one.

10 Destined for Pattern Arithmetic (scribble, v7 — a scoping refinement to Route 1 of Gap 1)

“Refuse outright” is a claim about $\boxplus$, not a permanent bar on ever combining unlike sorts by any mechanism. Raised as a caution on Section 1’s leading candidate: likely that only like-typed atoms will ever practically need merging, but unwise to state Route 1 in a way that forecloses future, explicitly-justified mechanisms for binding unlike sorts, should a practical need for one arise.

[The distinction that matters.] Route 1 was always a claim about $\boxplus$ specifically — that this particular operator, as defined, was never asked the cross-direction question and correctly declines to answer it. It was never a claim that no mechanism, under any name, could ever combine unlike sorts. Conflating the two would overclaim something the evidence never supported.

[Why this is not a new principle, but a confirmation of one already load-bearing in the paper.] Axiom 0 is already built this way: “a sort absent from this table is undecided, not admitted; treat it as inadmissible until a φ_τ is stated.” Inadmissible *now*, for lack of a stated proof — not inadmissible forever, by declaration. `chladni` is the existing demonstration: admitted later, on stated grounds, not pre-emptively barred. Even `chladni`’s own hedge on cross-mode combination, the strongest evidence behind Route 1, uses “may be closed negatively... rather than left waiting” — a leaning, not a verdict.

[How Route 1 should actually be worded, if merged.] Not: “ $\boxplus$ refuses states at differing directions, full stop.” Better: “ $\boxplus$, as currently defined, refuses states at differing directions; a cross-sort reading remains possible in principle, as its own stated mechanism, admitted on its own terms the way any new capability is — not as a modification of $\boxplus$ itself.” This keeps the refusal honest about what has actually been shown without overclaiming permanence the evidence does not support.

[Status.] Scribble. A scoping note on Section 1’s existing entry, not a new result and not yet written into that section’s own text. “Probably rare” (practical likelihood) and “permanently barred” (structural impossibility) are different claims; only the first has support, and Route 1’s eventual wording should reflect only that one.

11 Destined for Pattern Arithmetic (scribble, v8 — proposed terminology realignment, with cost assessment, not merged)

Proposed: Collapse, UnCollapse, Bundle, MixMerge, UnMix. A full realignment of five terms, proposed together:

- **Collapse** ($\oplus$) — one atom, one origin, several internal vectors.
- **UnCollapse** ($\ominus$) — subtract one atom’s vectors from another; not previously discussed in depth.
- **Bundle** — a bowl/set, many atoms, kept separate, separate origins.
- **MixMerge** ($\boxplus$) — one new atom, fused.
- **UnMix** — not possible.

This reassigns Collapse and Bundle relative to the reading settled two exchanges earlier in this project (Collapse = $\oplus$ = the bowl, confirmed three times directly from source text and from the bowl’s own definition as a separate finite set). The proposal here reverses that assignment. Recorded as a deliberate choice to evaluate, not adopted, and not written into Paper 1.

[Blocking: a direct symbol collision.] $\ominus$ is not available for UnCollapse. It already carries its own formal axiom: “Axiom 4: Gather and remove ($\oplus$, $\ominus$)... $\ominus$ undoes a gather. It is not anti-phase interference.” Two incompatible operators cannot share one symbol in the same document. A different symbol is needed before UnCollapse can be merged under any sequencing; this is not a matter of care in editing, the symbol itself is already spoken for.

[Instance counts, checked directly against the source.]

Term	Current meaning	Count	Title usage
Collapse	$\oplus$ ’s bowl-forming act	14	Yes — a subsection title
Mix / $\boxplus$	fused, single output	77 / 50	Yes — Axiom 5’s own title
Bundle	one atom, N internal vectors	10	No
Bowl	a set of separate atoms	80	No, used throughout

[Cost, per piece.]

- *Collapse as one atom*: high cost, and a reversal rather than a rename. The “Process principle: Collapse, path, result” subsection’s own stated reasoning — “collapse is the act that discards the order and count of that sequence... R = the bowl” — would need rewriting, not relabeling; its current logic explains why collapse yields a set, and that explanation would need to become its own opposite.
- *Bundle as bowl*: high cost. Collides with Bundle’s own existing, different meaning (Axiom 7’s N -vectors-in-one-atom). Bowl already carries 80 instances of its own, separate meaning; this is a reconciliation of two already-distinct, already-used terms, not an added synonym.
- *MixMerge, rename only, same underlying behavior*: moderate cost, mechanical. 77 instances is substantial, but nothing about $\boxplus$ ’s actual behavior changes — risk is missed instances or inconsistent phrasing, not logical breakage.
- *UnMix, named but still impossible*: low cost. The concept is already stated (“division as unmixing is shown ill-posed”) — this mostly formalizes an existing result with a name and a symbol, once a free symbol is chosen.

[**Overall, honest verdict.**] The $\ominus$ collision is a hard blocker, independent of sequencing, and needs resolving before anything else proceeds. Past that, Collapse-reversal and Bundle-reassignment are the expensive, high-risk pair, each touching foundational, load-bearing text and interacting with one another. MixMerge and UnMix are comparatively cheap and could move first, independently, as a lower-risk starting point if this is pursued.

[**Status.**] Scribble. Cost assessed, not adopted. No change made to v15_19. Awaiting a decision on sequencing, and a replacement symbol for UnCollapse, before any part of this could be merged.

12 Destined for Pattern Arithmetic (scribble, v9 — symbol resolution for Bundle and UnCollapse, plus when ids are actually needed)

Bundle gets a symbol and becomes a verb.

$$\text{Bundle : } A_1 \uplus A_2 \uplus \cdots \uplus A_n \longrightarrow B \quad (\text{many atoms, kept separate}).$$

Checked clean against the source: $\uplus$ (multiset union) does not appear anywhere in Paper 1. The choice is principled, not arbitrary — $\uplus$ ’s standard mathematical meaning, combine while preserving separate membership, is exactly Bundle’s job. Worth being precise about what this move actually is: with $\oplus$ reassigned to Collapse under the v8 scheme, Bundle-as-verb takes over the role Paper 1’s own $\oplus$ currently performs (the bowl-forming act, nothing merged, nothing lost). Not a new mechanism — a new name and symbol for an existing one, freed up because the old symbol moved.

UnCollapse gets a symbol. $\ominus$, also checked clean against the source. Visually related to $\ominus$, so the family resemblance between Collapse/UnCollapse reads correctly, while remaining a genuinely distinct symbol — no collision with Axiom 4’s own $\ominus$.

[**Precisely when ids are needed for UnCollapse, and when they are not.**] Collapse, under this scheme, is lossless by construction: every v_i stays individually intact inside the resulting atom. Splitting it back apart is *already* structurally possible without ids whenever every incoming vector has a distinct $(r_i, \hat{n}_i, \chi_i, \tau_i)$ — the bundle’s own listing already carries everything needed.

Ids earn their place in exactly one case: two *distinct* source atoms that happen to share identical values. Without an id, the two resulting entries inside the bundle are indistinguishable from each other — UnCollapse would return the right values, with no way to confirm whether this was genuinely two separate instances that happened to match, or something else that produced the same numbers. This is the same instance-versus-kind distinction already on file from the v5 scribble: “two distinct instances of the same pin... need two ids.”

So, stated precisely rather than as a blanket claim: UnCollapse is value-exact with or without ids. What ids add is identity-exact recovery — confirmation that what comes back out is traceably the same atom that went in, not merely one with matching numbers. For the ordinary, all-distinct case, this guarantee is free. For the degenerate, duplicate-value case, it does not exist without an id. That is the honest scope of “may become a possibility with ids” — a fix for one specific, real ambiguity, not a general precondition for UnCollapse to work at all.

[**Status.**] Scribble. Two symbol collisions from v8 are now resolved (Bundle: $\uplus$; UnCollapse: $\ominus$). The Collapse-reversal and Bundle-reassignment costs noted in v8 are unchanged by this — resolving the symbols does not reduce the cost of rewriting the foundational passages that currently use these terms the other way. Not written into v15_19.

13 Considered and found to need nothing filed

For completeness: the reference-atom and precursor-witness mechanism, the bundle-vector / uncapped V_{derived} distinction from Mix, the $\pi = (x; w_1, \dots, w_n)$ poem trajectory, and the precise definition of χ as internal phase-color (already stated in Axiom 1: “two identities may occupy the same direction without overwriting one another”) were all discussed at length and all turned out to already be fully present in Inference and Paper 1 respectively. Nothing from those threads is parked here, since there was nothing left unresolved to park.