

Pattern Arithmetic — Quick Reference, Index and Glossary

Extracted from *Pattern_Arithmetic_v15_28*. Section numbers (§) and equation numbers (eq.) refer to that document.
Every number in Part 9 was recomputed from the paper’s printed inputs (*reproduce_v15_28.py*).

0 · The soul of the paper, on one page

The soul in eight paragraphs

Thesis (§1). Mathematics need not be performed on numbers alone. Wherever a domain supplies a *unit of truth* — the smallest thing it treats as one — and a small set of rules by which such units combine, obeyed consistently, the unit and its rules form an *arithmetic of that domain*. Counting ($1 + 1 = 2$, “how many?”) is one instance, not the only one: the word “add” names a different rule, and therefore a different kind of unit, for a colour, a note, a word.

The unit (Axiom 1, §2.2). One typed point, not a field: $x = (r, \hat{n}, \chi, \tau)$ with strength $r \in [0, 1]$, direction $\hat{n} \in S^2$ (a *semantic* address, not a place), phase $\chi \in [0, 2\pi)$, sort τ (which buttons may be pressed). The sphere is the stage; the unit is the vector.

The spine (§3.1). A *bowl* of seated units ($\uplus$) is read by a *path* (T) to a *result*; a result may be seated again (*levels are closed*). A unit does not know what it constitutes: **a seated word does not know the poem; the path knows it**. One bowl, several ways to read it.

Six verbs, six questions. $\uplus$ *what is present?* · $\oplus$ *what is combined, nothing discarded?* · $\boxplus$ *what composite results?* · T *what happens, in what order?* · offset *what walk is reused?* · $\bowtie$ *how do two units relate while both still sit?* Three have a counterpart that takes something back ($\setminus$, $\ominus$); Mix’s own, $\boxminus$, is proven impossible.

Six laws worth remembering. $A \uplus A = A$ · $A \oplus A \neq A$ for every A · Mix is lossy and, under the cap, non-associative · $T_a T_b \neq T_b T_a$ yet $T_w * T_w = \text{id}$ · Assembly never repels ($U_{AB} < 0$) · UnMix does not exist.

What it refuses, on purpose. Mix across differing directions (declined, not left open) · UnMix · a geometric reading for any sort with no structure-preserving map (Axiom 0; most physical domains) · combining across sorts (no stated rule) · treating a costume (storm, forest, table and chair) as a model.

What it keeps honest. Every law is deterministic: the same printed starting units — read at the same time, where any carries an expiry — give the same result. Every choice (the clip, $\alpha = \pi r$, η , δ , the census cut, ϵ_{allow}) is a *named convention*, not a truth of the algebra.

How an atom is counted (§2.10). An atom is N *cores* holding Q informational *vectors*; $D \leq N$ of the cores are distinct. Teaching atom: $N = Q = 1$. Empty atom (black): $N = 1$, $Q = 0$. Beside its content an atom may carry *meta vectors* (time, expiry, place, links, V_{derived}) that describe it and are no part of what it says (§2.11).

1 · The verbs

Verb	Question	Defining rule	Counterpart	Where
Bundle $\uplus$	What is present?	Union on bowls: $B_1 \uplus B_2 = B_1 \cup B_2$. Idempotent, commutative, associative; sort-blind ; discards order and count. $A \uplus A = A$; $A \uplus B = \{A, B\}$.	Remove $\setminus$: $\{A\} \setminus A = \emptyset$ (empty bowl $\neq$ black 0).	§2.5, eq. 5–7
Collapse $\oplus$	What is combined, with nothing discarded?	Keeps every core of every operand, merges none: $N = \sum N_i$, $Q = \sum Q_i$. Not idempotent: $A \oplus A = A^{(2)}$. One sort per atom today; across sorts open. Each core keeps its meta vectors.	UnCollapse $\ominus$: returns the atom without one core; $N \rightarrow N - 1$ ($N \geq 2$); at $N = 1$ leaves black; Q falls by that core’s vector count.	§3.3, §3.4

Verb	Question	Defining rule	Counterpart	Where
Mix $\boxplus$	What composite state results?	Phasor sum at a shared $\hat{n}$: $z' = \sum r_i e^{i\chi_i}$, $r' = \min(1, z')$, $\chi' = \arg z'$; $\tau' = \tau$; black if $z' = 0$. Over a bowl, one step: $\boxplus(B)$, $z_B = \sum_{x \in B} r_x e^{i\chi_x}$. Lossy; commutative; binary form non-associative under the cap. Refused across differing $\hat{n}$.	UnMix $\boxminus$: impossible — a Mixed result does not determine its sources.	§2.6, eq. 8–9; §3.2; §8.2
Transform T	What happens, and in what order?	$T_w(x) = (r_x, \text{Rot}_{\hat{n}_w}(\alpha(r_w)) \hat{n}_x, (\chi_x + \chi_w) \bmod 2\pi, \tau_x)$, $\alpha(r_w) = \pi r_w$. Leaves r_x, τ_x alone. $T_0 = \text{id}$; composition associative; $T_a T_b \neq T_b T_a$.	Inverse actor $w^* = (r_w, -\hat{n}_w, -\chi_w \bmod 2\pi, \tau_w)$, so $T_w^* T_w = \text{id}$.	§4.2–4.3, eq. 20, 22
Offset	What walk is reused?	$v = \log_{\hat{n}_m}(\hat{n}_k)$; $\hat{n}_* = \exp_{\hat{n}_w}(\text{transport}_{m \rightarrow w}(v))$; $r_* = r_w$, $\chi_* = (\chi_w + \chi_k - \chi_m) \bmod 2\pi$, $\tau_* = \tau_w$. Copies a walk; does not mint the name. Not T .	—	§5.1, eq. 29–35
Assembly $\boxtimes$	How do two units relate while both still sit?	$A \boxtimes B = (A, B, \kappa_{AB}, U_{AB})$. Both operands survive unchanged; nothing is fused. $U_{AB} < 0$ always: the bracket $3 + \hat{n}_A \cdot \hat{n}_B$ stays in $[2, 4]$.	—	§2.9, eq. 12–14

Schoolbook + (“how many?”) lives only in **num** mode (decode, add in $\mathbb{Z}$, encode; §6.2). The paper reserves each verb name “the way a keyword is reserved in a programming language”: once given a meaning it is not reused elsewhere for an unrelated or informal sense (Table 1).

2 · The axioms

#	Name	One-line content	Where
0	Admissibility of Sorts	A sort τ may claim a geometric χ - or T -reading only through a map $\varphi_\tau : D_\tau \rightarrow \mathcal{S}$ with $\varphi_\tau(a \circ_\tau b) \approx \boxplus(\varphi_\tau(a), \varphi_\tau(b))$ or $T_{\varphi_\tau(a)}(\varphi_\tau(b))$ at a stated fidelity (Tier 1). Tier 0 (word , num) is inert. An inadmissible sort (e.g. snowflake, D_{6h} growth) may still be seated, at Tier 0 only. A sort absent from the table is undecided, not admitted.	§2.1, eq. 1
1	Normalized Unit Truth State	A pattern is one typed point, $x = (r, \hat{n}, \chi, \tau)$, $r \leq 1$ a normalized activation (not a probability). A bowl is a separate finite set of such atoms. The bound is imposed by clipping — a choice, with its trade stated (not injective; $r = 1$ is a value, not a limit).	§2.2, eq. 2
2	Absolute Null State (Black)	$r = 0 \Rightarrow x = \mathbf{0}$: direction and phase unused — an <i>empty seat</i> . As an atom: $N = 1$, $Q = 0$. A vector of strength zero is not an arrow.	§2.3, eq. 3
3	Saturation (White)	The upper pole of the same bounded stage: one full-spectrum name, or a saturated bowl. No unit is $r = \infty$. “Density” = many identities share the stage without overwrite.	§2.4, eq. 4
4	Bundle and Remove	$\uplus$ collects (union, idempotent); $\setminus$ removes one member. Sort-blind: sort discipline governs what may later be <i>done</i> with a bowl, not who may sit in it.	§2.5, eq. 5–7
5	Mix as a reading	A phasor operator: combines χ among states that already share $\hat{n}$; never moves an arrow. A reading returns a state and leaves the bowl intact.	§2.6, eq. 8–9
—	Mix across differing directions: refused; chladni inclusion	Cross-mode coexistence is $\uplus$, not $\boxplus$: mode space is not closed under addition. Refusal is the closure — settled by declining, not by finding a rule. chladni is Tier 1: exact within one mode, undefined across modes.	§2.7, eq. 10
—	Rotation preserves the unit	$\mathbf{R} \in \text{SO}(3)$ turns $\hat{n}$ and leaves r, χ, τ : the geometric engine of T , not a second blend.	§2.8
6	Assembly	Relates two intact units by a contact point and an interaction energy; never repels; completed by the enclosure’s seats $\vec{P}$.	§2.9, eq. 12–14
7	Informational vectors and derived per-atom metrics	An atom may hold N cores of Q informational vectors under one closure, each an instance of Axiom 1, all one sort. Defines V_{derived} , seven scalar metrics, Q_{eff} , the empty atom.	§2.10
—	Meta vectors (not an axiom)	What an atom carries <i>beside</i> its content: uncapped, not informational, in neither N nor Q ; rides with its core.	§2.11

3 · The unit, its coordinates, and the counts

Symbol	Domain	What it answers
r	$[0, 1]$	How strong / how confidently held is this unit?
$\hat{n}$ (or (θ, ϕ))	S^2 (higher: $S^{N_{\text{dim}}-1}$)	What direction — the semantic address?
χ	$[0, 2\pi)$	What internal phase / mood / hue?
τ	{word, num, color, note-pitch, chladni, ...}	What sort — which operators may legally act on it? (<i>au</i> does not move the arrow.)

Term	Meaning
teaching atom	The 4-tuple of Axiom 1: one arrow, $N = Q = 1$. One state among several an atom can be in.
atom (Collapsed)	N cores holding Q informational vectors $v_i = (r_i, \hat{n}_i, \chi_i)$, $0 < r_i \leq 1$, under one closure; all vectors share one sort τ . Arises by Collapse or is seated directly by whatever supplies the seat.
core	A group of vectors held together in an atom. Holds any number of vectors: one, many, or none. “Core” is reserved for this sense.
N	Number of cores . A count of the stored atom: does not change with the reading time t .
Q	Number of informational vectors — “the atom’s info number”. $Q \geq N$ is usual, not a rule. Read over the cores valid at t .
D	Number of distinct cores among the N : two cores coincide exactly when $\oplus$ would seat them together (content only; empty cores coincide; meta vectors never enter). $D \leq N$, equality iff every core is distinct. Stored count.
$N - D$	How many times some core repeated.
$A^{(k)}$	k identical copies of A ; has kN_A cores. $A \oplus A = A^{(2)}$.
empty atom	Black: $N = 1$, $Q = 0$. $S, \mathcal{V}, \mathcal{E}, V_{\text{derived}} = 0$; $\gamma_{\text{atom}} = 0$, $\sigma_{\text{atom}} = 0$ by convention; $\alpha_{\text{atom}}, \sigma_{\text{shell}}, Q_{\text{eff}}$ undefined. Collapse keeps a silent contribution: $A \oplus 0$ has one more core, same vectors; $0 \oplus 0 \neq 0$.
mass number / atomic number	N reads like a nucleus’s mass number (every core counted), D like its atomic number (distinct cores). An analogy of shape only.

4 · Metrics (Axiom 7, §2.10)

Named in the paper: *atom strength*, *atom volume*, *mean vector volume*, *atom coherence*, *effective vector count*; $\mathcal{E}_{\text{atom}}$ is “the atom’s intrinsic energy”; $\alpha_{\text{atom}}, \sigma_{\text{atom}}, \sigma_{\text{shell}}$ are its fifth, sixth and seventh metrics, described but not named (descriptive labels below are marked). All presuppose an atom of Q informational vectors $v_i = r_i \hat{n}_i$. None is used by any verb; each is read over the cores valid at the reading time. Identity: $|V_{\text{derived}}|^2 = \mathcal{E}_{\text{atom}} + 2 \sum_{i < j} r_i r_j (\hat{n}_i \cdot \hat{n}_j)$, the structure of a variance of a sum.

Symbol	Formula	What it measures	Requires
V_{derived}	$\sum_{i=1}^Q r_i \hat{n}_i$ (χ excluded)	The raw, uncapped vector sum; per core and per atom (the atom’s is the sum of its cores’). Reads as an effective pair $r = V $, $\hat{n} = V/r$ for a weight or a direction; not itself a unit; needs a cast into $[0, 1]$ before acting as T ’s actor.	any Q
S_{atom}	$\sum_i r_i$	Atom strength . Denominator of atom coherence.	any Q
$\mathcal{V}_{\text{atom}}$	$\sum_i r_i^3$	Atom volume . Strictly $< S_{\text{atom}}$ whenever any $r_i < 1$ — forced by the cap.	any Q
γ_{atom}	$\mathcal{V}_{\text{atom}}/Q$	Mean vector volume .	$Q \geq 1$ ($:= 0$ at $Q = 0$)
$\mathcal{E}_{\text{atom}}$	$\sum_i r_i^2$ (eq. 15)	Intrinsic energy (the paper’s phrase). Stands on its own; the energy the vectors would carry if every pair were orthogonal.	any Q

Symbol	Formula	What it measures	Requires
α_{atom}	$\frac{\sum_{i<j} r_i r_j (\hat{n}_i \cdot \hat{n}_j)}{\sum_{i<j} r_i r_j}$ (eq. 16)	<i>Fifth metric; unnamed.</i> Strength-weighted mean cosine similarity across all internal pairs; separates active cancellation (< 0) from mere non-reinforcement, which atom coherence cannot.	$Q \geq 2$
σ_{atom}	$\frac{Q^2}{Q-1} \text{Var}(s_1, \dots, s_Q),$ $s_i = r_i^3 / \mathcal{V}_{\text{atom}}$ (eq. 17)	<i>Sixth metric; unnamed</i> (concentration of volume). 0 when even, 1 when one vector holds all. Equals the normalized Herfindahl index of the cubed shares.	$Q \geq 2$ ($:= 0$ at $Q \leq 1$)
$\sigma_{\text{shell}}(w)$	1 if $K = 1$; else $\frac{K^2}{K-1} \text{Var}(p_1, \dots, p_K),$ $p_k = n_k / Q$, bins of width w (default 0.1) (eq. 18)	<i>Seventh metric; unnamed</i> (concentration across shells). Catches what σ_{atom} cannot: many vectors independently landing at the same magnitude.	$Q \geq 1$
Q_{eff}	$S_{\text{atom}}^2 / \mathcal{E}_{\text{atom}}$	Effective vector count — not an eighth metric. $1 \leq Q_{\text{eff}} \leq Q$; scale-free; the largest vector holds at least $1/Q_{\text{eff}}$ of the strength; a <i>count</i> , with Q_{eff}/Q the size-free form.	$S_{\text{atom}} > 0$
atom coherence	$ V_{\text{derived}} / S_{\text{atom}}$	How much strength survives into one resultant; 1 for aligned vectors, falling toward 0. Also a cast of V_{derived} into $[0, 1]$.	$S_{\text{atom}} > 0$

5 · Meta vectors (§2.11)

What they are. Values that describe an atom and are no part of what it says. Not capped; not informational; counted in neither N nor Q ; outside the seven metrics and D ; no admissibility of their own; **not a coordinate** (the four coordinates and the sort tag are unchanged). They ride with the cores they describe. Anything beyond that — what a meta vector *means* — is extracted by context or by the path that reads it.

Kind	Held	What it is
V_{derived}	per core and per atom	Sum of the informational vectors, recomputed whenever the content is. Already in the paper (Axiom 7).
seat $\vec{P}$	per unit or bowl	External position in unbounded Euclidean space; enters only $d_{AB} = \vec{P}_B - \vec{P}_A $ (§10.1). Already in the paper.
$V_{\text{timestamp}}$	per core	When the core was born, an absolute (non-cyclic) time, so $\text{age} = t - V_{\text{timestamp}}$ is derived, never stored. An optional clock-face vector (minute on ϕ , hour on χ , θ at the equator) groups by time of day but cannot replace it. A Collapse result also gets its own new timestamp.
V_{expiry}	per core	When the core stops being valid. Absence means no expiry. Valid at t iff t is earlier than every expiry it carries.
V_{location}	per core	A point on a sphere (latitude as polar angle, longitude as azimuth). Not a sort: a place is a coordinate, not a claim with a strength.
V_{link}	per core	Names another atom by one of that atom's meta vectors, never by content (a word is never linked to a poem). Only backward links are stored, at birth; forward and bidirectional links are views (a search for atoms whose backward link names a given one). A Collapse result gets a backward link naming its sources.

Rules that govern all of them

Rule	Statement
Form	Always vectors, with no constraint on form. A scalar meta vector takes a designated axis of its core and carries its value as the coordinate there (uncapped), so vector distance is the difference of values; more coordinates (a plane or more) where a kind needs them. Which axis suits which kind is a recommendation (a catalogue, §6.7), not an axiom.
Turning	Turn an atom as a whole (same rotation and phase shift on each vector): V_{derived} turns with it; scalar meta vectors are carried unchanged; angles, magnitudes and distances among the information vectors and V_{derived} are preserved.

Rule	Statement
Atoms are not edited	An atom is a value. Change is Collapse or UnCollapse, each returning a new atom; recorded meta vectors never change after birth. There is no “last modified”.
Equality ignores them	Equality of cores (for $\uplus$, D , UnCollapse) is on content only. Were the timestamp part of the value, $N - D$ would always be 0 and $A \uplus A = A$ would never apply.
Expiry is a reading	Everything is read at a time t (default now). An expired core contributes no vectors and reads as a black core; N , D are stored counts and do not change with t ; Q , V_{derived} , the metrics and every process read only valid cores. An atom’s expiry applies to all its cores; a contribution’s expiry is carried onto each of its cores at Collapse. A backward link is lineage, not participation, and stays readable.
Nothing loses identity	Under Collapse, Bundle and Mix the individual persists as meta vectors; this is persistence, not recovery. Mix leaves its sources unconsumed and names them by backward links; UnMix stays impossible.
Absence is not zero	For any meta vector: an absent location is not $(0, 0)$; “expiry 0 means none” would read as already expired.
No verb changes one	No verb changes a meta vector or takes one as an operand; a verb may read one (Assembly reads V_{derived} and the seat). A meta vector never becomes an actor.
Uses	Group atoms for a task (collect, order, assemble). Grouping needs a comparison stated with each kind (great-circle for a location, cyclic for a clock face, equality for an identifier); none is common to all. Direction is the cleaner assembly rule; magnitude enters as a weight. Links declare structure and Assembly’s energy produces it from geometry and a seat; the two are separate. Links can carry an order past $\uplus$, which does not read them, so order still comes from outside and the path is where it is walked. At contact the best partner of an atom of strength 0.60 has strength 0.30; at separation 3 it is the strongest available.

6 · Other named quantities, by operator

Symbol	Formula	Lives in	What it is
U_{AB}	$-\frac{r_{AB}}{2} \cdot \frac{3 + \hat{n}_A \cdot \hat{n}_B}{(d_{AB}^2 + \varepsilon_{AB}^2)^{3/2}}$ (eq. 13)	§2.9	Assembly’s interaction energy. Product r_{AB} : how strong; bracket in $[2, 4]$: how aligned (at most a factor of two); denominator: how far. Always < 0 for nonzero strengths; exactly 0 for a black unit.
ε_{AB}	$r_A + r_B$	§2.9	Softening length, derived by reading r as a radius.
κ_{AB}	$\hat{n}_A + \hat{n}_B$ (unnormalized; eq. 14)	§2.9	Contact point; defined at every angle, = $\mathbf{0}$ at antipodal. Its magnitude falls from 2 to 0.
polarity	$ r_A - r_B $	§2.9	Labels a bond as balanced or lopsided; does not change what the relation does.
$d_{AB}, \hat{r}_{AB}$	$ \vec{P}_B - \vec{P}_A , (\vec{P}_B - \vec{P}_A)/d_{AB}$ (eq. 46)	§10.1	Separation and separation axis; no axis derived from orientation is used.
$\alpha(r_w)$	πr_w	§4.2	Strength to rotation angle; $r_w = 1$ gives 180° .
z_B	$\sum_{x \in B} r_x e^{ix}$; $\boxplus(B) = (\min(1, z_B), \hat{n}, \arg z_B, \tau)$ (eq. 9)	§2.6	Mix’s one-step bowl reading; the reason binary Mix is non-associative under the cap.
$\log_{\hat{p}}, \exp_{\hat{p}}$	$\frac{\theta}{\sin \theta}(\hat{q} - (\hat{p} \cdot \hat{q})\hat{p}); \cos\ v\ \hat{p} + \sin\ v\ v/\ v\ $ (eq. 29–30)	§5.1	Walk from p to q as a tangent vector, and back.
transport	$\text{Rot}_{\hat{u}}(\theta_{mw}) v, \hat{u} = \hat{n}_m \times \hat{n}_w / \ \cdot\ $ (eq. 32–33); general N_{dim} : a rotation within the plane the two seats span (eq. 35)	§5.1	Carries the walk from seat m to seat w . The two seats name the turning plane, so no Collapse is needed. Exactly the cross-product form at $N_{\text{dim}} = 3$.
F_i	$-\nabla_{\vec{P}_i} \sum_{j \neq i} U_{ij}$; step $\vec{P}_i \leftarrow \vec{P}_i + \delta F_i$ (eq. 47)	§10.2	Force on a seated unit, descended with no mass and no momentum.
freeze at resolution η	$\max_i F_i < \eta$ (eq. 48)	§10.2	A stated checkpoint, not exact rest: the pull is positive at every finite separation, so left longer every configuration drifts toward one pile.

Symbol	Formula	Lives in	What it is
camp, body $\vec{P}_{c,k}$	$\frac{\sum_{i \in G_k} r_i \vec{P}_i}{\sum_{i \in G_k} r_i}$ (eq. 49)	§10.3	A frozen group and its strength-weighted centre, read <i>after</i> motion. A system of camps is the <i>set</i> of bodies, never one grand average.
ϵ_{allow}	named, not derived	§10.3	Checkpoints whose bodies differ by less are read as the same freeze.

7 · Sorts, tiers, and who may do what

Sort	Native law $\circ_\tau$	Mix/ T map?	Tier
num	schoolbook +	no (chart only)	0
word	none claimed	n/a	0
note-pitch	pitch-class sum	near-exact	1
color	additive light	phasor Mix (here; a toy $0^\circ/120^\circ/60^\circ$ chart — not CIE, not linear RGB)	1
chladni	modal superposition	exact within a mode; undefined across	1 (restricted)
snowflake	D_{6h} growth	no	Tier 1 no; Tier 0 yes

Pair	+	$\uplus$	$\oplus$	$\boxplus$	$T_w(x)$	$x - a + b$
(word,word)	no	yes	yes	yes [†]	yes	yes
(num,num)	yes	yes	yes	no (clock trap)	no	no
(word,num), (num,word)	no	yes	open	no	typed	no
(color,color)	no	yes	yes	yes	yes	open
(note-pitch,note-pitch)	no	yes	yes	yes	yes	open
(chladni,chladni)	no	yes	yes	yes (within a mode)	open	open
any other pair	no	yes	open	open	open	open

§6.4 (Table 3). Every Mix entry is also subject to the place restriction (shared $\hat{n}$). $\uplus$ is *yes* for every pair: a bowl records presence and makes no domain claim. $\oplus$: same sort yes, cross-sort open. [†]Mix on two words is geometry, not a χ -reading of language. **A document is a sequence, not a blend** (§6.3): words and numbers meet only by *juxtaposition*, *typed action* (T across sorts needs an explicit output sort) or *cast*; dawn $\uplus$ 37.4 has no meaning until one side is cast.

8 · Dimensions: how an actor grows beyond three

Space	Sphere	Fixed part (the “axis”)	What turns
$\mathbb{R}^2$	S^1	a point, dim 0	one plane
$\mathbb{R}^3$	S^2	a line, dim 1	one plane
$\mathbb{R}^4$	S^3	a plane, dim 2	one plane
$\mathbb{R}^5$	S^4	a 3-D space, dim 3	one plane
$\mathbb{R}^{N_{\text{dim}}}$	$S^{N_{\text{dim}}-1}$	dim $N_{\text{dim}} - 2$	one plane

A teaching atom ($Q = 1$) is a legal actor ****only at $N_{\text{dim}} = 3$ ****; beyond that the actor’s atom must supply $N_{\text{dim}} - 2$ mutually orthogonal arrows ($Q \geq N_{\text{dim}} - 2$ is necessary, not sufficient): $T_w(x) = (r_x, \hat{n}_x^F + \text{Rot}_{\alpha(r_w)} \hat{n}_x^P, \chi_x + \chi_w, \tau_x)$, eq. 23. Orientation of $F^\perp$ is a convention, stated once. Checked invertible at every angle tested and non-commutative at $N_{\text{dim}} = 4, 5, 8, 12, 20, 50$; exact at $N_{\text{dim}} = 3$. Offset’s transport generalizes without any Collapse. Mix, Bundle and α_{atom} are dimension-blind. The paper keeps $\mathbb{R}^3$ as its standing assumption (§4.4).

9 · Numbers worth remembering (all recomputed from the printed inputs)

Claim	Printed value	Where
$(1\angle 0^\circ) \boxplus (1\angle 120^\circ)$ (red, green)	$1\angle 60^\circ$, yellow	§2.6
$0.6\angle 0^\circ \boxplus 0.8\angle 120^\circ$	$r' \approx 0.721$, $\chi' \approx 73.9^\circ$	§2.6, §8.1
$A \boxplus A$	$r' = \min(1, 2r)$ — not idempotent	§2.6
$A = B = 1\angle 0^\circ$, $C = 1\angle 180^\circ$ (by hand)	$(A \boxplus B) \boxplus C = \text{black}$, $A \boxplus (B \boxplus C) = 1\angle 0^\circ$; one-step $z_B = 1$	§2.6
Clock, $M = 24$: 7 + 5 and Mix of the two marks	12 (decode, add, encode); Mix gives $\chi' = 90^\circ$, hour 6	§6.2
Sun, Wind, Sky ($r_{\text{Sun}} = 1$, $r_{\text{Wind}} = 0.7$)	Sun then Wind $(-0.866, -0.500, 0.011)$; Wind then Sun $(0.278, -0.500, 0.820)$; $\alpha_{\text{Wind}} = 126^\circ$	§4.6
Four-dimensional actor, $r_w = 0.5$	$T_w(\hat{n}_x) = (\frac{1}{2}, \frac{1}{2}, 0, \frac{1}{\sqrt{2}})$, recovered by w^*	§4.7
Order gap, $N_{\text{dim}} = 4, 5, 8, 12, 20, 50$	0.683, 0.688, 0.649, 0.584, 0.489, 0.331	§4.4
Storm: Deepen / Clear endpoints	$(0.7071, -0.6124, -0.3536)$, $(110.7^\circ, 319.1^\circ)$ / $(0.3536, -0.6124, -0.7071)$, $(135^\circ, 300^\circ)$; both $\chi = 135^\circ$	§9.1
Offset on an equator $(0^\circ, 30^\circ, 90^\circ)$	lands at $(-\frac{1}{2}, \frac{\sqrt{3}}{2}, 0)$, 120° — a protractor check, not the operator	§5.2
Unmixing: $0^\circ/120^\circ$ against $10^\circ/100^\circ$	same phasor sum (weights 0.643, 0.766); residual at rounding level	§8.2
Alignment, $r = 0.85$, pairs 5.7° apart	$Q = 2$: raw 0.72, $\alpha = 0.995$; $Q = 5$ on a cone of half-angle 5.7° : 7.14, 0.988	§2.10
Three aligned at $r = 0.5$	$ V = 1.5$, $\alpha(1.5) = 270^\circ$ — not a rotation T accepts uncast	§2.10
Dominant vector 0.95 against three at 0.2	volume share 0.973	§2.10
Best partner for an atom of $r = 0.60$ (Assembly energy)	at contact $r_B = 0.30$; at separation 3 the strongest available (1.0)	§2.11
σ_{shell} : 10@0.98, 3@0.3, 400@0.65	0.908 at $w = 0.1$ and $w = 0.05$	§2.10
Enclosure, six printed units, $\eta = 10^{-5}$	628, 1261, 2526, 6320 steps at $\delta = 0.02, 0.01, 0.005, 0.002$; two camps $\{1, 2, 3\}$, $\{4, 5, 6\}$; bodies $(0.33336, 0.33333, 0)$, $(40.33330, 0.33333, 0)$; runs agree to 2×10^{-7} ; r -weighted and plain bodies differ by 2.4×10^{-8}	§10.4
Enclosure, pair $r = 0.9$, $r = 0.1$, $\delta = 0.02$	separation never increases, $3 \rightarrow 1.80$; dividing each step by r^3 (a 729-fold weighting) makes it oscillate	§10.4

Where a printed number cannot be reproduced exactly: the σ_{atom} comparison (0.00032 clumped against 0.00214 uniform) and the “2.200 against 2.150” strength comparison, because their populations are not printed (a uniform spread over $(0, 1]$ gives 0.0032); the count of 804 of 3000 steps on which the pair’s separation increases, which is rounding-sensitive (about 830–855 in replays; the qualitative claim is robust); and shell binning at floating-point edges ($[0.3/0.1]$ is 2 in floating point), where the printed populations are unaffected.

10 · Named conventions — choices, not truths

Convention	What it fixes	Where
the clip $r' = \min(1, z)$	How the cap is imposed. An order-preserving bijection $ z \mapsto z /(1 + z)$ would bound without discarding but makes $r = 1$ a limit. The clip is kept; the trade is stated.	§2.2
$\alpha(r) = \pi r$	The simplest monotone map from strength to rotation angle.	§4.2
orientation of $F^\perp$	Which direction counts as $+\alpha$ beyond three dimensions; any consistent choice, stated once.	§4.4
toy colour chart	Phasor Mix on $0^\circ/120^\circ/60^\circ$; neither CIE nor linear RGB.	§2.1, §2.6
numeric chart	$\chi = 2\pi k/M$, $M = 24$, at a fixed direction with $r = 1$.	§6.1–6.2
shell width w	Default 0.1 (ten shells on $[0, 1]$); reported with σ_{shell} .	§2.10
η , δ , census cut, ϵ_{allow}	Freeze threshold; step size; how camps are counted; when two checkpoints are the same freeze. The enclosure’s four measurement conventions.	§10.2–10.3

Convention	What it fixes	Where
reading time t	Default now; fixing it keeps every law deterministic. “Now” is external context, as the lexicon is.	§1.4, §2.11
standard dictionaries and the meta-vector catalogue	Recommendations, recorded and left open; not axioms.	§6.7

11 · Stated open, or not claimed

Item	Status in the paper	Where
Collapse across differing sorts	Open. Axiom 7 requires every vector in one atom to share the atom’s single τ ; Collapse across sorts needs a per-vector τ_i extension that Axiom 7 does not currently grant.	§3.3, §12.1
Collapsed atom as the actor of T	An example is not claimed here; a cast is named at the call site.	§2.10
Identity handle id_i	Proposed, not adopted: tells same-valued sources apart in UnCollapse.	§3.4
Reducing a path by merging close actors	Sequential actors already compose into one rotation for free (closure). Replacing merely <i>close</i> actors is a different operation, coarse-graining, whose shape only is stated: it yields an additional derived actor <i>beside</i> its sources and removes none. Which merge rule, and what drift bound, belongs to Inference.	§12.1
Dimensionality	The sphere S^2 is where the idea is taught; embedding-scale dimensions are a later choice.	§6.8, §12.1
Offset on Tier-1 sorts; antipodal offset	Untested here; antipodal seats have no unique short geodesic and that convention is not used.	§5.1, §6.4
Combining across sorts	No rule stated at all (a colour with a charge; a chord with a valence bond).	§1, §2.1
Verdicts	Saying what a resting state <i>means</i> is a later reading, not a verb here.	§1

12 · Document map

§	Title	What you will find
1	Introduction	Thesis; examples of other arithmetics; Pāṇini; scope of the claim (1.4); determinism.
2	Axiomatic Foundations	Axioms 0–7, Mix refusal and chladni (2.7), rotation (2.8), Assembly (2.9), metrics (2.10), meta vectors (2.11).
3	Process Principles	Bundle–path–result and levels (3.1); UnMix (3.2); Collapse (3.3); UnCollapse (3.4).
4	Transformation Algebra	T : definition, properties, beyond three dimensions, poems as trajectories, worked examples.
5	Offset	Log, carry, exponential; the equator check; offset is not T .
6	Encodings, Sorts, Mixed Documents	num and word ; the clock; a document is a sequence; operator legality; lexicons; standard dictionaries.
7	A Sketch of Conventions	Case and language do not move the address; a bag is not a sentence; a numeral is not a word until cast.
8	Operational Algebra	Mix of two shells; why unmixing is ill-posed; spectral storage (later).
9	Higher-order systems	Costumes: storm, table and chair, forest, pathogen.
10	The Enclosure	Seats, rest and freeze, camps and bodies, a replayable configuration, what it does not supply.
11–12	Discussion; Conclusion	Related work (Bloch sphere, embeddings, transformers, RotatE, von Mises–Fisher, Riemann sphere, $SO(3)$, spherical harmonics); future work.

13 · A–Z index of terms with special meanings

Term	Meaning	Where
Actor w	The unit that acts in $T_w(x)$: supplies axis, strength ($\rightarrow$ angle) and phase. Beyond three dimensions it must supply $N_{\text{dim}} - 2$ orthonormal arrows.	§4.2, §4.4
Admissible (geometrically)	A sort with a stated map φ_τ at a stated fidelity (Tier 1). Absent from the table = undecided, not admitted.	§2.1, eq. 1
Age	$t - V_{\text{timestamp}}$; derived, never stored (a stored age is wrong once time passes).	§2.11
Allowance ϵ_{allow}	Checkpoints whose bodies differ by less are read as one freeze; a named convention.	§10.3
Arithmetic (of a domain)	A unit of truth plus a small set of rules for combining such units, obeyed consistently.	§1
Assembly $\bowtie$	Relates two intact units by (κ_{AB}, U_{AB}) without fusing them.	§2.9
Atom	The typed unit. <i>Teaching atom</i> : one arrow. <i>Collapsed atom</i> : N cores, Q vectors, one closure.	§2.2, §2.10
Atom coherence	$ V_{\text{derived}} /S_{\text{atom}}$.	§2.10
Axis ($\hat{n}_w$, fixed part F)	The line (3-D) or subspace of dimension $N_{\text{dim}} - 2$ that T leaves fixed.	§4.4
Black 0	$r = 0$: an empty seat; $N = 1$, $Q = 0$; $T_0 = \text{id}$; the identity of nothing else. Distinct from the empty bowl $\emptyset$.	§2.3, §2.5
Body $\vec{P}_{c,k}$	A frozen camp's strength-weighted centre, measured afterwards by an observer; a description, not a destination.	§10.3, eq. 49
Bowl	A finite set of seated units: who is present. No order, no reading, no representation of the reading to come. May carry meta vectors; $\uplus$ does not read them.	§2.2, §2.5, §3.1
Bundle $\uplus$	Union on bowls; idempotent; sort-blind.	§2.5
Camp	A separated group frozen at resolution η ; their number is whatever the law, the seats and η produce.	§10.3
Cast (three uses)	(a) lexicon cast E_L : token to state (eq. 42); (b) sort cast, e.g. $E_{\text{num} \rightarrow \text{word}}$ (§6.3); (c) the cast of the uncapped V_{derived} back into $[0, 1]$ so it may act as T 's actor — the clip, or atom coherence (§2.10).	§6.6, §6.3, §2.10
Census cut	The stated separation (or chain of short links) used to count camps; a convention.	§10.3
Chart	An exact, invertible encoding (e.g. num); not a Mix or T homomorphism.	§2.1, §6.2
Checkpoint	One freeze read at some η after some time; not a destination.	§10.3
Clip	$r' = \min(1, z)$; the chosen way to bound strength; not injective.	§2.2, §2.6
Clock trap	Mix on (num , num): geometrically defined, type-refused. 7 and 5 Mix to hour 6.	§6.2
Collapse $\oplus$	Count-preserving combination; keeps every core; not idempotent.	§3.3
Companion paper	<i>Pattern Arithmetic: Inference from Bowls and Paths</i> — reference atoms, classification, abstention; where unmixing and path reduction are left.	§1, §3.2, §12.1
Contact point κ_{AB}	$\hat{n}_A + \hat{n}_B$, unnormalized; 0 at antipodal.	§2.9, eq. 14
Core	A group of vectors held together in an atom; any number of vectors. Reserved word.	§2.10
Costume	An illustration of the $\uplus$ -versus- T split (storm, table and chair, forest, pathogen); not a model.	§1.4, §9
Counting trap	$A \uplus A \uplus A \uplus A \setminus A = \emptyset$: four gathers are one seat, one removal empties it.	§2.5, eq. 7
Density	Many identities can share the stage without overwrite. Not a metric and not infinite information.	§2.4
Determinism	Same printed starting units (read at the same time where any carries an expiry) give the same result, step for step. Printing makes a result replayable; it does not make it right.	§1.4
Dictionary pin / doorbell	A seated unit for a name (man, storm, calm, Night). The algebra does not mint it; a lexicon or learning rule seats it.	§9.1, §5.2, §6.6
Direction $\hat{n}$	A pattern's semantic address, not its physical position.	§2.2, §10.1
Effective vector count Q_{eff}	$S^2/\mathcal{E}$; how many equally strong vectors carry the same energy.	§2.10
Empty atom	Black: $N = 1$, $Q = 0$.	§2.3, §2.10
Empty bowl $\emptyset$	A bowl with no occupant; $\neq \mathbf{0}$. T_0 is defined, $T_\emptyset$ is not.	§2.5, eq. 6
Enclosure	Seats, rest and camps: where Assembly's separation comes from.	§10

Term	Meaning	Where
Energy (two senses)	The atom's intrinsic energy $\mathcal{E}_{\text{atom}} = \sum r_i^2$ (§2.10, eq. 15); the interaction energy U_{AB} (§2.9, eq. 13). Not interchangeable.	§2.10, §2.9
Equality (of cores)	On content only; meta vectors never enter.	§2.11
Expiry V_{expiry}	When a core stops being valid; absence = none; an expired core reads as black.	§2.11
F_i (force)	$-\nabla_{\vec{P}_i} \sum_{j \neq i} U_{ij}$.	§10.2, eq. 47
Freeze (at resolution η)	$\max_i F_i < \eta$. A freeze, not a stop.	§10.2, eq. 48
Identity handle id_i	Proposed, not adopted: tells same-valued sources apart.	§3.4
Individual vs value	Under Collapse, Bundle, Mix the individual (with its meta vectors) persists; value is content only.	§2.11
Informational vector	One of an atom's Q arrows $v_i = (r_i, \hat{n}_i, \chi_i)$, $0 < r_i \leq 1$.	§2.10
Levels are closed	A result may be seated again and Bundled, walked, read at the next level.	§3.1
E_L (lexicon)	A map from one language's tokens onto states; case and language do not move the address.	§6.6, §7, eq. 42
Link V_{link}	Backward provenance stored at birth; forward and bidirectional are views.	§2.11
Location V_{location}	A point on a sphere carried as a meta vector; not a sort.	§2.11
Meta vector	A value that describes an atom and is no part of what it says.	§2.11
Mix $\boxplus$	Phasor reading of states sharing a direction.	§2.6
Mix across differing directions	Refused. Settled by declining, not by finding a rule.	§2.7
Mode (chladni)	A vibration mode; Mix exact within one, undefined across.	§2.7
Non-associativity (Mix)	A consequence of the cap: the binary operator is not associative; the one-step bowl reading is.	§2.6
Non-commutativity (T)	Inherited from $SO(3)$; the only thing that lets two poems from one bowl diverge.	§4.3, eq. 22
Occupant (register occupant)	The single current unit on the sphere that T acts on.	§2.2, §3.1
Offset	Copies a walk; does not mint the landing's name.	§5
Order gap	$ T_{w_2} T_{w_1} \hat{n}_x - T_{w_1} T_{w_2} \hat{n}_x $.	§4.4
Path / walk / trajectory	An ordered sequence of actors on an occupant; a poem is a T -walk. Order enters here and not in the bowl.	§3.1, §4.5
Phase χ	Internal phase / mood / hue: adds under T , sums as a phasor under Mix; reserved for mood, not language.	§2.2, §7
Polarity	$ r_A - r_B $.	§2.9
Pratyāhāra; vipratīṣedha	Pāṇini's sound-class shorthand and his conflict-resolution rule; offered as kinship, not identity.	§1
Reading	An operation that returns a state and leaves its inputs intact (Mix, a T -path, Collapse); also, expiry is a reading.	§2.6, §3.3, §2.11
Remove $\setminus$	Takes one member out of a bowl.	§2.5
Rest	A freeze at a stated resolution, never exact stillness.	§10.2
Result	Whatever the path makes of the bowl.	§3.1
Rotation	$SO(3)$ turn of $\hat{n}$ only.	§2.8
Seat $\vec{P}$	External position vector; a meta vector; enters only d_{AB} .	§10.1, eq. 46
Seated unit	A unit loaded on the sphere, pinned by a lexicon or a learned map.	§1.4, §3.1
Shell	(a) a width- w bin of strength in σ_{shell} (§2.10); (b) one phased identity at a direction in Mix ("two shells, not an angle", §8.1).	§2.10, §8.1
Softening length ε_{AB}	$r_A + r_B$, derived by reading r as a radius.	§2.9
Sort τ	Says which operators may legally act on a unit; does not move the arrow.	§2.2, §6
Stage $\mathcal{S}$	The unit sphere (or its higher analogue): the sphere is the stage, the unit is the vector.	§2.2
Step δ	Numerical sampling convention for the descent.	§10.2
Strength r	Normalized activation in $[0, 1]$.	§2.2
Tape	An ordered sequence of tokens; a document.	§6.3
Teaching atom	One 4-tuple, $N = Q = 1$.	§2.2
Tier 0 / Tier 1	Inert seating; or explicit inclusion by a stated φ_τ .	§2.1
Timestamp $V_{\text{timestamp}}$	Absolute birth time of a core; a Collapse result has its own.	§2.11

Term	Meaning	Where
Transform T	Rotate the occupant about the actor's axis by πr_w and add the actor's phase.	§4
Transport	Carries a tangent vector from seat m to seat w in Offset.	§5.1, eq. 33, 35
Turn (an atom)	Same rotation and phase shift on each vector; V_{derived} turns, scalar meta vectors are carried.	§2.11
UnCollapse $\ominus$	Returns the atom without one core.	§3.4
UnMix $\boxminus$	Impossible: sources not determined by the result.	§3.2, §8.2
Valid (core)	At t iff t is earlier than every expiry it carries.	§2.11
Volume $\mathcal{V}_{\text{atom}}$	$\sum r_i^3$, "atom volume"; descriptive only, deliberately excluded from motion.	§2.10
White	Saturation of the bowl, or a full-spectrum name; never $r = \infty$.	§2.4
D	Distinct cores among the N .	§3.3
N	Cores.	§2.10
N_{dim}	Dimension of the ambient space; 3 is the standing assumption.	§4.4
Q	Informational vectors; "the atom's info number".	§2.10
U_{AB}	Assembly's interaction energy.	§2.9, eq. 13
V_{derived}	Uncapped sum of the informational vectors.	§2.10
z_B	Phasor sum of a bowl.	§2.6, eq. 9