***New Phytologist* Supporting Information**

Article title: Fluctuating warm temperatures decrease the effect of a key floral repressor on flowering time in *Arabidopsis thaliana*

Authors: Liana T. Burghardt, Daniel E. Runcie, Amity M. Wilczek, Martha D. Cooper, Judith L. Roe, Stephen M. Welch, and Johanna Schmitt

Article acceptance date: 11 November 2015

The following Supporting Information is available for this article:

Fig. S1: Scatter plot of the relationship between days from sowing to bolt and days from sowing to flower.

Figure S2: Summary of results for all non-vernalized Columbia background genotypes from Experiment 1.

Figure S3: Summary of results for all non-vernalized L*er* background genotypes from Experiment 1.

Figure S4: Summary of results for all vernalized genotypes from Experiment 1.

Figure S5: Summary of results for all non-vernalized genotypes from Experiment 3.

Figure S6: Summary of results for all vernalized genotypes from Experiment 3.

Figure S7: Number of rosette leaves at bolting for select genotypes.

Figure S8: Differences in blade morphology and hypocotyl length between environments.

Figure S9: Reanalysis of data from Lempe (2005) on flowering time of a diverse panel of ecotypes in multiple constant temperatures.

Figure S10: Comparison of days to bolt data from high irradiance and low irradiance experiments.

Table S1: Description of genotypes, sources, and in what experiment they were used.

Table S2: Likelihood ratio tests to determine if fluctuating temperatures influence bolting times of wildtype genotypes (Col and Ler).

Table S3: Likelihood ratio tests to determine which allelic changes altered plant responses to fluctuating treatments

Table S4: Likelihood ratio tests to determine whether the bolting time of a *VIN3* mutation in the Col FRI_{SF2} background differed from the Col FRI_{SF2} background.

Table S5: Likelihood ratio tests to determine if high average temperatures influence bolting time of Col and Columbia FRI_{SF2} genotypes.

Table S6: Likelihood ratio tests querying whether each genotype behaves significantly differently to warm fluctuating conditions as compared to Columbia wildtype

Table S7: Likelihood ratio tests to determine whether each mutation in the Col FRI_{SF2} background influenced bolting within each treatment.

Table S8: Likelihood ratio tests to determine if fluctuating temperatures influence blade ratio.

Table S9: Likelihood ratio tests for the effect of fluctuating temperatures on hypocotyl lengths for Col, L*er*, and L*er phyB-1* in warm treatments.

Table S10: Likelihood ratio tests to determine if fluctuations at warm temperatures alter above ground biomass at various times throughout development.

Table S11: Likelihood ratio tests to determine if fluctuations at warm temperatures alter above ground biomass at bolt.



Fig. S1: Scatter plot of the relationship between days from sowing to bolt and days from sowing to flower. Color-coding is as follows: 22ConLD/SD (gold/orange), 22VarLD/SD (red/dark red), 12ConLD/SD (light blue/dark blue), 12VarLD/SD (light green/dark green).



Figure S2: Summary of results for all non-vernalized Columbia background genotypes from Experiment 1. For all graphs, boxes indicate 25% and 75% quartiles and heavy black line is the median. Whiskers extend to the highest value that is within 1.5 x the Interquartile range. Data outside of this range are outliers and visualized as points. Colors reference the environmental

treatment: Blue and green refer to the cool average temp treatments while red and orange refer to the warm average temperatures. Darker versions of each color refer to the short day, as oppose to long day, treatment. Blue and orange refer to constant environmental conditions and red and green refer to the variable conditions. Col Ama and Col Lis refer to two different sources of the Columbia ecotype.



Figure S3: Summary of results for all non-vernalized genotypes in the Landsberg *erecta* background from Experiment 1. For all graphs, boxes indicate 25% and 75% quartiles and heavy black line is the median. Whiskers extend to the highest value that is within 1.5 x the Interquartile range. Data outside of this range are outliers and visualized as points. Colors

reference the environmental treatment: Blue and green refer to the cool average temp treatments while red and orange refer to the warm average temperatures. Darker versions of each color refer to the short day, as oppose to long day, treatment. Blue and orange refer to constant environmental conditions and red and green refer to the variable conditions. L*er* and L*er* Lis refer to two different sources of the L*er* genotype.



Figure S4: Summary of results for all vernalized genotypes from Experiment 1. For all graphs, boxes indicate 25% and 75% quartiles and heavy black line is the median. Whiskers extend to the highest value that is within 1.5 x the interquartile range. Data outside of this range are outliers and visualized as points. Colors reference the environmental treatment: Blue and green

refer to the cool average temp treatments while red and orange refer to the warm average temperatures. Darker versions of each color refer to the short day, as oppose to long day, treatment. Blue and orange refer to constant environmental conditions and red and green refer to the variable conditions.



Figure S5: Summary of results for all non-vernalized genotypes from Experiment 3. For all graphs, boxes indicate 25% and 75% quartiles and heavy black line is the median. Whiskers extend to the highest value that is within 1.5 x the Interquartile range. Data outside of this range are outliers and visualized as points. Colors reference the environmental treatment: Blue

refers to the cool average temp treatments while red and orange refer to the warm average temperature. Darker versions of each color refer to the short day, as oppose to long day, treatment. Blue and orange refer to constant environmental conditions and red refers to the variable conditions (the 12Var treatments were not assessed in this experiment). Col FRI Ama and Col FRI SM refer to two sources of Col FRI_{SF2} NILs.



Figure S6: Summary of results for all vernalized genotypes from Experiment 3. For all graphs, boxes indicate 25% and 75% quartiles and heavy black line is the median. Whiskers extend to the highest value that is within 1.5 x the Interquartile range. Data outside of this range are outliers and visualized as points. Colors reference the environmental treatment: Blue refers to

the cool average temp treatments while red and orange refer to the warm average temperature. Darker versions of each color refer to the short day, as oppose to long day, treatment. Blue and orange refer to constant environmental conditions and red refers to the variable conditions (the 12Var treatments were not assessed in this experiment). Col FRI Ama and Col FRI SM refer to two sources of Col FRI_{SF2} NILs.



Figure S7: Number of rosette leaves at bolting for select genotypes in warm constant (orange) and variable (red) conditions in long (left panel) and short (right panel) day lengths. Black line is the median and colored boxes denote lower and upper quartiles (25% and 75%). Whiskers indicate the most extreme data point within 1.5 times the interquartile range and circles are points that fall beyond that range.



Figure S8: Differences in blade morphology and hypocotyl length between environments. (a) Blade length/total leaf length ratio of plants measured at bolting in Exp. 1 across all genotypes in L*er* (left) and Col (right) genetic backgrounds. Values above .5 indicate plants where the majority of the total leaf length is attributed to the blade. Values below .5 indicate leaves increasingly dominated by petiole length. Black line is the median and colored boxes denote lower and upper quartiles (25% and 75%). Whiskers indicate the most extreme data point

within 1.5 times the interquartile range and circles are points that fall beyond that range. Box width is proportional to the square root of the sample size. (b) Boxplots of hypocotyl lengths measured in the 12Con, 22Con, and 22Var treatments 14 days after seeds were sown. 12 individuals were measured in each environmental treatment. Black line is the median and colored boxes denote lower and upper quartiles (25% and 75%). Whiskers indicate the most extreme data points.



Figure S9: Reanalysis of data from Lempe *et al.* on flowering time of a diverse panel of ecotypes in multiple constant temperatures (2005). The difference in days to flower between 23°C long day conditions and 16°C long day conditions plotted against the time to flower for each ecotype. Points are coded based on whether they are known to have an extremely weak or nonfunctional *FRIGIDA* allele. Unknown genotypes were assumed to be high *FRI* because that is the ancestral allelic state. Regressions were performed for each presumed *FRI* status category separately.

Lempe J, Balasubramanian S, Sureshkumar S, Singh A, Schmid M, Weigel D. 2005. Diversity of flowering responses in wild *Arabidopsis thaliana* strains. *PLoS Genetics* **1**(1): 109-118.



Figure S10: Comparison of days to bolt data from high irradiance (HI: 280-300 um) and low irradiance (LI: 130-150 um) experiments (Exp. 3 and 1 respectively). White backgrounds indicate the 22ConLD treatments, light gray backgrounds indicate 22VarLD treatment, and dark gray indicate the 12ConLD treatments. Because 12Var was not tested in Exp. 3 we cannot provide that comparison. Error bars refer to the standard error and asterisks indicate significant comparisons. As these data were collected in separate experiments they should be interpreted cautiously.

Table S1: Description of genotypes, sources, and in what experiment they were used.

Genotype Abr.	Detailed description	Back-ground	Locus	Exp 1	Exp 2	Exp 3	Source
agl24-1	AGAMOUS-LIKE 24	Col	AT4G24540	yes			R. Amasino
co (SM)	CONSTANS	Col	AT5G15840			yes	S. Michaels
co FRI (SM)	CONSTANS + FRIGIDA from San Feliu	Col FRI SF2	AT5G15840			yes	S. Michaels
co-2	CONSTANS	Ler	AT5G15840	yes			ABRC stock center
Col (Ama)	Columbia	Col		yes	yes	yes	R. Amasino
Col (Lis)	Columbia	Col-4		yes			C. Lister (RIL parent)
Col FRI	Columbia with FRIGIDA from San Feliu ecotype	Col FRI SF2	AT4G00650	yes	yes	yes	R. Amasino
Col FRI (SM)	Columbia with FRIGIDA from San Feliu ecotype	Col FRI SF2	AT4G00650			yes	S. Michaels
cry1	CRYPTOCHROME1	Ler	AT4G08920	yes			ABRC stock center
cry1-304	CRYPTOCHROME1	Col-0	AT4G08920	yes		yes	C. Lin
cry1-304/cry2	CRYPTOCHROME1/CRYPTOCHROME2	Col	see single mutants			yes	C. Lin
cry2-1	CRYPTOCHROME2	Col-4	AT1G04400	yes		yes	C. Lin
fca-6	FCA	Ler	AT4G16280	yes			ABRC stock center
fca-9	FCA	Col	AT4G16280			yes	S. Michaels
fca-9 FRI	FCA + FRIGIDA from San Feliu	Col FRI SF2	AT4G16280			yes	S. Michaels
fkf1-2	FLAVIN-BINDING KELCH DOMAIN F BOX PROTEIN	Col	AT1G68050	yes			T. Imaizumi (originally from B. Bartel)
flc3 fri	FLOWERING LOCUS C	Col	AT5G10140	yes			R. Amasino
flc3 FRI	FLOWERING LOCUS C/ FRI	Col FRI SF2	AT5G10140	yes			R. Amasino
fld-3	FLOWERING LOCUS D	Col	AT3G10390			yes	S. Michaels
fld-3 FRI	FLOWERING LOCUS D	Col FRI SF2	AT3G10390			yes	S. Michaels
flk-1	FLOWERING LOCUS KH DOMAIN	Col	AT3G04610	yes			C. Lin
flm-97	FLOWERING LOCUS M	Col	AT1G77080	yes			(?)
fpa-2	FPA	Ler	AT2G43410	yes			ABRC stock center
fpa-7	FPA	Col	AT2G43410			yes	S. Michaels
fpa-7 FRI	FPA + FRIGIDA from San Feliu	Col FRI SF2	AT2G43410			yes	S. Michaels
frl1-3 FRI	FRIGIDA LIKE 1	Col FRI SF2	AT5G16320	yes			R. Amasino
ft-10	FLOWERING LOCUS T	Col	AT1G65480	yes		yes	P. Wigge
ft-2	FLOWERING LOCUS T	Ler	AT1G65480	yes			ABRC stock center
fve-3	FVE	Col	AT2G19520	yes			J. Martinez-Zapater
fve-4	FVE	Col	AT2G19520			yes	S. Michaels
fve-4 FRI	FVE + FRIGIDA from	Col FR	AT2G19520			yes	S. Michaels

	San Feliu	SF2				
gai-d1	GIBBERELLIC ACID INSENSITIVE	Ler tt1	AT1G14920	yes		N. Harberd
gi-2	GIGANTEA	Col-1	AT1G22770	yes	yes	ABRC stock center
gi-6	GIGANTEA	Ler	AT1G22770	yes		ABRC stock center
hua2-3 FRI	HUA2+ FRIGIDA from San Feliu	Col FR SF2	AT5G23150	yes		R. Amasino
hua2-3 fri	HUA2	Col	AT5G23150	yes		R. Amasino
ld-1	LUMIDEPENDENS	Col	AT4G02560	yes		ABRC stock center
ld-1 (SM)	LUMIDEPENDENS	Col	AT4G02560		yes	S. Michaels
ld-1 FRI	LUMIDEPENDENS	Col FR SF2	AT4G02560		yes	S. Michaels
Ler (Lis)	Landsberg ecotype	Ler-0		yes		C. Lister (RIL parent)
Ler-1	Landsberg ecotype	Ler-1		yes		ABRC stock center
phyA-2	PHYTOCHROME A	Ler	AT1G09570	yes		G. Whitelam
phyB-1	PHYTOCHROME B	Ler	AT2G18790	yes		ABRC stock center (?)
phyC-2	PHYTOCHROME C	Col-0	AT5G35840	yes		C. Weinig
phyD-1	PHYTOCHROME D	Ler	AT4G16250	yes		R. Sharrock
phyE-1	PHYTOCHROME E	Ler	AT4G18130	yes		R. Sharrock
prmt5-3 FRI	PROTIEN METHYLTRANSFERASE5	Col FR SF2	AT4G31120	yes		R. Amasino
soc1-2	SUPPRESSOR OF OVEREXPRESION OF CONSTANS	Col	AT2G45660	yes		I. Lee
spy-3	SPINDLY-3	Col	AT3G11540	yes		N. Olszewski
svp-41	SHORT VEGETATIVE PHASE	Col	AT2G22540	yes		P. Huijser
tfl1-2	TERMINAL FLOWER1	Ler	AT5G03840	yes		ABRC stock center
tfl2-6	TERMINAL FLOWER 2	Col	AT5G17690	yes		ABRC stock center
tsf-1	TWIN SISTER OF CONSTANS	Col	AT4G20370	yes		R. Amasino
vin3-4 -fri	VERNALIZATION INSENSITIVE 4	Col	AT5G57380	yes	yes	R. Amasino
vin3-4 FRI	VERNALIZATION INSENSITIVE 3 + FRIGIDA from San Feliu	Col FR SF2	AT5G57380	yes	yes	R. Amasino

Notes: ABRC is the Arabidopsis Resource Center. Question marks indicate lines for which we are unsure of the source.

Table S2: Likelihood ratio test to determine if fluctuating temperatures influence bolting times of wild type genotypes (Col and Ler). Data are from Exp. 1 and chamber number was included as a random effect in the models. All tests had one degree of freedom. Significant contrasts (at the .05 level) after correction for multiple tests would be highlighted in grey but here none meet that criteria. Coef is the coefficient estimate, SE refers to the standard error, and X^2 to the Chi-squared statistic.

Genotype	Treatment	Con mean	Var mean	Coef	SE	X^2	P-value	Corrected P-value
Col Lis	22LDNV	20.9	20.8	-0.006	0.044	0.019	0.890	0.383
Col Lis	22SDNV	64.9	58.5	-0.105	0.056	2.709	0.100	1.000
Col Lis	12SDNV	81.8	75.0	-0.080	0.099	0.610	0.435	1.000
Col Lis	12SDNV	81.8	75.0	-0.080	0.099	0.610	0.435	1.000
Col Ama	22LDNV	20.0	20.9	0.042	0.036	1.323	0.250	1.000
Col Ama	22SDNV	65.6	60.3	-0.089	0.104	0.689	0.406	1.000
Col Ama	22LDV	16.9	18.4	0.085	0.029	5.604	0.018	1.000
Col Ama	22SDV	51.4	44.8	-0.137	0.047	4.892	0.027	1.000
Col Ama	12LDNV	31.6	32.6	0.034	0.037	0.762	0.383	1.000
Col Ama	12SDNV	80.6	73.3	-0.093	0.074	1.335	0.248	1.000
Col Ama	12LDV	26.3	27.3	0.038	0.036	0.976	0.323	1.000
Col Ama	12SDV	65.1	63.3	-0.026	0.077	0.110	0.741	1.000
Ler Lis	22LDNV	18.5	19.6	0.060	0.029	2.874	0.090	1.000
Ler Lis	22SDNV	49.6	36.4	-0.310	0.052	10.374	0.001	1.000
Ler Lis	12SDNV	66.0	63.4	-0.034	0.095	0.123	0.725	1.000
Ler Lis	12SDNV	66.0	63.4	-0.034	0.095	0.123	0.725	1.000
Ler-1	22LDNV	19.0	20.4	0.069	0.036	2.626	0.105	1.000
Ler-1	22SDNV	53.5	38.5	-0.330	0.056	10.337	0.001	1.000
Ler-1	12SDNV	69.1	63.4	-0.074	0.102	0.494	0.482	1.000
Ler-1	12SDNV	69.1	63.4	-0.074	0.102	0.494	0.482	1.000

Table S3: Likelihood ratio tests to determine which allelic changes altered plant responses to fluctuating treatments (Genotype x Fluctuation term). The response of each allelic variant was compared to the response of its respective wild type background (Col or Ler). Data from Experiment 1 was used and chamber type was included as a random effect. All tests have 1 degree of freedom. Significant contrasts (at the .05 level) after correction for multiple tests are highlighted in grey. Geno is short for genotype, Coef is the coefficient estimate, SE refers to the standard error, and X^2 to the Chi-squared statistic.

Treatment	Genotype	Background	Geno x Fluc Coef	Geno x Fluc.SE	X^2	P-value	Corrected P-value
12LDNV	agl24-1	Col	-0.001	0.045	0.000	0.986	1.000
12LDNV	co-2	Ler	-0.087	0.028	8.634	0.003	0.492
12LDNV	Col FRI	Col	-0.019	0.040	0.218	0.640	1.000
12LDNV	cry1	Ler	-0.035	0.030	1.350	0.245	1.000
12LDNV	cry1-304	Col	0.024	0.038	0.384	0.536	1.000
12LDNV	cry2-1	Col	-0.025	0.040	0.394	0.530	1.000
12LDNV	fca-6	Ler	-0.065	0.028	5.207	0.022	1.000
12LDNV	fkf1-2	Col	0.032	0.040	0.607	0.436	1.000
12LDNV	flc-3	Col	0.034	0.039	0.765	0.382	1.000
12LDNV	flc-3 FRI	Col	0.033	0.041	0.652	0.420	1.000
12LDNV	flk-1	Col	-0.043	0.039	1.187	0.276	1.000
12LDNV	flm-97	Col	-0.016	0.042	0.142	0.706	1.000
12LDNV	fpa-2	Ler	-0.028	0.027	1.091	0.296	1.000
12LDNV	frl1-3 FRI	Col	0.084	0.051	2.617	0.106	1.000
12LDNV	ft-2	Ler	-0.067	0.030	4.679	0.031	1.000
12LDNV	fve-3	Col	-0.028	0.038	0.540	0.462	1.000
12LDNV	gai-d1	Ler	-0.016	0.038	0.174	0.676	1.000
12LDNV	gi-2	Col	-0.092	0.049	3.357	0.067	1.000
12LDNV	gi-6	Ler	-0.081	0.029	7.043	0.008	1.000
12LDNV	hua2-3	Col	0.057	0.040	1.982	0.159	1.000
12LDNV	hua2-3 FRI	Col	0.024	0.058	0.164	0.685	1.000
12LDNV	ld-1	Col	0.119	0.040	7.991	0.005	0.686
12LDNV	phyA-2	Ler	-0.059	0.033	3.165	0.075	1.000
12LDNV	phyB-1	Ler	-0.014	0.029	0.222	0.638	1.000
12LDNV	phyC-2	Col	-0.018	0.038	0.222	0.637	1.000
12LDNV	phyD-1	Ler	0.048	0.030	2.524	0.112	1.000
12LDNV	phyE-1	Ler	-0.098	0.032	8.622	0.003	0.492

12LDNV	prmt5 FRI	Col	0.033	0.039	0.700	0.403	1.000
12LDNV	spy-3	Col	0.006	0.053	0.014	0.907	1.000
12LDNV	svp-41	Col	-0.074	0.037	3.782	0.052	1.000
12LDNV	tfl1-2	Ler	-0.020	0.032	0.377	0.539	1.000
12LDNV	tfl2-6	Col	-0.028	0.037	0.589	0.443	1.000
12LDNV	tsf-1	Col	0.055	0.041	1.771	0.183	1.000
12LDNV	vin3-4	Col	-0.022	0.037	0.353	0.552	1.000
12LDNV	vin3-4 FRI	Col	-0.020	0.037	0.285	0.593	1.000
12LDV	Col FRI	Col	-0.034	0.036	0.876	0.349	1.000
12LDV	flc-3	Col	-0.005	0.031	0.031	0.860	1.000
12LDV	flc-3 FRI	Col	0.053	0.046	1.267	0.260	1.000
12LDV	frl1-3 FRI	Col	0.003	0.037	0.007	0.935	1.000
12LDV	hua2-3	Col	-0.002	0.032	0.004	0.949	1.000
12LDV	hua2-3 FRI	Col	0.012	0.037	0.102	0.749	1.000
12LDV	prmt5 FRI	Col	-0.105	0.033	8.780	0.003	0.460
12LDV	vin3-4	Col	-0.026	0.042	0.399	0.527	1.000
12LDV	vin3-4 FRI	Col	-0.011	0.033	0.114	0.735	1.000
12SDNV	agl24-1	Col	0.044	0.024	3.239	0.072	1.000
12SDNV	co-2	Ler	0.008	0.038	0.045	0.831	1.000
12SDNV	Col FRI	Col	0.093	0.034	7.142	0.008	1.000
12SDNV	cry1	Ler	-0.071	0.045	2.465	0.116	1.000
12SDNV	cry1-304	Col	-0.018	0.023	0.584	0.445	1.000
12SDNV	cry2-1	Col	0.032	0.023	1.827	0.176	1.000
12SDNV	fca-6	Ler	0.020	0.035	0.341	0.559	1.000
12SDNV	fkf1-2	Col	0.027	0.026	1.090	0.297	1.000
12SDNV	flc-3	Col	0.008	0.023	0.135	0.713	1.000
12SDNV	flc-3 FRI	Col	0.023	0.023	0.919	0.338	1.000
12SDNV	flk-1	Col	0.080	0.025	9.555	0.002	0.307
12SDNV	flm-97	Col	0.039	0.024	2.707	0.100	1.000
12SDNV	fpa-2	Ler	0.003	0.038	0.006	0.936	1.000
12SDNV	frl1-3 FRI	Col	0.007	0.030	0.061	0.805	1.000
12SDNV	ft-2	Ler	-0.052	0.041	1.579	0.209	1.000
12SDNV	fve-3	Col	0.063	0.029	4.445	0.035	1.000
12SDNV	gai-d1	Ler	-0.018	0.037	0.232	0.630	1.000
12SDNV	gi-2	Col	0.075	0.034	4.699	0.030	1.000
12SDNV	gi-6	Ler	-0.029	0.037	0.601	0.438	1.000
12SDNV	hua2-3	Col	0.021	0.024	0.790	0.374	1.000
12SDNV	hua2-3 FRI	Col	0.022	0.039	0.308	0.579	1.000
12SDNV	ld-1	Col	0.083	0.027	8.739	0.003	0.467
12SDNV	phyA-2	Ler	-0.026	0.038	0.461	0.497	1.000
12SDNV	phyB-1	Ler	0.018	0.038	0.216	0.642	1.000
12SDNV	phyC-2	Col	0.037	0.029	1.587	0.208	1.000
12SDNV	phyD-1	Ler	0.038	0.038	0.999	0.318	1.000

12SDNV	phyE-1	Ler	-0.040	0.038	1.102	0.294	1.000
12SDNV	prmt5 FRI	Col	0.119	0.031	12.478	0.000	0.065
12SDNV	spy-3	Col	0.044	0.035	1.559	0.212	1.000
12SDNV	svp-41	Col	0.020	0.023	0.743	0.389	1.000
12SDNV	tfl1-2	Ler	0.036	0.042	0.723	0.395	1.000
12SDNV	tfl2-6	Col	-0.145	0.045	9.299	0.002	0.351
12SDNV	tsf-1	Col	0.052	0.026	3.816	0.051	1.000
12SDNV	vin3-4	Col	-0.044	0.030	2.030	0.154	1.000
12SDNV	vin3-4 FRI	Col	-0.026	0.033	0.618	0.432	1.000
12SDV	Col FRI	Col	-0.040	0.030	1.720	0.190	1.000
12SDV	flc-3	Col	-0.065	0.031	4.169	0.041	1.000
12SDV	flc-3 FRI	Col	-0.018	0.032	0.307	0.579	1.000
12SDV	frl1-3 FRI	Col	-0.042	0.033	1.545	0.214	1.000
12SDV	hua2-3	Col	-0.011	0.032	0.111	0.738	1.000
12SDV	hua2-3 FRI	Col	-0.009	0.025	0.127	0.722	1.000
12SDV	prmt5 FRI	Col	-0.020	0.035	0.319	0.572	1.000
12SDV	vin3-4	Col	-0.106	0.038	6.791	0.009	1.000
12SDV	vin3-4 FRI	Col	-0.008	0.043	0.038	0.845	1.000
22LDNV	agl24-1	Col	-0.078	0.051	2.262	0.133	1.000
22LDNV	co-2	Ler	-0.030	0.025	1.467	0.226	1.000
22LDNV	Col FRI	Col	-0.399	0.071	24.221	0.000	0.000
22LDNV	cry1	Ler	-0.008	0.026	0.083	0.774	1.000
22LDNV	cry1-304	Col	-0.036	0.049	0.540	0.462	1.000
22LDNV	cry2-1	Col	-0.204	0.059	10.615	0.001	0.175
22LDNV	fca-6	Ler	-0.190	0.030	28.657	0.000	0.000
22LDNV	fkf1-2	Col	-0.334	0.046	35.094	0.000	0.000
22LDNV	flc-3	Col	-0.029	0.047	0.370	0.543	1.000
22LDNV	flc-3 FRI	Col	-0.038	0.043	0.782	0.377	1.000
22LDNV	flk-1	Col	-0.058	0.051	1.238	0.266	1.000
22LDNV	flm-97	Col	0.094	0.061	2.294	0.130	1.000
22LDNV	fpa-2	Ler	-0.030	0.037	0.629	0.428	1.000
22LDNV	frl1-3 FRI	Col	-0.035	0.051	0.465	0.495	1.000
22LDNV	ft-2	Ler	0.191	0.038	20.157	0.000	0.001
22LDNV	fve-3	Col	-0.211	0.053	13.621	0.000	0.035
22LDNV	gai-d1	Ler	-0.025	0.029	0.747	0.387	1.000
22LDNV	gi-2	Col	-0.218	0.052	14.805	0.000	0.019
22LDNV	gi-6	Ler	-0.305	0.036	39.658	0.000	0.000
22LDNV	hua2-3	Col	0.085	0.044	3.617	0.057	1.000
22LDNV	hua2-3 FRI	Col	-0.045	0.065	0.483	0.487	1.000
22LDNV	ld-1	Col	-0.066	0.072	0.844	0.358	1.000
22LDNV	phyA-2	Ler	-0.070	0.027	6.212	0.013	1.000
22LDNV	phyB-1	Ler	0.142	0.035	14.137	0.000	0.027
22LDNV	phyC-2	Col	-0.127	0.054	5.309	0.021	1.000

22LDNV	phyD-1	Ler	-0.024	0.022	1.152	0.283	1.000
22LDNV	phyE-1	Ler	-0.073	0.033	4.577	0.032	1.000
22LDNV	prmt5 FRI	Col	-0.797	0.064	69.803	0.000	0.000
22LDNV	spy-3	Col	0.084	0.072	1.330	0.249	1.000
22LDNV	svp-41	Col	0.140	0.046	8.480	0.004	0.528
22LDNV	tfl1-2	Ler	-0.019	0.029	0.446	0.504	1.000
22LDNV	tfl2-6	Col	0.078	0.047	2.723	0.099	1.000
22LDNV	tsf-1	Col	-0.052	0.057	0.825	0.364	1.000
22LDNV	vin3-4	Col	0.000	0.051	0.000	0.995	1.000
22LDNV	vin3-4 FRI	Col	-0.552	0.072	38.563	0.000	0.000
22LDV	Col FRI	Col	0.026	0.044	0.347	0.556	1.000
22LDV	flc-3	Col	-0.028	0.044	0.396	0.529	1.000
22LDV	flc-3 FRI	Col	-0.070	0.040	2.901	0.089	1.000
22LDV	frl1-3 FRI	Col	0.028	0.049	0.319	0.572	1.000
22LDV	hua2-3	Col	-0.046	0.039	1.371	0.242	1.000
22LDV	hua2-3 FRI	Col	0.066	0.050	1.695	0.193	1.000
22LDV	prmt5 FRI	Col	-0.556	0.055	43.193	0.000	0.000
22LDV	vin3-4	Col	-0.039	0.040	0.931	0.335	1.000
22LDV	vin3-4 FRI	Col	-0.534	0.086	24.672	0.000	0.000
22SDNV	agl24-1	Col	-0.231	0.074	8.790	0.003	0.460
22SDNV	co-2	Ler	0.211	0.046	17.630	0.000	0.004
22SDNV	Col FRI	Col	-0.424	0.066	30.433	0.000	0.000
22SDNV	cry1	Ler	-0.151	0.052	7.898	0.005	0.718
22SDNV	cry1-304	Col	-0.052	0.046	1.282	0.258	1.000
22SDNV	cry2-1	Col	-0.025	0.045	0.326	0.568	1.000
22SDNV	fca-6	Ler	-0.321	0.058	22.635	0.000	0.000
22SDNV	fkf1-2	Col	-0.036	0.052	0.482	0.487	1.000
22SDNV	flc-3	Col	-0.066	0.043	2.381	0.123	1.000
22SDNV	flc-3 FRI	Col	-0.050	0.043	1.320	0.251	1.000
22SDNV	flk-1	Col	-0.324	0.078	14.779	0.000	0.019
22SDNV	flm-97	Col	0.043	0.073	0.348	0.555	1.000
22SDNV	fpa-2	Ler	0.139	0.083	2.717	0.099	1.000
22SDNV	frl1-3 FRI	Col	-0.170	0.067	6.057	0.014	1.000
22SDNV	ft-2	Ler	0.499	0.051	55.067	0.000	0.000
22SDNV	fve-3	Col	0.156	0.057	6.922	0.009	1.000
22SDNV	gai-d1	Ler	-0.012	0.067	0.032	0.858	1.000
22SDNV	gi-2	Col	-0.080	0.112	0.506	0.477	1.000
22SDNV	gi-6	Ler	0.241	0.067	9.787	0.002	0.272
22SDNV	hua2-3	Col	0.067	0.041	2.590	0.108	1.000
22SDNV	hua2-3 FRI	Col	-0.008	0.064	0.014	0.907	1.000
22SDNV	ld-1	Col	-0.059	0.061	0.913	0.339	1.000
22SDNV	phyA-2	Ler	-0.002	0.045	0.002	0.964	1.000
22SDNV	phyB-1	Ler	-0.063	0.048	1.676	0.195	1.000

22SDNV	phyC-2	Col	0.052	0.044	1.351	0.245	1.000
22SDNV	phyD-1	Ler	-0.044	0.051	0.738	0.390	1.000
22SDNV	phyE-1	Ler	-0.012	0.043	0.082	0.774	1.000
22SDNV	prmt5 FRI	Col	0.156	0.057	6.922	0.009	1.000
22SDNV	spy-3	Col	-0.053	0.061	0.722	0.396	1.000
22SDNV	svp-41	Col	0.023	0.047	0.237	0.626	1.000
22SDNV	tfl1-2	Ler	-0.120	0.058	4.174	0.041	1.000
22SDNV	tfl2-6	Col	0.023	0.074	0.099	0.753	1.000
22SDNV	tsf-1	Col	0.065	0.055	1.372	0.241	1.000
22SDNV	vin3-4	Col	-0.026	0.045	0.339	0.560	1.000
22SDNV	vin3-4 FRI	Col	-0.371	0.080	17.895	0.000	0.004
22SDV	Col FRI	Col	-0.001	0.034	0.002	0.966	1.000
22SDV	flc-3	Col	0.048	0.032	2.233	0.135	1.000
22SDV	flc-3 FRI	Col	-0.012	0.030	0.170	0.680	1.000
22SDV	frl1-3 FRI	Col	-0.022	0.037	0.353	0.552	1.000
22SDV	hua2-3	Col	0.134	0.053	5.886	0.015	1.000
22SDV	hua2-3 FRI	Col	0.033	0.036	0.870	0.351	1.000
22SDV	prmt5 FRI	Col	-0.399	0.053	32.270	0.000	0.000
22SDV	vin3-4	Col	-0.005	0.040	0.014	0.905	1.000
22SDV	vin3-4 FRI	Col	-0.210	0.077	5.828	0.016	1.000

Table S4: Likelihood ratio tests to determine whether the bolting time of a *VIN3* mutation in the Col FRI_{SF2} background differed from the Col FRI_{SF2} background. Data from Experiment 1 was used and vernalized treatments were omitted. All tests have one degree of freedom and chamber type was included as a random effect in all tests except the 22VarSD test which did not include random effects because one of the chamber replicates were truncated. Significant contrasts (at the .05 level) after correction for multiple tests are highlighted in grey. Geno is short for genotype, Est is the coefficient estimate, SE refers to the standard error, and X^2 to the Chi-squared statistic.

Treatment	Genotype	Background	Geno .Est	Geno .SE	X^2	P-value	Corrected P-value
12ConLD	Col *FRI vin3-4*	Col *FRI*	0.079	0.020	10.787	0.001	0.006
12ConSD	Col *FRI vin3-4*	Col *FRI*	0.384	0.030	35.988	0.000	0.000
12VarLD	Col *FRI vin3-4*	Col *FRI*	0.078	0.023	8.578	0.003	0.017
12VarSD	Col *FRI vin3-4*	Col *FRI*	0.265	0.042	19.569	0.000	0.000
22ConLD	Col *FRI vin3-4*	Col *FRI*	0.107	0.107	0.961	0.327	0.981
22ConSD	Col *FRI vin3-4*	Col *FRI*	-0.033	0.034	0.907	0.341	0.981
22VarLD	Col *FRI vin3-4*	Col *FRI*	-0.047	0.019	4.943	0.026	0.105
22VarSD	Col *FRI vin3-4*	Col *FRI*	0.028	0.153	.045	0.83	0.981

Table S5: Likelihood ratio tests to determine if high average temperatures influence bolting time of Col and Columbia FRI_{SF2} genotypes. All tests have one degree of freedom. Linear models without a random effect were used to reflect the experimental structure. Significant contrasts (at the .05 level) after correction for multiple tests would be highlighted in grey but here none meet that criteria. Coef refers to the estimate of the coeffient in the model, and X^2 to the Chi-squared statistic.

Genotype	Treatment Contrast	Coef	Sum.of.Sq	P-value X^2	Corrected P-value
Col *FRI*	22ConLD vs. 27ConLD	0.033	0.005	0.799	0.799
Col *FRI*	22ConLD vs. 27VarLD	0.121	0.075	0.022	0.086
Col Ama	22ConLD vs. 27ConLD	-0.110	0.045	0.104	0.312
Col Ama	22ConLD vs. 27VarLD	-0.099	0.039	0.139	0.312

Table S6: Likelihood ratio tests querying whether each genotype behaves significantly differently to warm fluctuating conditions as compared to Columbia wildtype (Genotype x Fluctuation). Data are from Experiment 3 and degrees of freedom are 1 for all tests. Significant contrasts (at the .05 level) after correction for multiple tests are highlighted in grey. Geno is short for genotype, Coef is the coefficient estimate, SE refers to the standard error, and X^2 to the Chi-squared statistic.

Treatment	Genotype	Background	Geno x Fluc Coef	Geno x Fluc.SE	X^2	P-value	Corrected P-value
22LDNV	co	Col	-0.087	0.044	3.751	0.053	1.000
22LDNV	co FRI	Col	-1.116	0.085	57.625	0.000	0.000
22LDNV	Col FRI Ama	Col	-0.610	0.078	34.239	0.000	0.000
22LDNV	Col FRI SM	Col	-0.651	0.086	33.024	0.000	0.000
22LDNV	cry1_cry2	Col	0.018	0.044	0.159	0.690	1.000
22LDNV	cry1-304	Col	-0.029	0.037	0.607	0.436	1.000
22LDNV	cry2-1	Col	-0.058	0.038	2.268	0.132	1.000
22LDNV	fca-9	Col	-0.488	0.052	41.869	0.000	0.000
22LDNV	fca-9 FRI	Col	-0.618	0.072	37.798	0.000	0.000
22LDNV	fld-3	Col	-0.383	0.066	22.916	0.000	0.000
22LDNV	fld-3 FRI	Col	-0.510	0.063	35.484	0.000	0.000
22LDNV	flk	Col	-0.177	0.119	2.144	0.143	1.000
22LDNV	fpa-7	Col	-0.264	0.118	4.640	0.031	1.000
22LDNV	fpa-7 FRI	Col	-0.464	0.062	31.355	0.000	0.000
22LDNV	ft-10	Col	0.017	0.043	0.164	0.686	1.000
22LDNV	fve-4	Col	-0.054	0.068	0.618	0.432	1.000
22LDNV	fve-4 FRI	Col	-0.719	0.082	39.428	0.000	0.000
22LDNV	gi-2	Col	-0.325	0.059	21.281	0.000	0.000
22LDNV	ld-1	Col	-0.382	0.101	11.879	0.001	0.033
22LDNV	ld-1 FRI	Col	-0.401	0.051	34.735	0.000	0.000
22LDNV	soc1-2	Col	0.035	0.042	0.688	0.407	1.000
22LDNV	vin3-4 FRI	Col	-0.733	0.079	41.591	0.000	0.000
22LDV	co	Col	-0.133	0.071	3.313	0.069	1.000
22LDV	co FRI	Col	-0.133	0.066	3.786	0.052	1.000
22LDV	Col FRI Ama	Col	-0.067	0.063	1.119	0.290	1.000
22LDV	Col FRI SM	Col	0.042	0.073	0.329	0.566	1.000
22LDV	fca-9	Col	-0.031	0.065	0.224	0.636	1.000
22LDV	fca-9 FRI	Col	-0.020	0.088	0.054	0.815	1.000
22LDV	fld-3	Col	0.006	0.063	0.009	0.926	1.000

22LDV	fld-3 FRI	Col	0.029	0.068	0.180	0.671	1.000
22LDV	flk	Col	0.014	0.068	0.046	0.831	1.000
22LDV	fpa-7	Col	-0.075	0.062	1.432	0.231	1.000
22LDV	fpa-7 FRI	Col	0.075	0.092	0.658	0.417	1.000
22LDV	fve-4	Col	0.088	0.064	1.821	0.177	1.000
22LDV	fve-4 FRI	Col	0.037	0.065	0.315	0.575	1.000
22LDV	ld-1	Col	0.049	0.075	0.436	0.509	1.000
22LDV	ld-1 FRI	Col	-0.024	0.087	0.078	0.780	1.000
22LDV	vin3-4 FRI	Col	-0.496	0.096	19.525	0.000	0.001
22SDNV	co	Col	0.018	0.063	0.084	0.771	1.000
22SDNV	co FRI	Col	-0.461	0.124	11.366	0.001	0.043
22SDNV	Col FRI Ama	Col	-0.546	0.099	21.083	0.000	0.000
22SDNV	Col FRI SM	Col	-0.364	0.108	9.500	0.002	0.117
22SDNV	cry1_cry2	Col	-0.054	0.142	0.142	0.706	1.000
22SDNV	cry1-304	Col	0.132	0.075	2.924	0.087	1.000
22SDNV	cry2-1	Col	0.072	0.060	1.431	0.232	1.000
22SDNV	fca-9	Col	-0.325	0.075	14.336	0.000	0.009
22SDNV	fca-9 FRI	Col	-0.237	0.074	8.609	0.003	0.187
22SDNV	fld-3	Col	-0.199	0.075	6.268	0.012	0.652
22SDNV	fld-3 FRI	Col	-0.096	0.089	1.136	0.286	1.000
22SDNV	flk	Col	-0.181	0.094	3.320	0.068	1.000
22SDNV	fpa-7	Col	-0.037	0.092	0.162	0.687	1.000
22SDNV	fpa-7 FRI	Col	0.083	0.084	0.964	0.326	1.000
22SDNV	ft-10	Col	0.021	0.061	0.116	0.733	1.000
22SDNV	fve-4	Col	-0.249	0.094	6.312	0.012	0.647
22SDNV	fve-4 FRI	Col	-0.050	0.103	0.220	0.639	1.000
22SDNV	gi-2	Col	-0.144	0.180	0.627	0.429	1.000
22SDNV	ld-1	Col	-0.203	0.118	2.826	0.093	1.000
22SDNV	ld-1 FRI	Col	-0.140	0.098	1.957	0.162	1.000
22SDNV	soc1-2	Col	-0.034	0.060	0.312	0.576	1.000
22SDNV	vin3-4 FRI	Col	-0.394	0.071	19.126	0.000	0.001
22SDV	co	Col	0.007	0.059	0.013	0.910	1.000
22SDV	co FRI	Col	0.082	0.053	2.331	0.127	1.000
22SDV	Col FRI Ama	Col	0.064	0.052	1.500	0.221	1.000
22SDV	Col FRI SM	Col	0.135	0.049	6.759	0.009	0.513
22SDV	fca-9	Col	0.030	0.075	0.162	0.687	1.000
22SDV	fca-9 FRI	Col	0.183	0.103	3.017	0.082	1.000
22SDV	fld-3	Col	0.019	0.057	0.107	0.744	1.000
22SDV	fld-3 FRI	Col	0.061	0.059	1.063	0.303	1.000
22SDV	flk	Col	0.094	0.058	2.548	0.110	1.000
22SDV	fpa-7	Col	-0.028	0.081	0.119	0.730	1.000
22SDV	fpa-7 FRI	Col	-0.065	0.092	0.496	0.481	1.000
22SDV	fve-4	Col	0.037	0.055	0.448	0.503	1.000

22SDV	fve-4 FRI	Col	0.019	0.074	0.066	0.798	1.000
22SDV	ld-1	Col	0.085	0.082	1.061	0.303	1.000
22SDV	ld-1 FRI	Col	0.159	0.080	3.719	0.054	1.000
22SDV	vin3-4 FRI	Col	-0.568	0.082	29.065	0.000	0.000

Table S7: Likelihood ratio tests to determine whether each mutation in the Col *FRI*$_{SF2}$ background influenced bolting within each treatment. Col *FRI* SM was used as reference to keep design balanced and because most genotypes were from that source. Tests of short day treatments for extremely late flowering genotypes were omitted due to truncation of data collection. Significant contrasts (at the .05 level) after correction for multiple tests are highlighted in grey. Geno is short for genotype, Coef is the coefficient estimate, SE refers to the standard error, and X^2 to the Chi-squared statistic.

Treatment	Genotype	Background	Geno Coef	Geno SE	X^2	P-value	Corrected P-value
22ConLD	co FRI	Col FRI SM	-0.835	0.092	27.207	0.000	0.000
22ConSD	co FRI	Col FRI SM	0.034	0.132	0.065	0.799	1.000
22VarLD	co FRI	Col FRI SM	-0.374	0.063	18.504	0.000	0.000
22VarSD	co FRI	Col FRI SM	0.131	0.060	3.280	0.070	0.631
22ConLD	Col FRI Ama	Col FRI SM	0.022	0.089	0.061	0.806	1.000
22ConSD	Col FRI Ama	Col FRI SM	-0.065	0.086	0.568	0.451	1.000
22VarLD	Col FRI Ama	Col FRI SM	-0.019	0.059	0.104	0.747	1.000
22VarSD	Col FRI Ama	Col FRI SM	0.116	0.107	1.102	0.294	1.000
22ConLD	fca-9 FRI	Col FRI SM	0.351	0.078	13.215	0.000	0.004
22VarLD	fca-9 FRI	Col FRI SM	0.384	0.069	16.769	0.000	0.001
22ConLD	fld-3 FRI	Col FRI SM	0.323	0.075	12.311	0.000	0.005
22VarLD	fld-3 FRI	Col FRI SM	0.464	0.061	24.459	0.000	0.000
22ConLD	fpa-7 FRI	Col FRI SM	0.555	0.073	21.940	0.000	0.000
22VarLD	fpa-7 FRI	Col FRI SM	0.743	0.071	33.117	0.000	0.000
22ConLD	fve-4 FRI	Col FRI SM	0.485	0.074	20.798	0.000	0.000
22VarLD	fve-4 FRI	Col FRI SM	0.417	0.081	15.756	0.000	0.001
22ConLD	ld-1 FRI	Col FRI SM	0.113	0.064	2.891	0.089	0.710
22VarLD	ld-1 FRI	Col FRI SM	0.364	0.062	18.290	0.000	0.000
22ConLD	vin3-4 FRI	Col FRI SM	0.188	0.090	3.863	0.049	0.543
22ConSD	vin3-4 FRI	Col FRI SM	0.135	0.069	3.488	0.062	0.618
22VarLD	vin3-4 FRI	Col FRI SM	0.106	0.059	2.896	0.089	0.710
22VarSD	vin3-4 FRI	Col FRI SM	0.151	0.087	1.348	0.246	1.000

Table S8: Likelihood ratio tests to determine if fluctuating temperatures influence blade ratio. Vernalized plants were excluded from this analysis. All tests had 1 degree of freedom. Significant contrasts (at the .05 level) after correction for multiple tests (8) are highlighted in grey. Coef is the coefficient estimate, SE refers to the standard error, and X^2 to the Chi-squared statistic. Con and Var refer to the constant and fluctuating temperature treatments respectively.

Genetic Backgrnd	Contrast	Con Mean	Var Mean	Coef	SE	X^2	P-value	Corrected P-value
Col	22ConLD vs. 22VarLD	0.627	0.541	-0.085	0.012	8.709	0.003	0.022
Col	22ConSD vs. 22VarSD	0.591	0.554	-0.033	0.010	5.357	0.021	0.103
Ler	22ConLD vs. 22VarLD	0.697	0.599	-0.102	0.016	7.994	0.005	0.028
Ler	22ConSD vs. 22VarSD	0.675	0.493	-0.151	0.021	9.113	0.003	0.020
Col	12ConLD vs. 12VarLD	0.607	0.606	-0.002	0.013	0.019	0.891	1.000
Col	12ConSD vs. 12VarSD	0.610	0.607	-0.004	0.013	0.080	0.778	1.000
Ler	12ConLD vs. 12VarLD	0.669	0.653	-0.014	0.014	0.921	0.337	1.000
Ler	12ConSD vs. 12VarSD	0.663	0.675	0.021	0.013	1.899	0.168	0.673

Table S9: Likelihood ratio tests for the effect of fluctuating temperatures on hypocotyl lengths for Col, L*er*, and L*er phyB-1* in warm treatments. Measurements were made on plants in Experiment 1. Vernalized plants were excluded from this analysis. The short day contrast for phyB-1 is omitted because hypocotyl length was highly correlated with developmental stage. All tests have 1 degree of freedom. Significant contrasts (at the .05 level) after correction for multiple tests are highlighted in grey. Coef is the coefficient estimate and X^2 to the Chi-squared statistic. Con and Var refer to the constant and fluctuating temperature treatments respectively.

Genotype	Contrast	Con mean	Var mean	Coef	X^2	P-value	Corrected P-value
Col (Lis)	22ConLD vs. 22VarLD	4.580	7.633	1.357	0.584	0.445	0.852
Col (Lis)	22ConSD vs. 22VarSD	5.070	12.978	8.597	27.051	0.000	0.000
Ler-1	22ConLD vs. 22VarLD	6.460	7.410	0.845	0.634	0.426	0.852
Ler-1	22ConSD vs. 22VarSD	7.785	12.978	5.882	20.748	0.000	0.000
phyB-1	22ConLD vs. 22VarLD	11.035	14.575	5.345	13.787	0.000	0.001

Table S10: Likelihood ratio tests to determine if fluctuations at warm temperatures alter above ground biomass at various times throughout development. All tests have 1 degree of freedom. Significant contrasts (at the .05 level) after correction for multiple tests would be highlighted in grey but here none meet that criteria. Coef is the coefficient estimate, SE refers to the standard error, and X^2 to the Chi-squared statistic. Con and Var refer to the constant and fluctuating temperature treatments respectively.

Genotype	Contrast	Plant age	Con mean	Var mean	Coef	SE	X^2	P-val	Corrected P-value
Col	22ConLD vs. 22VarLD	15	0.0014	0.0011	0.00139	0.00011	2.810	0.094	0.780
Col	22ConLD vs. 22VarLD	18	0.0025	0.0019	0.00252	0.00031	1.922	0.166	0.828
Col *FRI*	22ConLD vs. 22VarLD	15	0.0015	0.0012	0.00150	0.00023	0.940	0.332	1.000
Col *FRI*	22ConLD vs. 22VarLD	18	0.0031	0.0021	0.00305	0.00037	2.935	0.087	0.780
Col *FRI*	22ConLD vs. 22VarLD	29	0.0297	0.0304	0.02973	0.00485	0.010	0.920	1.000
Col	22ConSD vs. 22VarSD	18	0.0008	0.0005	0.00083	0.00011	3.410	0.065	0.648
Col	22ConSD vs. 22VarSD	29	0.0092	0.0058	0.00921	0.00127	2.608	0.106	0.780
Col	22ConSD vs. 22VarSD	41	0.0365	0.0275	0.03647	0.00210	6.267	0.012	0.148
Col *FRI*	22ConSD vs. 22VarSD	18	0.0007	0.0006	0.00073	0.00009	1.216	0.270	1.000
Col *FRI*	22ConSD vs. 22VarSD	29	0.0093	0.0065	0.00931	0.00102	2.704	0.100	0.780
Col *FRI*	22ConSD vs. 22VarSD	41	0.0377	0.0292	0.03765	0.00277	4.056	0.044	0.484
Col *FRI*	22ConSD vs. 22VarSD	56	0.0924	0.0782	0.09236	0.00825	1.256	0.262	1.000
Col	22ConLD vs. 22VarLD	15	0.0014	0.0011	0.00139	0.00011	2.810	0.094	0.780
Col	22ConLD vs. 22VarLD	18	0.0025	0.0019	0.00252	0.00031	1.922	0.166	0.828
Col *FRI*	22ConLD vs. 22VarLD	15	0.0015	0.0012	0.00150	0.00023	0.940	0.332	1.000
Col *FRI*	22ConLD vs. 22VarLD	18	0.0031	0.0021	0.00305	0.00037	2.935	0.087	0.780
Col *FRI*	22ConLD vs. 22VarLD	29	0.0297	0.0304	0.02973	0.00485	0.010	0.920	1.000
Col	22ConSD vs. 22VarSD	18	0.0008	0.0005	0.00083	0.00011	3.410	0.065	0.648
Col	22ConSD vs. 22VarSD	29	0.0092	0.0058	0.00921	0.00127	2.608	0.106	0.780

Table S11: Likelihood ratio tests to determine if fluctuations at warm temperatures alter above ground biomass at bolt. Significant contrasts (at the .05 level) after correction for multiple tests are highlighted in grey. All tests have 1 degree of freedom. Geno is short for genotype, Coef is the coefficient estimate, SE refers to the standard error, and X^2 to the Chi-squared statistic. Con and Var refer to the constant and fluctuating temperature treatments respectively.

Genotype	Contrast	Con mean	Var mean	Coef	SE	X^2	P-val	Corrected P-value
Col	22ConLD vs. 22VarLD	0.005	0.003	-0.001	0.001	4.457	0.035	0.045
Col	22ConSD vs. 22VarSD	0.101	0.046	-0.054	0.016	5.577	0.018	0.045
Col FRI	22ConLD vs. 22VarLD	0.312	0.035	-0.277	0.060	7.678	0.006	0.022
Col FRI	22ConSD vs. 22VarSD	0.765	0.292	-0.473	0.141	5.906	0.015	0.045