

1)

```
> 2+2
[1] 4
> exp(-2)
[1] 0.1353353
> log(100, base=10)
[1] 2
> runif(10)
[1] 0.58838130 0.13067830 0.00422777 0.58009178
[5] 0.63687474 0.42720776 0.05906346 0.71045549
[9] 0.23241068 0.54768403
> runif(10)
[1] 0.3512883 0.7293163 0.3892706 0.9280767 0.2861784
[6] 0.2182457 0.8600011 0.6526888 0.3025214 0.2948526
> |
```

2)

```
> x = 2
> x + x
[1] 4
> y = x + 2
> print y
Error: unexpected symbol in "print y"
> Print y
Error: unexpected symbol in "Print y"
> print(y)
[1] 4
> s = "this is a char str"
> print(s)
[1] "this is a char str"
> |
```

3)

```

> temperatures=c(30, 32, 31, 29, 28)
> x <- temperatures[2]
> y <- temperatures[4]
> print(paste("second -", x, "forth -", y))
[1] "second - 32 forth - 29"
>
> sales_Q1 = c(100, 150, 200)
> sales_Q2 = c(120,180, 240)
> sum_1 = sum(sales_Q1)
> sum_2 = sum(sales_Q2)
> difference = sum_1 - sum_2
> print(paste("difference is ", difference))
[1] "difference is  -90"
> grades = c(85, 72, 90, 65, 88)
> grades_above_80 <- grades[grades > 80]
>
> print(grades_above_80)
[1] 85 90 88
>
> time_intervals = seq(0, 24, 3)
> print(time_intervals)
[1] 0 3 6 9 12 15 18 21 24
> prices = c(20, 30, 40, 50)
> prices[3] = 35
> print(prices)
[1] 20 30 35 50
> |

```

4)

```

> x <- c(5, 10, 15, 20)
> y <- c(1, 2, 3, 4)
> print(x+y)
[1] 6 12 18 24
> print(x - y)
[1] 4 8 12 16
> a <- c(2, 4, 6)
> b <- c(1, 3, 5)
> print(a*b)
[1] 2 12 30
> print(a*10)
[1] 20 40 60
> p <- c(100, 200, 300)
> q <- c(2, 4, 5)
>
> print(p/q)
[1] 50 50 60
> m <- c(10, 20, 30)
> n <- c(3, 5, 7)
>
> print(m%%n)
[1] 1 0 2
> v <- c(1, 2, 3, 4)
> print(v^2)
[1] 1 4 9 16
> |


```

```


> data <- c(5, 10, 15, 20, 25, 30, 35)
>
> mean_value <- mean(data)
> print(mean_value)
[1] 20
>
> data <- c(18, 22, 30, 40, 50)
> median_value <- median(data)
> print(median_value)
[1] 30
>
> data <- c(5, 7, 10, 15, 20)
> std_dev <- sd(data)
> print(std_dev)
Error in print(std_dev) : could not find function "print"
> std_dev <- sd(data)
> print(std_dev)
[1] 6.107373
> data <- c(12, 18, 25, 30, 36)
> variance <- var(data)
> print(variance)
[1] 90.2
> data <- c(3, 5, 7, 9, 11, 13, 15)
> quantiles = quantile(data)
> print(quantiles)
 0%  25%  50%  75% 100%
 3   6   9  12  15
> data <- c(100, 200, 300, 400, 500)
> summary = summary(data)
> print(summary)
      Min. 1st Qu.  Median    Mean 3rd Qu.    Max.
    100     200     300     300     400     500
> data <- c(2, 4, 6, 8, 10, 12)
> iqr_value <- IQR(data)
> print(iqr_value)
[1] 5
.
>
> data <- c(10, 20, 30, 40, 50, 60, 70)
> fivenumsum = fivenum(data)
> print(fivenumsum)
[1] 10 25 40 55 70
> data <- c(1, 2, 3, 4, 5, 6, 7, 8, 9, 10)
> summary_data <- summary(data)
> print(summary_data)
      Min. 1st Qu.  Median    Mean 3rd Qu.    Max.
     1.00   3.25   5.50   5.50   7.75  10.00
> data <- c(7, 8, 9, 10, 12, 14, 16)
> specific_quantiles <- quantile(data, probs = c(0.25, 0.5, 0.75))
> print(specific_quantiles)
 25%  50%  75%
 8.5 10.0 13.0
> |

```

6)



7)



8)

```

> 1.1L
[1] 1.1
Warning message:
integer literal 1.1L contains decimal; using numeric value
> typeof(2)
[1] "double"
> typeof(2L)
[1] "integer"
> name = "Anne"
Error: unexpected invalid token in "name = ""
> name = "Anne Mary"
Error: unexpected invalid token in "name = ""
> name = "nne Mary"
> name
[1] "nne Mary"
> name = "Anne"
> name
[1] "Anne"
> 1/0
[1] Inf
> 0/1
[1] 0
> 0/0
[1] NaN
> -2/0
[1] -Inf
> |

```