



The Cost of Control: Cost-effectiveness Analysis of Hybrid Closed-Loop Therapy in Youth

Diabetes Care 2022;45:1971–1980 | <https://doi.org/10.2337/dc21-2019>

Anthony Pease,^{1,2} Emily Callander,¹
Ella Zomer,¹ Mary B. Abraham,^{3–5}
Elizabeth A. Davis,^{3–5}
Timothy W. Jones,^{3–5} Danny Liew,^{1,6} and
Sophia Zoungas^{1,2,6}

OBJECTIVE

Hybrid closed-loop (HCL) therapy is an efficacious management strategy for young people with type 1 diabetes. However, high costs prevent equitable access. We thus sought to evaluate the cost-effectiveness of HCL therapy compared with current care among young people with type 1 diabetes in Australia.

RESEARCH DESIGN AND METHODS

A patient-level Markov model was constructed to simulate disease progression for young people with type 1 diabetes using HCL therapy versus current care, with follow-up from 12 until 25 years of age. Downstream health and economic consequences were compared via decision analysis. Treatment effects and proportions using different technologies to define “current care” were based primarily on data from an Australian pediatric randomized controlled trial. Transition probabilities and utilities for health states were sourced from published studies. Costs were considered from the Australian health care system’s perspective. An annual discount rate of 5% was applied to future costs and outcomes. Uncertainty was evaluated with probabilistic and deterministic sensitivity analyses.

RESULTS

Use of HCL therapy resulted in an incremental cost-effectiveness ratio of Australian dollars (AUD) \$32,789 per quality-adjusted life year (QALY) gained. The majority of simulations (93.3%) were below the commonly accepted willingness-to-pay threshold of AUD \$50,000 per QALY gained in Australia. Sensitivity analyses indicated that the base-case results were robust.

CONCLUSIONS

In this first cost-effectiveness analysis of HCL technologies for the management of young people with type 1 diabetes, HCL therapy was found to be cost-effective compared with current care in Australia.

Type 1 diabetes is challenging to optimally manage. Hypoglycemia poses an acute risk to health and may seriously impact quality of life and productivity, while prolonged exposure to hyperglycemia increases the risk of microvascular and macrovascular complications (1–6). Advancements in insulin pump therapy, continuous glucose monitoring, and educational strategies for self-management have the potential to improve glycemia and reduce complications of diabetes. However, only a minority of young people internationally are meeting glycemic targets despite the progress in technology (7,8). Furthermore, type 1 diabetes is still associated with

¹School Public Health and Preventive Medicine, Monash University, Melbourne, Victoria, Australia

²Monash Health, Melbourne, Victoria, Australia

³Children’s Diabetes Centre, Telethon Kids Institute, The University of Western Australia, Perth, Western Australia, Australia

⁴Department of Endocrinology and Diabetes, Perth Children’s Hospital, Perth, Western Australia, Australia

⁵Division of Paediatrics, within the Medical School, The University of Western Australia, Perth, Western Australia, Australia

⁶Alfred Health, Melbourne, Victoria, Australia

Corresponding author: Sophia Zoungas, sophia.zoungas@monash.edu

Received 27 September 2021 and accepted 18 May 2022

This article contains supplementary material online at <https://doi.org/10.2337/figshare.20063300>.

© 2022 by the American Diabetes Association. Readers may use this article as long as the work is properly cited, the use is educational and not for profit, and the work is not altered. More information is available at <https://www.diabetesjournals.org/journals/pages/license>.

increased mortality, with the highest risk among those with a younger age of onset (9–12).

Hybrid closed-loop (HCL) therapy is currently the most advanced management strategy available for type 1 diabetes. Current-generation HCL therapy integrates insulin pumps and continuous glucose monitoring devices with decision algorithms to automatically increase or decrease basal insulin to help correct hyperglycemia and mitigate hypoglycemia. While the automation of HCL therapy does not eliminate the need for fastidious user oversight, HCL systems do appear to provide the best potential for optimal glycemic outcomes among pediatric and adult populations (13–18).

The high acquisition cost of HCL therapy is a barrier to equitable access for people with type 1 diabetes (18–21). The cost of technology must also be weighed against the significant costs of diabetes and its complications for people with type 1 diabetes, their carers, health care systems, and society in general (22–25). At present, published economic data regarding HCL therapy are scarce (26). Two economic evaluations reported that HCL therapy was cost-effective compared with insulin pump therapy primarily among adults in Sweden and the U.K. (27,28). Our recent cohort-based cost-effectiveness analysis also reported that HCL therapy was likely cost-effective compared with insulin injections and capillary glucose testing among adults with type 1 diabetes in Australia (29). However, we are unaware of cost-effectiveness analyses regarding HCL therapy among young people or that consider a comparator group using insulin pumps or injections with any glucose monitoring system.

We thus aimed to evaluate the cost-effectiveness of HCL therapy compared with current care among young people with type 1 diabetes in Australia by using a patient-level simulation model. Facilitating and informing key aspects of our analyses were the results of an Australian, multicenter, parallel-group, randomized controlled trial (RCT) for the Australian JDRF Closed Loop Research Group (30).

RESEARCH DESIGN AND METHODS

Model Description

Development and reporting of our model followed the Consolidated Health Economic

Evaluation Reporting Standards (CHEERS) statement, the Diabetes Modeling Input Checklist, as well as the Assessment of the Validation Status of Health-Economic decision models (AdViSHE) checklist (Supplementary “Reporting and validation,” Supplementary Tables 4–6, and Supplementary Figs. 1 and 2) (31–33). A patient-level Markov model with annual cycles was constructed to evaluate the cost-effectiveness of HCL therapy compared with current care among young people aged 12 years, with follow-up until 25 years of age. This age range was chosen to mirror the Australian pediatric RCT that informed this economic analysis (30). A secondary analysis was performed to evaluate the cost-effectiveness of HCL therapy compared with current care from 12 years of age until death. Microsimulation was used to allow transition probabilities to vary based on different characteristics. The perspective was that of the Australian health care system. Future costs and outcomes were discounted by 5% annually based on guidelines by the Australian Government Department of Health (34). The primary outcome of interest was the incremental cost-effectiveness ratio (ICER) in terms of net cost per quality-adjusted life year (QALY) gained.

Model Structure

Health states considered by the model comprised type 1 diabetes with no complications or else any combination of diabetic eye disease, diabetic nephropathy, diabetic foot disease, acute myocardial infarction (AMI), stroke, congestive heart failure (CHF), unstable angina, and death. Diabetic eye disease was modeled with stages including “diabetic retinopathy” (level 20–35 according to the Early Treatment Diabetic Retinopathy Study [ETDRS] grading scale), “sight-threatening diabetic retinopathy” (level ≥ 43 on the ETDRS scale), “macular edema” (extrafoveal and/or clinically significant macular edema according to ETDRS classification), or blindness (both eyes) (35). Diabetic nephropathy was modeled with stages including moderately increased albuminuria, severely increased albuminuria, and end-stage renal disease. Regression of eye disease or renal disease was not modeled directly. Diabetic foot disease was modeled with health states including uncomplicated diabetic foot ulcer, complicated/infected

diabetic foot ulcer, healed diabetic foot ulcer, minor amputation (at or below the ankle), infected amputation, major amputation (above the ankle), and healed amputation (36). The interrelated nature of complications was captured by transition probabilities for health states using the shared risk factor of HbA_{1c} as well as incorporating renal health states as part of annual cardiovascular risk prediction. Tracker variables were implemented to facilitate the independent modeling of different health states. Mortality was modeled to be a result of cardiovascular disease, renal causes, or “other” causes. Simulated individuals continued to cycle through the model until 25 years of age in the primary analysis and up to death with a theoretical limit of 100 years of age in the secondary analysis. Supplementary Fig. 2 graphically presents the model structure.

Transition Probabilities and Mortality Risk

Transition probabilities were derived from published sources to simulate disease progression (Supplementary Table 1). Contemporary published data were sought for each modeled health state, and age-specific all-cause mortality was drawn from published Australian data for people with type 1 diabetes (37).

Intervention Being Modeled

The treatment effect of HCL therapy on HbA_{1c} determined the difference in transition probabilities between treatment groups for all vascular complications. Rates of severe and nonsevere hypoglycemia were modeled to reflect the possibility of having multiple hypoglycemic events in a year rather than annual transition probabilities per event. All hypoglycemic events were assumed to be nonfatal. To predict the annual probability of the first macrovascular event, the Steno T1 Risk Engine model was used and involved the variables of age, sex, diabetes duration, systolic blood pressure, LDL cholesterol, HbA_{1c}, presence or absence of moderately or severely increased albuminuria, estimated glomerular filtration rate (eGFR) categorized as >40 or <40 mL/min/1.73 m², smoking status, and participation in regular exercise (38). Owing to the population-level data source for renal health states reporting on albuminuria or end-stage renal disease rather than eGFR levels, it was pragmatically

assumed that eGFR was >40 mL/min/ 1.73 m² when using the Steno T1 Risk Engine model unless patients transitioned to end-stage renal disease. Recurrent cardiovascular events had to be preceded by a cardiovascular event at least 1 year earlier, and renal death could only occur in the presence of renal pathology arising at least 1 year prior. To predict the probability of recurrent cardiovascular events, the results of “A new model for 5-year risk of CVD in Type 1 diabetes” were annualized (39). This cardiovascular risk prediction model includes variables of age, diabetes duration, HbA_{1c}, systolic blood pressure, total cholesterol-to-HDL cholesterol ratio, smoking status, presence or absence of severe albuminuria, and the presence of a previous event (39). Random numbers were generated to determine patient sex, smoking status, and participation in regular exercise according to cohort-level probabilities as inputs for the cardiovascular risk prediction models. Results from the Diabetes Control and Complications Trial (DCCT)/Epidemiology of Diabetes Interventions and Complications (EDIC) study were used to define the proportion of events comprising AMI, stroke, or CHF (40). Case fatality data were used to derive mortality rates for AMI, stroke, and CHF. Unstable angina and lower-extremity amputations were assumed to be nonfatal. Because projected cardiovascular and renal mortality rates were modeled separately in relation to glycemic control, previously reported estimates for these causes of death were subtracted from the published aggregate of Australian all-cause mortality data to avoid double counting.

Costs

All costs were reported in 2021 Australian dollars (AUD), with adjustment for inflation using the health consumer price index (41). Multiple sources were used to estimate costs (Supplementary Table 2). For type 1 diabetes without any complications, costs were sourced from the largest Australian cost-of-illness study comprising 2,200 people with type 1 diabetes selected from an Australian diabetes register (National Diabetes Services Scheme [NDSS]) (22). For diabetic retinopathy among those with type 1 diabetes in Australia, costs were estimated based on Medicare Benefits Schedule item numbers, the Pharmaceutical Benefits Scheme for medications, and the intensity of

monitoring and intervention was based on clinical guidelines, landmark clinical trials, and the expert opinion of ophthalmologists. For diabetic nephropathy, costs were derived from Australian economic projections published by Kidney Health Australia reflecting the cost of kidney disease for the general population (42,43). For diabetic foot disease, costs were derived from an Australian cost-effectiveness analysis regarding the optimal care for diabetic foot ulcers among those with type 1 or type 2 diabetes (36). Costs for the inpatient and subsequent outpatient management of stroke, regardless of diabetes status, were derived from Australian cost-effectiveness analyses (44,45), and the costs for other cardiovascular diseases among the general population were derived from a report by the Australian Heart Foundation (46). A conservative 50% reduction was applied to costs for fatal cardiovascular events to account for only 50% of deaths occurring in hospitals throughout Australia, regardless of diabetes status (47,48). For severe hypoglycemia, an Australian economic evaluation was used that comprised costs for ambulance callout, hospital admission, accident and emergency visit, glucagon injection, and follow-up visits (49). Nonsevere hypoglycemic events were assumed to have no associated costs, but an annual disutility was applied.

Regarding intervention costs, the analysis included incremental costs of an HCL system over current care, which was defined by the baseline utilization rates of insulin pump therapy or multiple daily injections of insulin with either capillary glucose testing or continuous glucose monitoring in the Australian pediatric RCT (30). The only commercially available HCL systems in Australia at the time of writing comprised the MiniMed 670G/770G (Medtronic), infusion sets, reservoirs, batteries, Guardian Sensor 3 (Medtronic), transmitter, and transmitter charger. Twice-daily capillary glucose testing was assumed for calibration when using the HCL system or continuous glucose monitoring compared with eight tests per day for capillary glucose testing. Among those in the comparator arm using continuous glucose monitoring apart from HCL therapy, it was assumed that available and subsidized brands (AMSL Diabetes and Medtronic Diabetes Australia) were used in equal proportions. Cohort-level utilization rates of insulin pumps,

injections, capillary glucose testing, and continuous glucose monitoring of those ≤ 25 and >25 years of age derived an overall weighted cost for management in the comparator arm. The previously described Australian cost of illness study presented aggregated costs of type 1 diabetes that captured the cost of multiple daily injections and capillary glucose testing. Consistent with current government subsidies, the cost of insulin pump consumables, but not insulin pumps, was added to the aggregated cost of type 1 diabetes among those using insulin pump therapy in the comparator arm.

Utilities

Utilities for the health state of type 1 diabetes without complications were drawn from the same Australian cost of illness study described above (22). In the absence of Australian data sources and sparsity of detail internationally, utilities for the different stages of diabetic eye disease were drawn from Japanese and Canadian patient preference-based time trade-off analyses that informed prior cost-effectiveness analyses (50–52). To facilitate the inclusion of separate utilities for albuminuria as well as hemodialysis and peritoneal dialysis, European analyses of people with type 2 diabetes or on chronic dialysis were implemented and were consistent with the available more general data regarding people with type 1 diabetes (53–55). To facilitate the inclusion of separate utilities for diabetic foot ulceration, infection, as well as minor or major amputations, with or without infection, European time trade-off analyses were chosen that had been used previously in Australian cost-effectiveness analyses (36,56). Disutilities for AMI, stroke, CHF, and angina were taken from a population-based time trade-off analysis among people with type 2 diabetes in the U.K. (57). Annual disutilities of -0.0475 and -0.0041 were used for severe and nonsevere hypoglycemia, respectively. Disutility for hypoglycemia was derived as the average of daytime and nocturnal events from a large Canadian time trade-off survey among people with type 1 diabetes (58). Disutilities for multihealth conditions were summed. See Supplementary Table 3 for a list of utilities/disutilities used in the analysis.

Simulation Cohort and Treatment Effects

The model was populated by convention with 10,000 individual patient-level simulations profiled largely from a 6-month Australian pediatric RCT of HCL therapy compared with current care (30). Individuals entered the model at 12 years of age, with diabetes duration of 7 years, and with no vascular complications. Rates of severe and nonsevere hypoglycemia among the pediatric population and those modeled to continue through adulthood were drawn from large registries, the DCCT/EDIC study, cohort studies as well as the 6-month Australian RCT (2,30,59–67). Treatment effects for HCL therapy on HbA_{1c} and nonsevere hypoglycemia reduction were drawn from the same Australian pediatric RCT and were assumed to be constant over follow-up for primary and secondary analyses (30). HCL therapy was also conservatively assumed to reduce severe hypoglycemia by 50% in the base-case. Baseline utilization of technology types among participants in the Australian pediatric RCT defined the comparator arm for our economic evaluation, were assumed to remain constant until 25 years of age, and comprised insulin pump therapy (81%), multiple daily injections of insulin (19%), continuous glucose monitoring (59%), and capillary glucose testing (41%) (30). For the secondary analyses that followed up individuals until death, clinical profiles and utilization rates of technology types after 25 years of age were based on data from the Australian National Diabetes Audit, which has reported that 27% and 23% of adults with type 1 diabetes used insulin pumps and continuous glucose monitoring, respectively, and were also assumed to remain constant over time (20). Cohort-level mean (\pm SD) values for biomarkers required for cardiovascular risk prediction in adulthood were also derived from the Australian National Diabetes Audit and were assumed to be equivalent in the treatment and comparator groups (68). Broadly in keeping with current funding initiatives in Australia, those using continuous glucose monitoring were assumed to receive no government subsidy after 25 years of age, unless part of the modeled intervention of HCL therapy (20). Those using insulin pumps apart from HCL therapy were also assumed to receive the current government subsidy that is limited to consumables (reservoirs

and cannulas) and insulin regardless of age. A detailed profile of the simulated individuals and treatment effects is provided in Table 1.

Sensitivity Analyses

Probabilistic sensitivity analyses with 10,000 iterations were performed to account for uncertainty across the input parameters. Univariate scenario analyses were also considered to explore model assumptions that comprised extending the horizon from 13 years to a lifetime; reducing discount rates for future costs and outcomes to 3.5%, 1.5%, and 0%; increasing and decreasing the cost of HCL therapy by 10%; reducing the treatment effect of HCL therapy on nonsevere hypoglycemia by 25% and 50%; increasing the baseline rate of nonsevere hypoglycemia to 104 events per person/year; reducing the treatment effect of HCL therapy on severe hypoglycemia by 50% and 100%; assuming no treatment effect on HbA_{1c}; assuming the baseline HbA_{1c} was 7.0% (53 mmol/mol) or 9.0% (75 mmol/mol); assuming the baseline incidence of severe hypoglycemia was 16.6 or 6.7 per 100 person-years based on local population studies (64,69); and assuming the treatment effect on HbA_{1c} increased from 0.3% (3.3 mmol/mol) to 1.0% (10.9 mmol/mol) over a lifetime horizon with the baseline HbA_{1c} unchanged or else increased to 9.0% (75 mmol/mol). All univariate scenario analyses were replicated over a lifetime horizon, except for assuming reduced base rates of severe hypoglycemia based on results reported among a pediatric population. In addition, the treatment effect of HCL therapy on HbA_{1c} was only increased from 0.3% (3.3 mmol/mol) to 1.0% (10.9 mmol/mol) among adults over a lifetime horizon based on results of meta-analyses among adults (13,14). Microsoft Excel (Microsoft Corp., Redmond, WA) and TreeAge Pro Healthcare 2021 R1.1 (TreeAge Software LLC, Williamstown, MA) were used for the economic analyses.

RESULTS

Compared with current care, the use of HCL therapy was associated with gains of 1.15 (discounted) QALYs per person until 25 years of age, at a net cost (discounted) of AUD \$37,827, resulting in

an ICER of AUD \$32,789 per QALY gained. At the commonly accepted willingness-to-pay (WTP) threshold of AUD \$50,000 per QALY gained in Australia (70–72), 93.3% of simulations predicted the HCL system to be cost-effective.

Sensitivity analyses found that results were most sensitive to the base rate and treatment effect on nonsevere hypoglycemia, the cost of HCL systems, the time horizon, and the treatment effect of HCL therapy on HbA_{1c}. Assuming the base rate of nonsevere hypoglycemia was increased to 104 events per person/year in keeping with the rates reported by the International Society for Pediatric and Adolescent Diabetes (ISPAD) Clinical Practice Consensus Guidelines for hypoglycemia (66), led to an ICER of AUD \$18,436 per QALY gained while reducing the treatment effect of HCL therapy on nonsevere hypoglycemia by 50% increased the ICER to AUD \$54,668 per QALY gained. In contrast, assuming no treatment effect on severe hypoglycemia for people aged 12 to 25 years increased the ICER only slightly to AUD \$37,452 per QALY gained. Reducing or increasing the cost of HCL therapy by 10% led to ICERs of AUD \$24,234 or AUD \$40,864 per QALY gained, respectively.

Extending the time horizon to a lifetime increased the ICER to AUD \$48,882 per QALY gained. In the context of a lifetime horizon, increasing the relative treatment effect of HCL therapy on HbA_{1c} from -0.3% (3.3 mmol/mol) to -1.0% (10.9 mmol/mol) reduced the ICER to AUD \$42,271 per QALY gained, or AUD \$43,136 per QALY gained if the baseline HbA_{1c} was also increased to 9.0% (75 mmol/mol). Results over a lifetime were also particularly sensitive to the cost of HCL therapy as well as the base rate of nonsevere hypoglycemia and the treatment effect of HCL therapy on nonsevere hypoglycemic events. Reducing the recommended 5% discount rate for future costs and health outcomes increased the ICER. Results of the base-case and scenario analyses are presented in Tables 2 and 3, respectively.

The cost-effectiveness plane and scatter plot (Fig. 1) presents the variation in modeled outcomes, with HCL therapy tending to offer greater effectiveness in QALYs but at a greater cost than current care. The ellipse in Fig. 1 represents the 95% confidence for modeled results. The cost-effectiveness acceptability curves for the primary and secondary analyses

Table 1—Characteristics of patient-level simulations

Cohort characteristics	Mean (variance)	α	β	Distribution	Reference
Initial age, years	12.00			Fixed	Abraham et al. (30)
Duration of diabetes, years	7.00			Fixed	Abraham et al. (30)
Female sex, %	55.00			Fixed	Abraham et al. (30)
Total-to-HDL cholesterol ratio	3.09			Fixed	Pease et al. (68)
LDL cholesterol, mean (SD), mmol/L	2.55 (0.95)			Normal	Pease et al. (68)
HbA _{1c} (age ≤ 21 years), mean (SD), %	8.0 (1.0)			Normal	Abraham et al. (30)
HbA _{1c} (age ≤ 21 years), mmol/mol	64 (10.9)			Normal	Abraham et al. (30)
HbA _{1c} (age > 21 years), mean (SD), %	8.5 (1.5)			Normal	Pease et al. (68)
HbA _{1c} (age > 21 years), mean (SD), mmol/mol	69 (16.4)			Normal	Pease et al. (68)
Systolic blood pressure, mmHg	124 (17)			Normal	Pease et al. (68)
Smoker, %	38.50	397	634	β	Pease et al. (68)
Regular exercise, %	68.20	2,937	1,369	β	Vistisen et al. (38)
Nonsevere hypoglycemia (overall), n/person/year	55.70	100.00	0.56	γ	Ratzki-Leewing et al. (60)
Severe hypoglycemia, n/person/year					
Total—age ≤ 21 years	0.38	100.00	0.00	γ	Gubitosi-Klug et al. (67)
Total—age > 21 years*	1.98	100.00	0.02	γ	Pease et al. (29), Geddes et al. (59)
Hypoglycemia awareness—age > 21 years					
Normal	1.00				
Impaired	6.00				
Proportion with impaired hypoglycemia awareness (age > 21 years), %	19.50				
Treatment effects of HCL therapy					
HbA _{1c} reduction, %	0.3			Fixed	Abraham et al. (30)
HbA _{1c} reduction, mmol/mol	3.3			Fixed	Abraham et al. (30)
Nonsevere hypoglycemia reduction, %	54			Fixed	Abraham et al. (30), Pease et al. (29)
Severe hypoglycemia reduction, %					
Age ≤ 25 years	50			Fixed	Assumption
Age > 25 years	95			Fixed	Pease et al. (29)

*The number of severe hypoglycemic events per person/year represented the average weighted number of events estimated from the prevalence of impaired hypoglycemia awareness (19.5%) and the number of severe hypoglycemic events for adults > 21 years of age with and without impaired awareness of hypoglycemia.

(Supplementary Fig. 3 and 4, respectively) present the proportion of 10,000 iterations that were cost-effective for HCL therapy and current care at various WTP thresholds. Tornado diagrams of sensitivity analyses of HCL therapy compared with current care are presented in Supplementary Figs. 5 and 6 and provide

further support for our choice of variables and scenarios for deterministic sensitivity analyses presented in Table 3.

CONCLUSIONS

Our modeled economic evaluation produced an ICER of AUD \$32,789 per QALY

gained for HCL therapy relative to current care. HCL therapy is thus likely to be cost-effective compared with current care among young people with type 1 diabetes in Australia based on the commonly accepted WTP threshold of AUD \$50,000 per QALY gained (70–72). Sensitivity analyses demonstrated the results were robust.

Our analyses regarding HCL therapy had a unique focus on young people as well as the comparison with a range of preexisting management devices. However, results were consistent with the few published economic evaluations, primarily among adults, that similarly reported HCL therapy to be cost-effective despite different health care systems, WTP thresholds, modeling perspectives, models, and different clinical comparators (27–29). HCL therapy has been consistently modeled

Table 2—Overall base-case results

Results	Current care	HCL therapy
Costs, mean (SD), \$	90,507 (11,391)	128,334 (12,611)
QALYs, mean (SD)	6.56 (0.81)	7.71 (0.82)
Incremental cost, \$		37,827
Incremental QALYs		1.15
ICER, \$		32,789

The results represent discounted values reported in 2021 AUD.

Table 3—Deterministic scenario analyses

Scenarios	ICER (mean), \$
Primary analysis (12 to 25 years of age)	
Discount rate 3.5% for costs and outcomes	32,816
Discount rate 1.5% for costs and outcomes	32,102
Discount rate 0% for costs and outcomes	31,952
Assuming the cost of HCL therapy increased by 10%	40,864
Assuming the cost of HCL therapy decreased by 10%	24,234
Assuming treatment effect on nonsevere hypoglycemia was reduced by 25%	41,110
Assuming treatment effect on nonsevere hypoglycemia was reduced by 50%	54,668
Assuming baseline rate of nonsevere hypoglycemia was 102 per person/year	18,436
Assuming treatment effect on severe hypoglycemia was reduced by 50%	35,184
Assuming treatment effect on severe hypoglycemia was reduced by 100%	37,452
Assuming baseline rate of severe hypoglycemia was 16.6 per 100 person/years	35,469
Assuming baseline rate of severe hypoglycemia was 6.7 per 100 person/years	36,817
Assuming no treatment effect on HbA _{1c}	32,928
Assuming baseline HbA _{1c} was 1.0% (10.9 mmol/mol) lower	32,928
Assuming baseline HbA _{1c} was 1.0% (10.9 mmol/mol) higher	32,990
Over a lifetime horizon:	48,882
Discount rate 3.5% for costs and outcomes	50,454
Discount rate 1.5% for costs and outcomes	57,909
Discount rate 0% for costs and outcomes	63,691
Assuming the cost of HCL therapy increased by 10%	61,885
Assuming the cost of HCL therapy decreased by 10%	43,501
Assuming treatment effect on nonsevere hypoglycemia was reduced by 25%	65,443
Assuming treatment effect on nonsevere hypoglycemia was reduced by 50%	82,920
Assuming baseline rate of nonsevere hypoglycemia was 102 per person/year	27,882
Assuming treatment effect on severe hypoglycemia was reduced by 50%	49,662
Assuming treatment effect on severe hypoglycemia was reduced by 100%	53,748
Assuming no treatment effect on HbA _{1c}	51,302
Assuming the treatment effect on HbA _{1c} was −1% (10.9 mmol/mol)	42,271
Assuming baseline HbA _{1c} was 1.0% (10.9 mmol/mol) lower	49,609
Assuming baseline HbA _{1c} was 1.0% (10.9 mmol/mol) higher	48,939
Assuming baseline HbA _{1c} was 1.0% higher and the treatment effect on HbA _{1c} was −1%	43,136

The results represent discounted values with annual discounting of 5% for costs and outcomes unless otherwise stated. All costs are reported in 2021 AUD.

to increase QALYs over a lifetime, with the increment ranging from 1.73 and 1.90 compared with insulin pump therapy (27,28), through to 3.72 compared with insulin injections (29) in the U.K., Sweden, and Australia, respectively. This is in keeping with our sensitivity analyses over a lifetime horizon that modeled an increment of 2.15 QALYs compared with current care. However, prior modeled cohorts have been much older, with analyses starting at 18 years of age in Australia (29), or a mean age of 37.8 (SD 16.5) years in the U.K. and Sweden (27,28). Previous analyses have also limited their comparison of HCL systems to insulin pump therapy or insulin injections. Therefore, our focus on the cost-effectiveness of HCL therapy among young people compared with current care adds important breadth to the growing landscape of economic evidence.

In addition to the primary analysis among young people, a secondary analysis using a lifetime horizon was included due to the lifelong nature of type 1 diabetes. Consistent between the primary and secondary analyses was that nonsevere hypoglycemia and the cost of technology remained key drivers for cost-effectiveness. The important role of utility gains from reduced hypoglycemia or “fear” of hypoglycemia as well as technology costs have also been reported by the few other economic evaluations of HCL therapy to date (27–29). Furthermore, because most vascular complications develop in adulthood, the relative impact of HbA_{1c} reductions from HCL therapy on ICER values was also much greater over the lifetime horizon compared with the primary analysis over 13 years. Our conservative analyses over a lifetime horizon produced an ICER of

AUD \$48,882 per QALY gained. Combined with our primary analysis, the results suggest that HCL therapy may be cost-effective in young people and across the life span compared with current care in Australia. Any reductions in technology costs over time would also be predicted to decrease the ICER further. However, base rates of nonsevere hypoglycemia and treatment effects of HCL therapy on HbA_{1c} and nonsevere hypoglycemia over a lifetime may impact these conclusions.

The strengths of our study included patient-level simulations providing more granular detail than cohort-based models and a diverse multidisciplinary team developing the model and ensuring face validity (Supplementary File; Reporting and validation). Treatment effects of HCL therapy for HbA_{1c} and nonsevere hypoglycemia were derived from an Australian RCT (30). The comparator arm used for our analyses may also be more generalizable than other studies because participants and their clinicians were free to use any management technology apart from HCL therapy throughout the trial, reflecting real-world clinical practice (30). In addition, conservative estimates for baseline glycemia were implemented, and extensive sensitivity analyses for key drivers of cost-effectiveness supported our base-case results. Costs and utilities for the baseline health state of type 1 diabetes without complications were sourced from Australia's largest cost of illness study (22), and all other cost data were taken from Australian sources.

We also acknowledge that there were limitations of the available data sources and model assumptions. Outcome projections over a number of years were extrapolated from short-term clinical data in the absence of long-term studies for modern technologies. Like most economic evaluations, generalizability of results may be limited by the implementation of RCT data with recruited and often highly motivated participants. Treatment effects and base rates of severe hypoglycemia were also unclear in the literature. However, data from the landmark DCCT/EDIC study were implemented for base rates of severe hypoglycemia as they were considered conservative compared with population studies and meta-regression of RCTs (1,2,64–67,73,74). Sensitivity analyses also considered lower baseline rates of severe hypoglycemia as reported by other Australian studies (64,69).




Figure 1—Incremental cost-effectiveness plane and scatter plot (12 to 25 years of age). The solid circles and open triangles represent iterations for which HCL therapy compared with current care was the optimal and suboptimal strategy in relation to the WTP threshold, respectively. The bold dotted line represents the WTP threshold, and the ellipse represents 95% confidence.

Regarding the treatment-effect of HCL therapy on severe hypoglycemia, the majority of recent trials have been of short duration and reported zero events among young people, although similarly low rates were present in comparator arms as well. In contrast to other economic evaluations that assumed HCL therapy completely prevented severe hypoglycemia over a lifetime in the base-case (27,28), we were more conservative and considered a treatment effect of 50% with sensitivity analyses ranging from 0 to 95%. As with other published economic models, utility values derived from populations with type 2 diabetes and/or from different countries were implemented when comparable detail was not available for those with type 1 diabetes in Australia. Experienced ophthalmologists also assisted with the development of model assumptions regarding the investigation and management of diabetic eye disease in the absence of adequate publicly available data. Owing to limited available data sources, a number of assumptions were made during model development. The costs of end-stage renal disease and of CVD were based on populations irrespective of comorbid diabetes

status, and the cost of diabetic foot disease was based on a population with any diabetes type. Fatal cardiovascular event costs were conservatively reduced by 50% to account for only 50% of deaths occurring in hospitals throughout Australia (47,48); however, assuming no adjustment for death costs or assuming zero costs for death had minimal impact on the primary or secondary analysis (data not shown). Results were limited to the health care system perspective, although any societal benefits from HCL therapy would likely reduce the ICER further. Similar to other validated models, historical results from the DCCT/EDIC study were used to apportion modeled cardiovascular events as AMI, unstable angina, stroke, or CHF. In addition, eGFR was assumed to be <40 mL/min/1.73 m² only for those modeled to have end-stage renal disease. Deterministic and probabilistic sensitivity analyses addressed uncertainty in the model, but it was not feasible to address all clinically relevant scenarios.

In addition to cost-effectiveness analyses, further research is needed into the indirect costs and budget impact of HCL therapy as well as real-world and patient-

relevant outcomes. Clinical trials and existing diabetes registries could be leveraged to collect standardized economic data to strengthen models, address economic uncertainty, and appraise implementation strategies. The increasing focus on time-in-range over HbA_{1c} as a metric of glycemic control also requires longer-term data regarding the complications of diabetes and quality of life to facilitate inclusion in future economic modeling. Furthermore, while most economic evaluations focus on the perspective of the health care system for funding decisions, the burden of health care costs on people with type 1 diabetes and their carers should also drive further research into health equity.

Overall, HCL therapy is likely cost-effective compared with current care by reducing HbA_{1c} and the incidence of hypoglycemia among those with type 1 diabetes in Australia. This first positive finding among young people adds to the weight of growing international economic evidence among adults that HCL therapy appears to represent good value for money in a relatively short time frame and across the life span. Future research into diabetes management technologies

should also focus on budget impact and incorporating health equity into funding decisions.

Acknowledgments. The authors thank the participating sites for the randomized controlled trial, which comprised Perth Children's Hospital, Nedlands, Western Australia; The Children's Hospital at Westmead, Westmead, New South Wales; Women and Children's Hospital, North Adelaide, South Australia; Royal Children's Hospital, Parkville, Victoria; and John Hunter Children's Hospital, New Lambton Heights, New South Wales. The authors also thank Prof. Peter van Wijngaarden (Centre for Eye Research Australia, Royal Victorian Eye and Ear Hospital, East Melbourne, Victoria, Australia; Ophthalmology, Department of Surgery, The University of Melbourne, Parkville, Victoria, Australia) and Dr. Peter Heydon (Faculty of Medicine and Health, The University of Sydney, Camperdown, New South Wales, Australia; Royal Prince Alfred Hospital, Camperdown, New South Wales, Australia; Liverpool Hospital, Liverpool, New South Wales, Australia) for their expert advice regarding clinical assumptions in the model for diabetic eye disease. This economic evaluation was developed through collaborations as part of the JDRF Global Centre of Excellence in Diabetes Research.

Funding. A.P. was supported by the Royal Australasian College of Physicians/Diabetes Australia Research Establishment Fellowship. Funding and support for the randomized controlled trial, "Effect of a hybrid closed-loop system on glycaemic and psychosocial outcomes in children and adolescents with type 1 diabetes: a randomized clinical trial" comprised: the JDRF Australia Type 1 Diabetes Clinical Research Network grant 4-SRA-2016-350-M-B (T.W.J.), Special Research Initiative of the Australian National Health and Medical Research Council grant APP1078190 (T.W.J.), and PCH Foundation grant 9729 (T.W.J.). The National Health and Medical Research Council Clinical Trials Centre was supported by its National Health and Medical Research Council Program grant. In-kind support was provided by Medtronic through the provision of insulin pumps, sensors, and transmitters for the study. Roche Diabetes Care provided the glucometers for the study. For the randomized controlled trial, M.B.A. was supported by the Department of Health/Raine Clinical Research Fellowship from Western Australia.

JDRF Australia provided input into the study design of the randomized controlled trial. The funders of the trial had no role in collection, analysis, and interpretation of the data and in manuscript preparation.

Duality of Interest. E.Z. has received grants from Amgen, AstraZeneca, Pfizer, and Shire outside the submitted work. M.B.A. reports receiving speaker honoraria fees from Medtronic outside the submitted work. E.A.D. reports participation in educational meetings on behalf of the Telethon Kids Institute for Eli Lilly (payment to institution). D.L. has received grants from AstraZeneca, Pfizer, AbbVie, and Bristol Myer Squibb, and has received personal fees

from AstraZeneca, Astellas, and Bayer outside the submitted work. S.Z. reports participation in advisory boards, expert committees, or educational meetings outside the submitted work on behalf of Monash University for Boehringer-Ingelheim, Eli Lilly, Sanofi, AstraZeneca, Novo Nordisk, and MSD Australia (payment to institution). T.W.J. reports receiving grants from the JDRF for study costs and grants from Perth Children's Hospital Foundation for study costs during the conduct of the study. No other potential conflicts of interest relevant to this article were reported.

Author Contributions. A.P. performed analysis and drafted the manuscript. E.C., E.Z., M.B.A., E.A.D., T.W.J., D.L., and S.Z. assisted manuscript drafting and review. E.C., E.Z., and D.L. also provided advice regarding analysis. All authors read and approved the manuscript. S.Z. is the guarantor of this work and, as such, had full access to all the data in the study and takes responsibility for the integrity of the data and the accuracy of the data analysis.

Prior Presentation. Parts of this study were presented as an abstract at The Australasian Diabetes Congress 2021, virtual event, 11–13 August 2021, and at the First Annual International Hypoglycaemia Study Group Advances in Hypoglycaemia Scientific Conference (<https://www.xcdsystem.com/adsc/program/vPZbWzx/index.cfm?pgid=2327&print=1&printmode=1>), 17–18 November 2021. An abstract was published after submission to The Australasian Diabetes Congress 2021. Pease A, Callander E, Zomer E, et al. Cost-effectiveness analysis of hybrid closed-loop therapy versus current care among young people with type 1 diabetes: a patient-level simulation.

References

- Nathan DM, Genuth S, Lachin J, et al.; Diabetes Control and Complications Trial Research Group. The effect of intensive treatment of diabetes on the development and progression of long-term complications in insulin-dependent diabetes mellitus. *N Engl J Med* 1993;329:977–986
- DCCT/EDIC Research Group. The Diabetes Control and Complications Trial/Epidemiology of Diabetes Interventions and Complications study at 30 years: overview. *Diabetes Care* 2014;37:9–16
- Nathan DM, Zinman B, Cleary PA, et al.; Diabetes Control and Complications Trial/Epidemiology of Diabetes Interventions and Complications (DCCT/EDIC) Research Group. Modern-day clinical course of type 1 diabetes mellitus after 30 years' duration: the diabetes control and complications trial/epidemiology of diabetes interventions and complications and Pittsburgh epidemiology of diabetes complications experience (1983–2005). *Arch Intern Med* 2009;169:1307–1316
- Brod M, Christensen T, Thomsen TL, Bushnell DM. The impact of non-severe hypoglycemic events on work productivity and diabetes management. *Value Health* 2011;14:665–671
- Aronson R, Galstyan G, Goldfracht M, Al Sifri S, Elliott L, Khunti K. Direct and indirect health economic impact of hypoglycaemia in a global population of patients with insulin-treated diabetes. *Diabetes Res Clin Pract* 2018;138:35–43

- Davis RE, Morrissey M, Peters JR, Wittrup-Jensen K, Kennedy-Martin T, Currie CJ. Impact of hypoglycaemia on quality of life and productivity in type 1 and type 2 diabetes. *Curr Med Res Opin* 2005;21:1477–1483
- Foster NC, Beck RW, Miller KM, et al. State of type 1 diabetes management and outcomes from the T1D Exchange in 2016–2018. *Diabetes Technol Ther* 2019;21:66–72
- Phelan H, Clapin H, Bruns L, et al. The Australasian Diabetes Data Network: first national audit of children and adolescents with type 1 diabetes. *Med J Aust* 2017;206:121–125
- Lind M, Svensson AM, Kosiborod M, et al. Glycemic control and excess mortality in type 1 diabetes. *N Engl J Med* 2014;371:1972–1982
- Livingstone SJ, Levin D, Looker HC, et al.; Scottish Diabetes Research Network epidemiology group; Scottish Renal Registry. Estimated life expectancy in a Scottish cohort with type 1 diabetes, 2008–2010. *JAMA* 2015;313:37–44
- Morgan E, Cardwell CR, Black CJ, McCance DR, Patterson CC. Excess mortality in type 1 diabetes diagnosed in childhood and adolescence: a systematic review of population-based cohorts. *Acta Diabetol* 2015;52:801–807
- Rawshani A, Sattar N, Franzén S, et al. Excess mortality and cardiovascular disease in young adults with type 1 diabetes in relation to age at onset: a nationwide, register-based cohort study. *Lancet* 2018;392:477–486
- Pease A, Lo C, Earnest A, Kiriakova V, Liew D, Zoungas S. The efficacy of technology in type 1 diabetes: a systematic review, network meta-analysis, and narrative synthesis. *Diabetes Technol Ther* 2020;22:411–421
- Pease A, Lo C, Earnest A, Kiriakova V, Liew D, Zoungas S. Time in range for multiple technologies in type 1 diabetes: a systematic review and network meta-analysis. *Diabetes Care* 2020;43:1967–1975
- Bekiari E, Kitsios K, Thabit H, et al. Artificial pancreas treatment for outpatients with type 1 diabetes: systematic review and meta-analysis. *BMJ* 2018;361:k1310
- Weisman A, Bai JW, Cardinez M, Kramer CK, Perkins BA. Effect of artificial pancreas systems on glycaemic control in patients with type 1 diabetes: a systematic review and meta-analysis of outpatient randomised controlled trials. *Lancet Diabetes Endocrinol* 2017;5:501–512
- Boughton CK, Hovorka R. New closed-loop insulin systems. *Diabetologia* 2021;64:1007–1015
- Australian Diabetes Society. Living Evidence for Diabetes Consortium. Australian Evidence-Based Clinical Guidelines for Diabetes 2020. New Recommendations: Medical device technology for the management of type 1 diabetes. Melbourne, Australia. Accessed 22 March 2022. Available from <https://diabetessociety.com.au/living-guidelines.asp>
- Addala A, Auzanneau M, Miller K, et al. A decade of disparities in diabetes technology use and HbA1c in pediatric type 1 diabetes: a transatlantic comparison. *Diabetes Care* 2021;44:133–140
- Pease A, Szwarcbard N, Earnest A, Andrikopoulos S, Wischer N, Zoungas S. Glycaemia and utilisation of technology across the lifespan of adults with type 1 diabetes: results of the Australian National Diabetes Audit (ANDA). *Diabetes Res Clin Pract* 2021;171:108609

21. Pease AJ, Andrikopoulos S, Abraham MB, et al. Utilisation, access and recommendations regarding technologies for people living with type 1 diabetes: consensus statement of the ADS/ADEA/APEG/ADIPS Working Group. *Med J Aust* 2021;215:473–478
22. Colagiuri S, Brnabic A, Gomez M, Fitzgerald B, Buckley A, Colagiuri R. *DiabCoSt Australia Type 1: assessing the burden of type 1 diabetes in Australia*. Canberra, Diabetes Australia, 2009
23. Tao B, Pietropaolo M, Atkinson M, Schatz D, Taylor D. Estimating the cost of type 1 diabetes in the U.S.: a propensity score matching method. *PLoS One* 2010;5:e11501
24. American Diabetes Association. Economic costs of diabetes in the U.S. in 2007. *Diabetes Care* 2008;31:596–615
25. Tanenbaum ML, Hanes SJ, Miller KM, Naranjo D, Bensen R, Hood KK. Diabetes device use in adults with type 1 diabetes: barriers to uptake and potential intervention targets. *Diabetes Care* 2017;40:181–187
26. Pease A, Zomer E, Liew D, Lo C, Earnest A, Zoungas S. Cost-effectiveness of health technologies in adults with type 1 diabetes: a systematic review and narrative synthesis. *Syst Rev* 2020;9:171
27. Jendle J, Pöhlmann J, de Portu S, Smith-Palmer J, Roze S. Cost-effectiveness analysis of the MiniMed 670G hybrid closed-loop system versus continuous subcutaneous insulin infusion for treatment of type 1 diabetes. *Diabetes Technol Ther* 2019;21:110–118
28. Roze S, Buompiersiere MI, Ozdemir Z, de Portu S, Cohen O. Cost-effectiveness of a novel hybrid closed-loop system compared with continuous subcutaneous insulin infusion in people with type 1 diabetes in the UK. *J Med Econ* 2021;24:883–890
29. Pease A, Zomer E, Liew D, et al. Cost-effectiveness analysis of a hybrid closed-loop system versus multiple daily injections and capillary glucose testing for adults with type 1 diabetes. *Diabetes Technol Ther* 2020;22:812–821
30. Abraham MB, de Bock M, Smith GJ, et al.; Australian Juvenile Diabetes Research Fund Closed-Loop Research Group. Effect of a hybrid closed-loop system on glycemic and psychosocial outcomes in children and adolescents with type 1 diabetes: a randomized clinical trial. *JAMA Pediatr* 2021;175:1227–1235
31. Husereau D, Drummond M, Petrou S, et al.; CHEERS Task Force. Consolidated Health Economic Evaluation Reporting Standards (CHEERS) statement. *BMJ* 2013;346:f1049
32. Palmer AJ, Si L, Tew M, et al. Computer modeling of diabetes and its transparency: a report on the eighth Mount Hood Challenge. *Value Health* 2018;21:724–731
33. Vemer P, Corro Ramos I, van Voorn GAK, Al MJ, Feenstra TL. AdViSE: a validation-assessment tool of health-economic models for decision makers and model users. *PharmacoEconomics* 2016;34:349–361
34. The Department of Health. The Pharmaceutical Benefits Advisory Committee Guidelines. Section 3A.1 Overview and rationale of the economic evaluation. Australian Government Department of Health. 2016. Accessed 19 March 2022. Available from <https://pbac.pbs.gov.au/section-3a/3a-1-overview-and-rationale-of-economic-evaluation.html>
35. Early Treatment Diabetic Retinopathy Study Research Group. Grading diabetic retinopathy from stereoscopic color fundus photographs—an extension of the modified Airlie House classification. ETDRS report number 10. *Ophthalmology* 1991;98(Suppl.):786–806
36. Cheng Q, Lazzarini PA, Gibb M, et al. A cost-effectiveness analysis of optimal care for diabetic foot ulcers in Australia. *Int Wound J* 2017;14:616–628
37. Australian Institute of Health and Welfare. Deaths among people with diabetes in Australia, 2009–2014. Cat. no. CVD 79. Canberra: AIHW. 2017. Accessed 19 March 2022. Available from <https://www.aihw.gov.au/reports-data/health-conditions-disability-deaths/diabetes/overview>
38. Vistisen D, Andersen GS, Hansen CS, et al. Prediction of first cardiovascular disease event in type 1 diabetes mellitus: the Steno Type 1 Risk Engine. *Circulation* 2016;133:1058–1066
39. Cederholm J, Eeg-Olofsson K, Eliasson B, Zethelius B; Swedish National Diabetes Register. A new model for 5-year risk of cardiovascular disease in Type 1 diabetes; from the Swedish National Diabetes Register (NDR). *Diabet Med* 2011;28:1213–1220
40. Diabetes Control and Complications Trial (DCCT)/Epidemiology of Diabetes Interventions and Complications (EDIC) Study Research Group. Intensive diabetes treatment and cardiovascular outcomes in type 1 diabetes: the DCCT/EDIC study 30-year follow-up. *Diabetes Care* 2016;39:686–693
41. Australian Bureau of Statistics. 2021. Consumer Price Index, Australia, March 2021. Australian Bureau of Statistics. 2021. Accessed 19 March 2022. Available from <https://www.abs.gov.au/statistics/economy/price-indexes-and-inflation/consumer-price-index-australia/latest-release#data-download>
42. Parliament of Australia. 2014. Senate Standing Committee on Community Affairs: Out-of-pocket costs in Australian healthcare - Submissions received by the Committee (Submission 49 - Kidney Health Australia). Accessed 19 March 2022. Available from https://www.aph.gov.au/Parliamentary_Business/Committees/Senate/Community_Affairs/Australian_healthcare/Submissions
43. Cass A, Chadban S, Gallagher M, et al. *The economic impact of end stage kidney disease in Australia: projections to 2020*. Melbourne, Australia, Kidney Health Australia, 2010
44. Arora N, Makino K, Tilden D, Lobotesis K, Mitchell P, Gillespie J. Cost-effectiveness of mechanical thrombectomy for acute ischemic stroke: an Australian payer perspective. *J Med Econ* 2018;21:799–809
45. Gloede TD, Halbach SM, Thrift AG, Dewey HM, Pfaff H, Cadilhac DA. Long-term costs of stroke using 10-year longitudinal data from the North East Melbourne Stroke Incidence Study. *Stroke* 2014;45:3389–3394
46. National Heart Foundation of Australia. 2018. Economic Cost of Acute Coronary Syndrome in Australia: The Cost to Governments. Accessed 19 March 2022. Available from https://www.heartfoundation.org.au/getmedia/8ba34ebc-e4b7-44d7-a5a6-15feac1b86d8/Economic_cost_of_ACS_in_Australia_the_cost_to_governments.pdf
47. Australian Institute of Health and Welfare. Deaths in Australian hospitals 2014–15. Cat. no. WEB 186. Canberra: AIHW. 2017. Accessed 8 April 2022. Available from <https://www.aihw.gov.au/reports/hospitals/deaths-in-australian-hospitals-2014-15/contents/summary>
48. Jiang J, May P. Proportion of deaths in hospital in European countries: trends and associations from panel data (2005–2017). *Eur J Public Health* 2021;31:1176–1183
49. Ly TT, Brnabic AJ, Eggleston A, et al. A cost-effectiveness analysis of sensor-augmented insulin pump therapy and automated insulin suspension versus standard pump therapy for hypoglycemic unaware patients with type 1 diabetes. *Value Health* 2014;17:561–569
50. Kawasaki R, Akune Y, Hiratsuka Y, Fukuhara S, Yamada M. Cost-utility analysis of screening for diabetic retinopathy in Japan: a probabilistic Markov modeling study. *Ophthalmic Epidemiol* 2015;22:4–12
51. Szabo SM, Beusterien KM, Pleil AM, et al. Patient preferences for diabetic retinopathy health states. *Invest Ophthalmol Vis Sci* 2010;51:3387–3394
52. Hiratsuka Y, Yamada M, Akune Y, et al.; Eye Care Comparative Effectiveness Research Team (ECCERT). Cost-utility analysis of cataract surgery in Japan: a probabilistic Markov modeling study. *Jpn J Ophthalmol* 2013;57:391–401
53. Bagust A, Beale S. Modelling EuroQol health-related utility values for diabetic complications from CODE-2 data. *Health Econ* 2005;14:217–230
54. Wasserfallen JB, Halabi G, Saudan P, et al. Quality of life on chronic dialysis: comparison between haemodialysis and peritoneal dialysis. *Nephrol Dial Transplant* 2004;19:1594–1599
55. Smith-Palmer J, Bae JP, Boye KS, Norrbacka K, Hunt B, Valentine WJ. Evaluating health-related quality of life in type 1 diabetes: a systematic literature review of utilities for adults with type 1 diabetes. *Clinicoecon Outcomes Res* 2016;8:559–571
56. Redekop WK, Stolk EA, Kok E, Lovas K, Kalo Z, Busschbach JVV. Diabetic foot ulcers and amputations: estimates of health utility for use in cost-effectiveness analyses of new treatments. *Diabetes Metab* 2004;30:549–556
57. Clarke P, Gray A, Holman R. Estimating utility values for health states of type 2 diabetic patients using the EQ-5D (UKPDS 62). *Med Decis Making* 2002;22:340–349
58. Harris S, Mamdani M, Galbo-Jørgensen CB, Bøgelund M, Gundgaard J, Groleau D. The effect of hypoglycemia on health-related quality of life: Canadian results from a multinational time trade-off survey. *Can J Diabetes* 2014;38:45–52
59. Geddes J, Schopman JE, Zammitt NN, Frier BM. Prevalence of impaired awareness of hypoglycaemia in adults with Type 1 diabetes. *Diabet Med* 2008;25:501–504
60. Ratzki-Leewing A, Harris SB, Mequanint S, et al. Real-world crude incidence of hypoglycemia in adults with diabetes: results of the InHypo-DM Study, Canada. *BMJ Open Diabetes Res Care* 2018;6:e000503
61. Gold AE, MacLeod KM, Frier BM. Frequency of severe hypoglycemia in patients with type 1 diabetes with impaired awareness of hypoglycemia. *Diabetes Care* 1994;17:697–703
62. Pedersen-Bjergaard U, Pramming S, Heller SR, et al. Severe hypoglycaemia in 1076 adult patients with type 1 diabetes: influence of risk markers and selection. *Diabetes Metab Res Rev* 2004;20:479–486
63. ter Braak EW, Appelman AM, van de Laak M, Stolk RP, van Haeften TW, Erkelens DW. Clinical characteristics of type 1 diabetic patients with

- and without severe hypoglycemia. *Diabetes Care* 2000;23:1467–1471
64. Haynes A, Hermann JM, Miller KM, et al.; T1D Exchange, WACDD and DPV registries. Severe hypoglycemia rates are not associated with HbA1c: a cross-sectional analysis of 3 contemporary pediatric diabetes registry databases. *Pediatr Diabetes* 2017;18:643–650
65. Bulsara MK, Holman CD, Davis EA, Jones TW. The impact of a decade of changing treatment on rates of severe hypoglycemia in a population-based cohort of children with type 1 diabetes. *Diabetes Care* 2004;27:2293–2298
66. Abraham MB, Jones TW, Naranjo D, et al. ISPAD Clinical Practice Consensus Guidelines 2018: assessment and management of hypoglycemia in children and adolescents with diabetes. *Pediatr Diabetes* 2018;19(Suppl. 27):178–192
67. Gubitosi-Klug RA, Braffett BH, White NH, et al.; Diabetes Control and Complications Trial (DCCT)/Epidemiology of Diabetes Interventions and Complications (EDIC) Research Group. Risk of severe hypoglycemia in type 1 diabetes over 30 years of follow-up in the DCCT/EDIC study. *Diabetes Care* 2017;40:1010–1016
68. Pease A, Earnest A, Ranasinha S, et al. Burden of cardiovascular risk factors and disease among patients with type 1 diabetes: results of the Australian National Diabetes Audit (ANDA). *Cardiovasc Diabetol* 2018;17:77
69. Davis EA, Keating B, Byrne GC, Russell M, Jones TW. Hypoglycemia: incidence and clinical predictors in a large population-based sample of children and adolescents with IDDM. *Diabetes Care* 1997;20:22–25
70. Wang S, Gum D, Merlin T. Comparing the ICERs in medicine reimbursement submissions to NICE and PBAC—does the presence of an explicit threshold affect the ICER proposed? *Value Health* 2018;21:938–943
71. Taylor C, Jan S. Economic evaluation of medicines. *Aust Prescr* 2017;40:76–78
72. Senate Standing Committees on Community Affairs. Parliament of Australia. Chapter 3: Affordable access – assessing cost effectiveness. In Report: Availability of new, innovative and specialist cancer drugs in Australia. Commonwealth of Australia 2015. Accessed 28 March 2022. Available from https://www.aph.gov.au/Parliamentary_Business/Committees/Senate/Community_Affairs/Cancer_Drugs/Report
73. Pickup JC, Sutton AJ. Severe hypoglycaemia and glycaemic control in Type 1 diabetes: meta-analysis of multiple daily insulin injections compared with continuous subcutaneous insulin infusion. *Diabet Med* 2008;25:765–774
74. Khunti K, Alsifri S, Aronson R, et al.; HAT Investigator Group. Rates and predictors of hypoglycaemia in 27 585 people from 24 countries with insulin-treated type 1 and type 2 diabetes: the global HAT study. *Diabetes Obes Metab* 2016;18:907–915