

Expectation Maximisation (EM)

Erfan Loweimi

Speech Group, Machine Intelligence Lab, University of Cambridge

Outline ... EM


- Importance
- Goal
- Idea
- Derivation
- Visualisation

Importance ...

Maximum Likelihood from Incomplete Data via the *EM* Algorithm

By A. P. DEMPSTER, N. M. LAIRD and D. B. RUBIN

Harvard University and Educational Testing Service




Dempster, A. P., Laird, N. M. & Rubin, D. B. (1977). *Maximum likelihood from incomplete data via the EM algorithm*. **Journal of the Royal Statistical Society**: Series B, 39, 1-38.

Importance ...

Maximum Likelihood from Incomplete Data via the *EM* Algorithm

By A. P. DEMPSTER, N. M. LAIRD and D. B. RUBIN

Harvard University and Educational Testing Service



<https://www.jstor.org> › stable

Maximum Likelihood from Incomplete Data via the ... - JSTOR

by AP Dempster · 1977 · Cited by 69463 — A broadly applicable algorithm for computing maximum likelihood estimates from incomplete data is presented at various levels of generality.

21, Feb, 2023


Importance ...

Keywords ...

Maximum Likelihood from Incomplete Data via the *EM* Algorithm

By A. P. DEMPSTER, N. M. LAIRD and D. B. RUBIN

Harvard University and Educational Testing Service



<https://www.jstor.org> › stable

Maximum Likelihood from Incomplete Data via the ... - JSTOR

by AP Dempster · 1977 · Cited by 69463 — A broadly applicable algorithm for computing maximum likelihood estimates from incomplete data is presented at various levels of generality.

21, Feb, 2023

Setup

X : *observable* rv^{*}

$$\mathbf{X} = \{\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_N\}, \quad \mathbf{x}_i \in \mathbb{R}^{D_1}$$

Z : *latent* rv

$$\mathbf{Z} = \{\mathbf{z}_1, \mathbf{z}_2, \dots, \mathbf{z}_N\}, \quad \mathbf{z}_i \in \mathbb{R}^{D_2}$$

\mathbf{X} : *incomplete* data

$\{\mathbf{X}, \mathbf{Z}\}$: *complete* data

θ : model parameters

* rv: random variable

Setup

independent

identically distributed

$$p(\mathbf{X}|\theta) = p(\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_N|\theta) \stackrel{i.i.d}{=} \prod_{i=1}^N p(\mathbf{x}_i|\theta)$$

$$\log p(\mathbf{X}|\theta) = \sum_{i=1}^N \log p(\mathbf{x}_i|\theta)$$

$$p(\mathbf{X}, \mathbf{Z}|\theta) = p(\mathbf{x}_1, \mathbf{z}_1, \mathbf{x}_2, \mathbf{z}_2, \dots, \mathbf{x}_N, \mathbf{z}_N|\theta) \stackrel{i.i.d}{=} \prod_{i=1}^N p(\mathbf{x}_i, \mathbf{z}_i|\theta)$$

$$\log p(\mathbf{X}|\theta) = \log \sum_{\mathbf{Z}} p(\mathbf{X}, \mathbf{Z}|\theta)$$

Setup

$P(\mathbf{X}|\theta)$: *incomplete data* likelihood

$P(\mathbf{X}, \mathbf{Z}|\theta)$: *complete data* likelihood

$P(\mathbf{Z}|\mathbf{X}, \theta)$: *posterior* probability

Marginalisation

$$p(\mathbf{X}|\theta) = \sum_{\mathbf{Z}} p(\mathbf{X}, \mathbf{Z}|\theta)$$

Chain rule (probability)

$$p(\mathbf{X}|\theta) = \frac{p(\mathbf{X}, \mathbf{Z}|\theta)}{p(\mathbf{Z}|\mathbf{X}, \theta)}$$

Goal ...

$$\theta_{ML}^* = \operatorname{argmax}_{\theta \in \Theta} p(\mathbf{X}|\theta) = \operatorname{argmax}_{\theta \in \Theta} \log p(\mathbf{X}|\theta)$$


- **log** is *strictly increasing* → argmax remains identical
- **Advantages:**
 - ✓ Mathematical convenience → $\log[\exp(.)]$
 - ✓ Numerical Stability

Goal ...

$$\begin{aligned}\theta_{ML}^*(\mathbf{X}|\theta) &= \operatorname{argmax}_{\theta \in \Theta} \log p(\mathbf{X}|\theta) \\ &= \operatorname{argmax}_{\theta \in \Theta} \log \sum_{\mathbf{Z}} p(\mathbf{X}, \mathbf{Z}|\theta)\end{aligned}$$

EM assumes model includes *latent variables* (\mathbf{Z}).


Latent Variable (Z)



Interpretation: *latent variable* is a part of a model ... *explains X*.

Latent Variable (Z)

$$\mathbf{x} \sim p(\mathbf{x})$$





Consider clustering ...

Interpretation: *latent variable* is a part of a model ... *explains X*.

Latent Variable (Z)

$$\mathbf{x} \sim p(\mathbf{x})$$



$$(\mathbf{x}, \mathbf{z}) \sim p(\mathbf{x}, \mathbf{z})$$

Interpretation: *latent variable* is a part of a model ... *explains* X.

Direct Solution

$$\theta_{ML}^* = \operatorname{argmax}_{\theta \in \Theta} \underbrace{\log \sum_{\mathbf{Z}} p(\mathbf{X}, \mathbf{Z} | \theta)}_{\log p(\mathbf{X} | \theta)}$$

$$\frac{\partial \log p(\mathbf{X} | \theta)}{\partial \theta} = 0$$

Step 1

Step 2

$$\left. \frac{\partial^2 \log p(\mathbf{X} | \theta)}{\partial \theta^2} \right|_{\theta=\theta_0} < 0$$

Step 3

θ_0 : derivative roots

Direct Solution

$$\theta_{ML}^* = \operatorname{argmax}_{\theta \in \Theta} \underbrace{\log \sum_{\mathbf{Z}} p(\mathbf{X}, \mathbf{Z} | \theta)}_{\log p(\mathbf{X} | \theta)}$$

Intractable (no closed-form solution)

$$\frac{\partial \log p(\mathbf{X} | \theta)}{\partial \theta} = 0$$

Step 1

Step 2

$$\left. \frac{\partial^2 \log p(\mathbf{X} | \theta)}{\partial \theta^2} \right|_{\theta = \theta_0} < 0$$

Step 3

θ_0 : derivative roots

Direction solution does not work!

$$\underbrace{\log \sum_z p(x, z|\theta)}_{\log p(x|\theta)} = \log(\dots + w_{z_i} e^{\frac{(x-\mu_i)^2}{2\sigma_i^2}} + \dots + w_{z_j} e^{\frac{(x-\mu_j)^2}{2\sigma_j^2}} + \dots)$$

Consider simplest case ... Gaussian

$$\frac{\partial \log p(x|\theta)}{\partial \theta} = 0$$

Intractable

Direction solution does not work!

$$\log \sum_z p(x, z|\theta) = \log(\dots + w_{z_i} e^{\frac{(x-\mu_i)^2}{2\sigma_i^2}} + \dots + w_{z_j} e^{\frac{(x-\mu_j)^2}{2\sigma_j^2}} + \dots)$$

... = 0 → Intractable

Direction solution does not work!

$$\log \sum_z p(x, z|\theta) = \log(\dots + w_{z_i} e^{\frac{(x-\mu_i)^2}{2\sigma_i^2}} + \dots + w_{z_j} e^{\frac{(x-\mu_j)^2}{2\sigma_j^2}} + \dots)$$

... = 0 → Intractable

If we could swap log & Σ ...

$$\sum_z \log p(x, z|\theta) = (\dots + w_i \log e^{\frac{(x-\mu_i)^2}{2\sigma_i^2}} + \dots + w_j \log e^{\frac{(x-\mu_j)^2}{2\sigma_j^2}} + \dots)$$

... = 0 → Tractable

$$= (\dots + c_i(x - \mu_i)^2 + \dots + c_j(x - \mu_j)^2 + \dots)$$

Optimise ... θ^*

$$\log \sum_{\mathbf{Z}} p(\mathbf{X}, \mathbf{Z} | \theta)$$

Intractable



Optimise ... θ^*

$$\sum_{\mathbf{Z}} \log p(\mathbf{X}, \mathbf{Z} | \theta)$$

Tractable

Optimise ... θ^*

$$\log \sum_{\mathbf{Z}} p(\mathbf{X}, \mathbf{Z} | \theta)$$



Optimise ... θ^*

$$\sum_{\mathbf{Z}} \log p(\mathbf{X}, \mathbf{Z} | \theta)$$

Intractable

Tractable

*Your problem is to bridge the gap which exists between
where you are now and the goal you intend to reach.*

*Earl Nightingale
(1921-1989)*

EM

Optimise ... θ^*

$$\log \sum_{\mathbf{Z}} p(\mathbf{X}, \mathbf{Z} | \theta)$$



Optimise ... θ^*

$$\sum_{\mathbf{Z}} \log p(\mathbf{X}, \mathbf{Z} | \theta)$$

Intractable

Tractable

*Your problem is to bridge the gap which exists between
where you are now and the goal you intend to reach.*

*Earl Nightingale
(1921-1989)*

EM Derivation – Step 0

$$p(\mathbf{X}|\theta) = \frac{p(\mathbf{X}, \mathbf{Z}|\theta)}{p(\mathbf{Z}|\mathbf{X}, \theta)}$$

Step 0: write $p(\mathbf{X}|\theta)$ using *chain rule*

EM Derivation – Step 1

$$\begin{aligned} p(\mathbf{X}|\theta) &= \frac{p(\mathbf{X}, \mathbf{Z}|\theta)}{p(\mathbf{Z}|\mathbf{X}, \theta)} \frac{q(\mathbf{Z})}{q(\mathbf{Z})} \\ &= \frac{p(\mathbf{X}, \mathbf{Z}|\theta)}{q(\mathbf{Z})} \frac{q(\mathbf{Z})}{p(\mathbf{Z}|\mathbf{X}, \theta)} \end{aligned}$$

Step 1: Multiply right-hand side in a *special 1*

EM Derivation – Step 2

$$\begin{aligned}\log p(\mathbf{X}|\theta) &= \log \left[\frac{p(\mathbf{X}, \mathbf{Z}|\theta)}{q(\mathbf{Z})} \frac{q(\mathbf{Z})}{p(\mathbf{Z}|\mathbf{X}, \theta)} \right] \\ &= \log \left[\frac{p(\mathbf{X}, \mathbf{Z}|\theta)}{q(\mathbf{Z})} \right] + \log \left[\frac{q(\mathbf{Z})}{p(\mathbf{Z}|\mathbf{X}, \theta)} \right]\end{aligned}$$

Step 2: Take *log* from both sides

EM Derivation – Step 3

$$q(\mathbf{Z}) \log p(\mathbf{X}|\theta) = q(\mathbf{Z}) \log \left[\frac{p(\mathbf{X}, \mathbf{Z}|\theta)}{q(\mathbf{Z})} \right] + q(\mathbf{Z}) \log \left[\frac{q(\mathbf{Z})}{p(\mathbf{Z}|\mathbf{X}, \theta)} \right]$$

Step 3: Multiply both sides by $q(\mathbf{Z})$

EM Derivation – Step 4

$$\sum_{\mathbf{Z}} q(\mathbf{Z}) \log p(\mathbf{X}|\theta) = \sum_{\mathbf{Z}} q(\mathbf{Z}) \log \left[\frac{p(\mathbf{X}, \mathbf{Z}|\theta)}{q(\mathbf{Z})} \right] + \sum_{\mathbf{Z}} q(\mathbf{Z}) \log \left[\frac{q(\mathbf{Z})}{p(\mathbf{Z}|\mathbf{X}, \theta)} \right]$$

Step 4: *Marginalise* over \mathbf{Z}

EM Derivation – Step 4

$$\sum_{\mathbf{Z}} q(\mathbf{Z}) \log p(\mathbf{X}|\theta) = \sum_{\mathbf{Z}} q(\mathbf{Z}) \log \left[\frac{p(\mathbf{X}, \mathbf{Z}|\theta)}{q(\mathbf{Z})} \right] + \sum_{\mathbf{Z}} q(\mathbf{Z}) \log \left[\frac{q(\mathbf{Z})}{p(\mathbf{Z}|\mathbf{X}, \theta)} \right]$$

$\log p(\mathbf{X}|\theta) = \dots$

Step 4: *Marginalise* over \mathbf{Z}

EM Derivation – Step 5

$$\log p(\mathbf{X}|\theta) = \sum_{\mathbf{Z}} q(\mathbf{Z}) \log \left[\frac{p(\mathbf{X}, \mathbf{Z}|\theta)}{q(\mathbf{Z})} \right] + \sum_{\mathbf{Z}} q(\mathbf{Z}) \log \left[\frac{q(\mathbf{Z})}{p(\mathbf{Z}|\mathbf{X}, \theta)} \right]$$

EM Derivation – Step 5.1

$$\log p(\mathbf{X}|\theta) = \sum_{\mathbf{Z}} q(\mathbf{Z}) \log \left[\frac{p(\mathbf{X}, \mathbf{Z}|\theta)}{q(\mathbf{Z})} \right] + \sum_{\mathbf{Z}} q(\mathbf{Z}) \log \left[\frac{q(\mathbf{Z})}{p(\mathbf{Z}|\mathbf{X}, \theta)} \right]$$

EM Derivation – Step 5.1

$$\log p(\mathbf{X}|\theta) = \sum_{\mathbf{Z}} q(\mathbf{Z}) \log \left[\frac{p(\mathbf{X}, \mathbf{Z}|\theta)}{q(\mathbf{Z})} \right] + \sum_{\mathbf{Z}} q(\mathbf{Z}) \log \left[\frac{q(\mathbf{Z})}{p(\mathbf{Z}|\mathbf{X}, \theta)} \right]$$

$$D_{KL}(q \parallel p) \triangleq \sum_y q(y) \log \frac{q(y)}{p(y)}$$


- D_{KL} (KL Divergence) properties:
 - ✓ $D_{KL}(q \parallel p) \geq 0$
 - ✓ $D_{KL}(q \parallel p) = 0 \iff q = p$

EM Derivation – Step 5.2

$$\log p(\mathbf{X}|\theta) = \sum_{\mathbf{Z}} q(\mathbf{Z}) \log \left[\frac{p(\mathbf{X}, \mathbf{Z}|\theta)}{q(\mathbf{Z})} \right] + \underbrace{\sum_{\mathbf{Z}} q(\mathbf{Z}) \log \left[\frac{q(\mathbf{Z})}{p(\mathbf{Z}|\mathbf{X}, \theta)} \right]}_{D_{KL}(q(\mathbf{Z}) \parallel p(\mathbf{Z}|\mathbf{X}, \theta))}$$



EM Derivation – Step 5.2

$$\log p(\mathbf{X}|\theta) = \text{whatever} + \underbrace{D_{KL}(q(\mathbf{Z}) || p(\mathbf{Z}|\mathbf{X}, \theta))}_{\geq 0}$$



EM Derivation – Step 5.2



$$\log p(\mathbf{X}|\theta) = \text{whatever} + \underbrace{D_{KL}(q(\mathbf{Z}) || p(\mathbf{Z}|\mathbf{X}, \theta))}_{\geq 0}$$



EM Derivation – Step 5.2

$$\log p(\mathbf{X}|\theta) = \text{whatever} + \underbrace{D_{KL}(q(\mathbf{Z}) || p(\mathbf{Z}|\mathbf{X}, \theta))}_{\geq 0}$$

$$\log p(\mathbf{X}|\theta) \geq \text{whatever}$$




EM Derivation – Step 5.2

$$\log p(\mathbf{X}|\theta) = \text{ELBO} + \underbrace{D_{KL}(q(\mathbf{Z}) || p(\mathbf{Z}|\mathbf{X}, \theta))}_{\geq 0}$$

$$\log p(\mathbf{X}|\theta) \geq \text{ELBO}$$

Evidence **L**ower **B**ound (ELBO)

Evidence \equiv log-likelihood



EM Derivation – Step 5

$$\underbrace{\log \sum_{\mathbf{Z}} p(\mathbf{X}, \mathbf{Z})}_{\log p(\mathbf{X}|\theta)} = \underbrace{\sum_{\mathbf{Z}} q(\mathbf{Z}) \log \left[\frac{p(\mathbf{X}, \mathbf{Z}|\theta)}{q(\mathbf{Z})} \right]}_{ELBO(q, \theta)} + \underbrace{\sum_{\mathbf{Z}} q(\mathbf{Z}) \log \left[\frac{q(\mathbf{Z})}{p(\mathbf{Z}|\mathbf{X}, \theta)} \right]}_{D_{KL}(q, \theta)}$$

EM Derivation – Step 6

$$\underbrace{\log \sum_{\mathbf{Z}} p(\mathbf{X}, \mathbf{Z})}_{\log p(\mathbf{X}|\theta)} = \underbrace{\sum_{\mathbf{Z}} q(\mathbf{Z}) \log \left[\frac{p(\mathbf{X}, \mathbf{Z}|\theta)}{q(\mathbf{Z})} \right]}_{ELBO(q, \theta)} + \underbrace{\sum_{\mathbf{Z}} q(\mathbf{Z}) \log \left[\frac{q(\mathbf{Z})}{p(\mathbf{Z}|\mathbf{X}, \theta)} \right]}_{D_{KL}(q, \theta)}$$

$$\theta_{ML}^* = \operatorname{argmax}_{\theta \in \Theta} \log \sum_{\mathbf{Z}} p(\mathbf{X}, \mathbf{Z}|\theta)$$

Intractable!

Direction solution does not work!

$$\log \sum_z p(x, z|\theta) = \log(\dots + w_{z_i} e^{\frac{(x-\mu_i)^2}{2\sigma_i^2}} + \dots + w_{z_j} e^{\frac{(x-\mu_j)^2}{2\sigma_j^2}} + \dots)$$

... = 0 → Intractable

... = 0 → Tractable

RECALL slide 9

IFF we could swap log & Σ ...

$$\begin{aligned} \sum_z \log p(x, z|\theta) &= (\dots + w_i \log e^{\frac{(x-\mu_i)^2}{2\sigma_i^2}} + \dots + w_j \log e^{\frac{(x-\mu_j)^2}{2\sigma_j^2}} + \dots) \\ &= (\dots + c_i(x - \mu_i)^2 + \dots + c_j(x - \mu_j)^2 + \dots) \end{aligned}$$

EM Derivation – Step 5

$$\underbrace{\log \sum_{\mathbf{Z}} p(\mathbf{X}, \mathbf{Z})}_{\log p(\mathbf{X}|\theta)} = \underbrace{\sum_{\mathbf{Z}} q(\mathbf{Z}) \log \left[\frac{p(\mathbf{X}, \mathbf{Z}|\theta)}{q(\mathbf{Z})} \right]}_{ELBO(q,\theta)} + \underbrace{\sum_{\mathbf{Z}} q(\mathbf{Z}) \log \left[\frac{q(\mathbf{Z})}{p(\mathbf{Z}|\mathbf{X}, \theta)} \right]}_{D_{KL}(q,\theta)}$$

Intractable!

$$\log \sum_{\mathbf{Z}} p(\mathbf{X}, \mathbf{Z}|\theta)$$

RECALL



EM

Tractable

$$\sum_{\mathbf{Z}} \log p(\mathbf{X}, \mathbf{Z}|\theta)$$

EM Derivation – Step 6




$$\underbrace{\log \sum_{\mathbf{Z}} p(\mathbf{X}, \mathbf{Z})}_{\log p(\mathbf{X}|\theta)} = \underbrace{\sum_{\mathbf{Z}} q(\mathbf{Z}) \log \left[\frac{p(\mathbf{X}, \mathbf{Z}|\theta)}{q(\mathbf{Z})} \right]}_{ELBO(q, \theta)} + \underbrace{\sum_{\mathbf{Z}} q(\mathbf{Z}) \log \left[\frac{q(\mathbf{Z})}{p(\mathbf{Z}|\mathbf{X}, \theta)} \right]}_{D_{KL}(q, \theta)}$$

$$\theta^* = \operatorname{argmax}_{\theta \in \Theta} ELBO$$

Tractable


EM → Instead of **log-likelihood** ... maximise **ELBO** ...

Best ELBO to optimise ...




Which ELBO is better?

Best ELBO to optimise ...



- Lower D_{KL} \rightarrow Better ELBO (closer to $\log p(\mathbf{X}|\theta)$)
- **IDEAL:** $D_{KL} = 0 \rightarrow$ Best ELBO = $\log p(\mathbf{X}|\theta)$


EM Procedure



E-Step

$$D_{KL}(q, \theta^{(n-1)}) = \sum_{\mathbf{Z}} q(\mathbf{Z}) \log \left[\frac{q(\mathbf{Z})}{p(\mathbf{Z}|\mathbf{X}, \theta^{(n-1)})} \right]$$

$$\text{Min } D_{KL} = 0 \Leftrightarrow q^{(n)}(\mathbf{Z}) = p(\mathbf{Z}|\mathbf{X}, \theta^{(n-1)})$$



M-Step

$$\begin{aligned} ELBO(q^{(n)}, \theta) &= \sum_{\mathbf{Z}} q^{(n)}(\mathbf{Z}) \log \left[\frac{p(\mathbf{X}, \mathbf{Z} | \theta)}{q^{(n)}(\mathbf{Z})} \right] \\ &= \underbrace{\sum_{\mathbf{Z}} q^{(n)}(\mathbf{Z}) \log p(\mathbf{X}, \mathbf{Z} | \theta)}_{\text{constant}} - \underbrace{\sum_{\mathbf{Z}} q^{(n)}(\mathbf{Z}) \log q^{(n)}(\mathbf{Z})}_{\text{constant}} \end{aligned}$$

M-Step

$$\begin{aligned} ELBO(q^{(n)}, \theta) &= \sum_{\mathbf{Z}} q^{(n)}(\mathbf{Z}) \log \left[\frac{p(\mathbf{X}, \mathbf{Z} | \theta)}{q^{(n)}(\mathbf{Z})} \right] \\ &= \underbrace{\sum_{\mathbf{Z}} q^{(n)}(\mathbf{Z}) \log p(\mathbf{X}, \mathbf{Z} | \theta)}_{\text{constant}} - \underbrace{\sum_{\mathbf{Z}} q^{(n)}(\mathbf{Z}) \log q^{(n)}(\mathbf{Z})}_{\text{constant}} \end{aligned}$$


$$\theta^* = \operatorname{argmax}_{\theta \in \Theta} ELBO(q^{(n)}, \theta)$$

$$= \operatorname{argmax}_{\theta \in \Theta} \sum_{\mathbf{Z}} q^{(n)}(\mathbf{Z}) \log p(\mathbf{X}, \mathbf{Z} | \theta)$$

M-Step



M-step

$$\begin{aligned}\theta^* &= \operatorname{argmax}_{\theta \in \Theta} ELBO(q^{(n)}, \theta) \\ &= \operatorname{argmax}_{\theta \in \Theta} \sum_{\mathbf{Z}} q^{(n)}(\mathbf{Z}) \log p(\mathbf{X}, \mathbf{Z} | \theta)\end{aligned}$$





Visualisation

E-step: $q^{(1)} = \operatorname{argmin} D_{KL}(q, \theta^{(0)})$



Visualisation



M-step: $\theta^{(1)} = \operatorname{argmax} \text{ELBO}(q^{(1)}, \theta)$



Visualisation



E-step: $q^{(2)} = \operatorname{argmin} D_{KL}(q, \theta^{(1)})$

$$\log p(X|\theta^{(1)}) = \text{ELBO}(q^{(2)}, \theta^{(1)})$$





Visualisation

M-step: $\theta^{(2)} = \operatorname{argmax} \text{ELBO}(q^{(2)}, \theta)$




Visualisation

M-step: $\theta^{(N)} = \operatorname{argmax} \text{ELBO}(q^{(N)}, \theta)$




Initialisation Matters (1)




Initialisation Matters (2)

- $p(\mathbf{X}|\theta^{(n)})$ is **ALWAYS** non-decreasing
 - ✓ $ELBO^*(q^{(n+1)}, \theta^{(n+1)}) \geq ELBO^*(q^{(n)}, \theta^{(n)})$
 - ✓ Convergence is guaranteed ... but to local optimum ...



Initialisation Matters (3)


- $p(\mathbf{X}|\theta^{(n)})$ is **ALWAYS** non-decreasing
- Initialisation affects ... **convergence rate** and **final log-likelihood**



Initialisation Matters (3)

- $p(X|\theta^{(n)})$ is **ALWAYS** non-decreasing
- Initialisation affects ... convergence rate and final log-likelihood

Try **multiple** initialisations and pick up the **best** local optimum (**highest log-likelihood**).



Other Considerations

- EM is closely related to **Coordinate Ascent** [App B]
 - E-step: fix θ , optimise q
 - M-step: fix q , optimise θ
- EM shines when M-step can be solved analytically
- Alternatives when M-step is intractable ...
 - Generalised EM (**GEM**) \rightarrow (conjugate) gradient ascent in M-Step
 - Expectation Conditional Maximisation (**ECM**) \rightarrow coordinate ascent in M-step

Wrap-up ... EM ...

- **Goal:** estimate θ_{ML} for probabilistic models with latent var
- **How:** an iterative two-stage (E-step, M-step) procedure
- **Applications:** GMM, HMM, Computational biology, ...
- **Assignment:** estimate θ_{MAP} using EM
- **Appendices**
 - (A) Further Reading
 - (B) Coordinate Ascent

(A) Further Reading

- **Murphy**, Chapter 8, Section 7.2, Pages 306-310
- **Bishop**, Chapter 9, Section 4, Pages 450-455
- **Andrew Ng's** [Lecture Notes](#), Chapter 11, Pages 142-147
- Others: [blog1](#) [blog2](#)

(B) Coordinate Ascent

- **Iterative** optimisation method for **multi-variate** functions $f(\mathbf{x})$
- **Idea:** **Maximise** over one variable (or a block of variables) at a time, assuming others are constants

Procedure:

INITIALISE $\mathbf{x}^{(0)} = [x_0, x_1, \dots, x_{D-1}]$

FOR i in range(**#iterations**):

FOR d in range(D):

$\mathbf{x}^{(i)}[d] \leftarrow \underset{x_d}{\operatorname{argmax}} f(\mathbf{x}^{(i)}[:d], x_d, \mathbf{x}^{(i-1)}[d+1:])$

constants

axis-aligned movements

