

DD704731

CI-05874929-4

DD704731

CISTI ICIST

CI-05874929-4

Document Delivery Service
in partnership with the Canadian Agriculture LibraryService de fourniture de Documents
en collaboration avec la Bibliothèque canadienne de l'agriculture**THIS IS NOT AN INVOICE / CECI N'EST PAS UNE FACTURE**ANTHONY ARTALE
MED LIB NATHAN CUMMINGS CTR (S-46)
MEMORIAL SLOAN KETTERING CANCER CTR
1275 YORK AVENUE
NEW YORK, NY 10021
UNITED STATESORDER NUMBER: CI-05874929-4
Account Number: DD704731
Delivery Mode: ARI
Delivery Address: arielsf.infotrieve.com/140.16
3.217.217
Submitted: 2005/11/23 10:33:29
Received: 2005/11/23 10:33:29
Printed: 2005/11/23 12:05:41

Direct	Periodical	OPENURLOPAC	UNITED STATES
--------	------------	-------------	---------------

Client Number: DDS36611/GANGI-DINO, RITA
Title: TRENDS IN BIOCHEMICAL SCIENCES.
DB Ref. No.: IRN10020949
ISSN: ISSN09680004
Vol./Issue: 17/4/APR
Date: 1992
Pages: 129-EOA
Article Title: A MODULE OF THE DNAJ HEAT SHOCK PROTEINS FOUND IN MALARIA
Article Author: BORK, J
Report Number: IRN10020949
Publisher: PUBLISHED FOR THE INTERNATIONAL UNION OF BIOCHEMISTRY BY ELSEVIER,**INSTRUCTIONS: PATRON REQUESTS COLOUR IF AVAILABLE. THANK YOU.****Estimated cost for this 1 page document: \$10.2 document supply fee +
\$30 copyright = \$40.2**

The attached document has been copied under license from Access Copyright/COPIBEC or other rights holders through direct agreements. Further reproduction, electronic storage or electronic transmission, even for internal purposes, is prohibited unless you are independently licensed to do so by the rights holder.

Phone/Téléphone: 1-800-668-1222 (Canada - U.S./E.-U.) (613) 998-8544 (International)
www.nrc.ca/cisti Fax/Télécopieur: (613) 993-7619 www.cnrc.ca/icist
info.cisti@nrc.ca info.icist@nrc.ca

TIBS 17 - APRIL 1992

PROTEIN SEQUENCE MOTIFS

A module of the DnaJ heat shock proteins found in malaria parasites

The DnaJ heat shock protein of *Escherichia coli* plays an essential role in the chaperone function of the hsp70-like DnaK protein¹. DnaJ stimulates the ATPase activity of DnaK² and also directly interacts with specific substrates of the DnaK chaperone machinery^{1,3}. Several proteins of *Saccharomyces cerevisiae* possess extensive sequence homologies to DnaJ⁴⁻⁷ and a human homologue has also been identified⁸. Sequence alignments suggest four areas of homology^{5,6}: a low homology region generally found at the carboxy-terminal end, a glycine-rich region, a domain containing four CXXCXGXG motifs and a more highly conserved region often found near the amino terminus. The yeast protein Sec63 contains only the region of highest homology and this is located away from the amino terminus, between two of its three putative membrane-spanning segments⁴ (Fig. 1a). Thus, the DnaJ family shows typical features of mosaic proteins⁹ which contain different building units (modules) with separate functions.

A database search with sequence consensus patterns¹⁰ of the most conserved (amino-terminal) domain of DnaJ revealed a surprising similarity to the ring-infected erythrocyte surface antigen (RESA)¹¹ of the malaria parasite *Plasmodium falciparum*. The sequence identity to DnaJ is as high as 39% over a length of 70 residues (Fig. 1b), only slightly below that of Sec63 (42%), higher than that of the human DnaJ-like protein (37%), and clearly above the threshold for structural homology of globular proteins¹². RESA contains two glutamate-rich regions, one next to the homologous segment and the other one at the carboxyl terminus (Fig. 1a). An equivalent acidic region is also found in Sec63 (Fig. 1a) confirming the modular architecture of both proteins.

RESA is an important non-polymorphic malaria antigen¹³ that has been shown to confer some degree of protective immunity on monkeys¹⁴. A fragment of RESA that includes the DnaJ region of homology was shown to be effective in providing partial immunity¹⁴. RESA is synthesized prior to *P. falciparum* merozoite differentiation and later becomes associated with the red-cell membrane skeleton of newly invaded erythrocytes where it binds to spectrin¹⁵. Membrane association in turn is one of




Figure 1

(a) Modular architecture of DnaJ-like proteins. Both the glycine-rich and the acidic regions differ in detail between the proteins, but nevertheless suggest similar functions. SCJ1 and Sec63 are involved in protein sorting^{4,7}, YDJ1 participates in chromosome segregation⁵ and SIS1 is required for nuclear migration during mitosis⁶. The occurrence of the highly conserved motif in all these proteins suggests a common function for this domain. (b) Multiple alignment of the proteins sharing the conserved motif. Residues that are conserved in all except one sequence are shown in bold type. The top consensus line indicates conserved features (capitals, conserved amino acids; lower cases, conserved properties such as h, hydrophobic; a, aromatic; t, turn-like or polar).

the common features known for most DnaJ-like proteins¹⁴⁻⁸. The homology to *E. coli* DnaJ suggests that RESA also may participate in a molecular complex similar to the DnaJ-DnaK-GrpE chaperone system¹ with the motif participating in protein-protein interactions.

As more members are added to the DnaJ family, the number of possible key residues of the motif involved in these interactions can be confined. In addition to several positions in which only aromatic or hydrophobic amino acids seem to be accepted (Fig. 1b) there is one conserved aspartate as well as an invariant tripeptide HPD (YPY in the malaria parasite).

References

- Georgopoulos, C., Ang, D., Liberek, K. and Zylicz, M. (1990) in *Stress Proteins in Biology and Medicine* (Morimoto, R. I., Tissieres, A. and Georgopoulos, C., eds), pp. 191-221, Cold Spring Harbor Laboratory Press
- Liberek, K. et al. (1991) *Proc. Natl Acad. Sci. USA* 88, 2874-2878
- Wickner, S. H. (1990) *Proc. Natl Acad. Sci. USA* 87, 2690-2694
- Sadler, I. et al. (1989) *J. Cell Biol.* 109, 2665-2675

- Caplan, A. J. and Douglas, M. G. (1991) *J. Cell Biol.* 114, 609-621
- Luke, M. M., Sutton, A. and Arndt, K. T. (1991) *J. Cell Biol.* 114, 623-638
- Blumberg, H. and Silver, P. A. (1991) *Nature* 349, 627-630
- Raabe, T. and Manley, J. L. (1991) *Nucleic Acids Res.* 19, 6645
- Doolittle, R. F. (1985) *Trends Biochem. Sci.* 10, 233-237
- Bork, P. and Grunwald, C. (1990) *Eur. J. Biochem.* 191, 347-358
- Favaloro, J. M. et al. (1986) *Nucleic Acids Res.* 14, 8265-8277
- Sander, C. and Schneider, R. (1991) *Proteins* 9, 56-68
- Coppel, R. L. et al. (1984) *Nature* 310, 789-792
- Collins, W. E. et al. (1986) *Nature* 323, 259-262
- Foley, M., Tilley, L., Sawyer, W. H. and Anders, R. F. (1991) *Mol. Biochem. Parasitol.* 46, 137-148

PEER BORK, CHRIS SANDER AND ALFONSO VALENCIA

EMBL, W-6900 Heidelberg, Germany.

BERND BUKAU

University of Heidelberg, Center of Molecular Biology, W-6900 Heidelberg, Germany.