

This material may be protected by copyright law (Title 17 US Code)

DD704731

CISTI ICIST

CI-05870482-8

Document Delivery Service
in partnership with the **Canadian Agriculture Library**Service de fourniture de Documents
en collaboration avec la **Bibliothèque canadienne de l'agriculture****THIS IS NOT AN INVOICE / CECI N'EST PAS UNE FACTURE**ANTHONY ARTALE
MED LIB NATHAN CUMMINGS CTR (S-46)
MEMORIAL SLOAN KETTERING CANCER CTR
1275 YORK AVENUE
NEW YORK, NY 10021
UNITED STATES

ORDER NUMBER:	CI-05870482-8
Account Number:	DD704731
Delivery Mode:	ARI
Delivery Address:	arielsf.infotrieve.com/140.16 3.217.217
Submitted:	2005/11/21 16:14:06
Received:	2005/11/21 16:14:06
Printed:	2005/11/22 09:36:34

Direct	Periodical	OPENURLOPAC	UNITED STATES
---------------	-------------------	--------------------	----------------------

Client Number:	DDS36481/GANGI-DINO, R
Title:	TRENDS IN BIOCHEMICAL SCIENCES.
DB Ref. No.:	IRN10020949
ISSN:	ISSN09680004
Vol./Issue:	19/5/MAY
Date:	1994
Pages:	199-200
Article Title:	THE MODULAR STRUCTURE OF NIFU PROTEINS
Article Author:	OUZOUNIS C
Report Number:	IRN10020949
Publisher:	PUBLISHED FOR THE INTERNATIONAL UNION OF BIOCHEMISTRY BY ELSEVIER,

Estimated cost for this 2 page document: \$10.2 document supply fee + \$30 copyright = \$40.2

The attached document has been copied under license from Access Copyright/COPIBEC or other rights holders through direct agreements. Further reproduction, electronic storage or electronic transmission, even for internal purposes, is prohibited unless you are independently licensed to do so by the rights holder.

Phone/Téléphone: 1-800-668-1222 (Canada - U.S./E.-U.) (613) 998-8544 (International)
www.nrc.ca/cisti Fax/Télécopieur: (613) 993-7619 www.cnrc.ca/icist
info.cisti@nrc.ca info.icist@nrc.ca

National Research
Council CanadaConseil national
de recherches Canada

histidine, stabilizing the motif into a single folded domain^{12,14}. As well as the conservation of the Zn²⁺ ligands, there are additional conserved residues between MDM2 and the rest of the RING-finger family, including a number of hydrophobic residues implicated in stabilizing the RING-finger fold (Fig. 1a; Ref. 14). Interestingly, two of the proteins shown in the alignment are implicated in the control of cell fate, namely IAP¹⁵ and DROC3H¹⁶. ICPO is involved in gene regulation¹⁴ whereas the functions of CELR05 (a *Caenorhabditis elegans* protein)¹⁷ and RING1 (Ref. 12) are not known. MDM2 and CELR05 show significant sequence homologies outside the RING finger, and both can be aligned (17% identity over 491 residues; Fig. 1b). To further assess the significance of the alignment, the zinc finger region of MDM2 (434–491) was used to search the OWL database (version 22.1) with the program PROSRCH¹⁸ using PAM matrices of 100 and 250. Statistically significant matches (excluding itself) were observed only between MDM2 and IAP, DROC3H, CELR05 and ICPO. Furthermore, no matches were observed in the top 100 alignments between MDM2 and any C2H2 or C4 zinc-finger-containing protein, supporting the

proposal that MDM2 contains a RING finger rather than two tandemly arranged zinc fingers as previously postulated^{8–10}.

The extensive RING-finger family contains a number of known oncoproteins including, for example, MEL18, BMI-1, RFP, PML and T18 (reviewed in Ref. 13). A large number of proteins in this family are also implicated in the control of cell growth, differentiation and development¹³. The presence of the RING-finger domain in MDM2 may infer a common functionality with other members of the RING-finger family. This function may involve protein–protein or protein–nucleic acid interaction, which would be distinct from the proposed inactivation of p53, but necessary for the other important biological activities of MDM2.

References

- 1 Eliyahu, D. *et al.* (1989) *Proc. Natl Acad. Sci. USA* 86, 8763–8767
- 2 Finlay, C. A., Hinds, P. W. and Levine, A. J. (1989) *Cell* 57, 1083–1093
- 3 Kern, S. E. *et al.* (1991) *Science* 252, 1708–1711
- 4 Zambetti, G. P. *et al.* (1992) *Genes Dev.* 6, 1143–1152
- 5 Levine, A. J. (1990) *Virology* 177, 419–426
- 6 Momand, J. *et al.* (1992) *Cell* 69, 1237–1245
- 7 Fakhrazadeh, S. S., Trusko, S. P. and George, D. L. (1991) *EMBO J.* 10, 1565–1569
- 8 Oliner, J. D. *et al.* (1992) *Nature* 358, 80–83

- 9 Chen, J., Marechal, V. and Levine, A. J. (1993) *Mol. Cell. Biol.* 13, 4107–4114
- 10 Oliner, J. D. *et al.* (1993) *Nature* 362, 857–860
- 11 Freeman, P. S., Hanson, I. M. and Trowsdale, J. (1991) *Cell* 64, 483–484
- 12 Lovering, R. *et al.* (1993) *Proc. Natl Acad. Sci. USA* 90, 2112–2116
- 13 Freemont, P. S. (1993) *Anal. NY Acad. Sci.* 684, 174–192
- 14 Everett, R. D. *et al.* (1993) *J. Mol. Biol.* 234, 1038–1047
- 15 Crook, N. E., Clem, R. J. and Miller, K. L. (1993) *J. Virol.* 67, 2168–2174
- 16 Price, B. D. *et al.* (1993) *EMBO J.* 12, 2411–2418
- 17 Sulston, J. *et al.* (1992) *Nature* 356, 37–41
- 18 Collins, J. F., Coulson, A. F. W. and Lyall, A. (1988) *CABIOS* 4, 67–71
- 19 Barton, G. J. and Sternberg, M. J. E. (1990) *J. Mol. Biol.* 212, 389–402
- 20 Needleman, S. B. and Wunsch, C. D. (1970) *J. Mol. Biol.* 48, 443–453

MICHAEL N. BODDY AND PAUL S. FREEMONT

Protein Structure Laboratory, Imperial Cancer Research Fund, 44 Lincoln's Inn Fields, London, UK WC2A 3PX.

KATHERINE L. B. BORDEN

Laboratory of Molecular Structure, National Institute for Medical Research, The Ridgeway, Mill Hill, London, UK NW7 1AA.

The modular structure of NifU proteins

The *nif* cluster contains genes responsible for nitrogen fixation in several prokaryotes^{1–3}. With the exception of the essential genes coding for and regulating the expression of the nitrogenase subunits, the functions of other *nif* genes appear to be associated with cofactor biosynthesis, amino acid interconversion and regulation of ammonia storage. However, not all the biochemical activities of the *nif* gene products have been identified yet. Some of these proteins might represent specialized versions of general metabolic enzymes; thus their presence cannot be precluded from non-nitrogen fixing organisms. A recent example is the identification of the NifS family as pyridoxal-phosphate-dependent aminotransferases⁴.

Here we report the identification by sequence analysis of two distinct domains in the NifU protein family (Fig. 1a). Although the biochemical function of NifU proteins remains unknown, it has been shown that NifU exists as a homodimer, containing one redox 2Fe–2S cluster per subunit⁵.

The carboxy-terminal domain, known to be the only common region of NifU between nitrogen-fixing bacteria and several rhodobacterial species^{6,7}, is shown here to be present in: the open reading

frame (ORF) YKL253 of chromosome XI of yeast⁸; a hypothetical 22 kDa protein from *Haemophilus influenzae*⁹; and a short ORF (ORF2) in *Azotobacter vinelandii*¹⁰ (Fig. 1b). Thus, *A. vinelandii* has two NifU-like proteins, one of which contains only the carboxy-terminal domain (Fig. 1a, b).

An internal domain of NifU proteins from nitrogen-fixing bacteria, not present in the so-called NifU proteins from two *Rhodobacter* species or the NifU-like protein from yeast, shows significant sequence similarity (Fig. 1c) to two internal repeats of nitrite reductases from *Klebsiella pneumoniae* (NasB)¹¹, *Escherichia coli* (NirB)¹², *Emericella nidulans* (NiiA)¹³ and *Neurospora crassa* (Nit-6)¹⁴, as well as to a single carboxy-terminal domain of nitrate reductase of *K. pneumoniae* (NasA)¹¹, the carboxyl terminus of NifE from *Bradyrhizobium japonicum*¹⁵ and a short ORF (gp64)¹⁶ from *E. coli*. Interestingly, the domain is present in NifE of *B. japonicum* at its extreme carboxy-terminal end and is absent from the other NifE homologues. In *E. nidulans*, the domain is encoded by the fifth exon of the *niiA* (nitrite reductase) gene¹³. This domain contains two pairs of conserved cysteines that might participate in the formation of the 2Fe–2S cluster in NifU and the other proteins identified (Fig. 1a). This cluster might serve as an electron carrier in the nitrate and nitrite reductases.

Modular proteins, frequently found in higher eukaryotes, are rare in prokaryotes, probably because of a lack of appropriate genetic shuffling mechanisms (see Refs 17, 18 and references therein). The mobility of the NifU-like domains in both prokaryotes and lower eukaryotes therefore comes as a surprise. It will be a challenge to identify the genetic mechanism(s) that enabled the various species to acquire the two domains. In addition, knowledge of the modular structure of NifU-like proteins might contribute towards elucidating the functions of their domains.

References

- 1 Beynon, J. *et al.* (1987) *J. Bacteriol.* 169, 4024–4029
- 2 Arnold, W. *et al.* (1988) *J. Mol. Biol.* 203, 715–738
- 3 Jacobson, M. R. *et al.* (1989) *J. Bacteriol.* 171, 1017–1027
- 4 Ouzounis, C. and Sander, C. (1993) *FEBS Lett.* 322, 159–164
- 5 Dean, D. R. *et al.* (1993) *J. Bacteriol.* 175, 6737–6744
- 6 Meijer, W. G. and Tabita, F. R. (1992) *J. Bacteriol.* 174, 3855–3866
- 7 Masepohl, B. *et al.* (1993) *Mol. Gen. Genet.* 238, 369–382
- 8 Purnelle, B., Skala, J., Van Dyck, L. and Goffeau, A. (1992) *Yeast* 8, 977–986
- 9 Tomb, J. F., El-Hajj, H. and Smith, H. O. (1991) *Gene* 104, 1–10
- 10 Joerger, R. D., Jacobson, M. R. and Bishop, P. E. (1989) *J. Bacteriol.* 171, 3258–3267
- 11 Lin, J. T., Goldman, B. S. and Stewart, V. (1993) *J. Bacteriol.* 175, 2370–2378




Figure 1

(a) Modular architecture of proteins that contain NifU-like domains. For large proteins (fragments only shown), the length of sequence shown is indicated.

(b) Alignment of the approximately 65-residue carboxy-terminal domain. Proteins are indicated by their Swissprot names if available [these contain an underscore (_) in the entry code]. Nfu1/Rhoca and Nfu2/Rhoca, *R. capsulatus* Nfu1 and Nfu2; Nifu_Rhosh, *R. sphaeroides* Nifu; Orf2/Azovi, *A. vinelandii* ORF 2; Nifu_Anas1, Nifu_Anasp, Nifu/Plebo, Nifu_Azoch, Nifu_Azovi and Nifu_Klepn, NifU proteins from *Anabaena* (two species), *Plectonome boryanum*, *A. chroococcum*, *A. vinelandii* and *K. pneumoniae*, respectively; Yk53_Yeast, yeast NifU-like protein; Yorg_Haein, *H. influenzae* hypothetical 22 kDa protein encoded by ORF G. The position of the domain within the corresponding proteins is indicated, as is the EMBL database accession number. Residues conserved in more than half of the sequences are boxed and displayed as consensus (bottom line). The newly identified domains show significant sequence similarity to known NifU proteins (amino acid identity: Yk53_Yeast, 28-36%; Yorg_Haein, 24-33%; Orf2/Azovi 35-50%). Database searches with the candidate proteins result in BLASTP P-values¹⁹ that are below 10×10^{-6} for at least one NifU. In order to verify these similarities, various sequence analysis methods were used including profile and pattern searches²⁰.

(c) Alignment of the approximately 60-residue middle domain. Nirb_Ecoli, *E. coli* nitrite reductase B; Nasb/Klepn, *K. pneumoniae* nitrite reductase B; Nir_Emeni, *E. nidulans* nitrite reductase; Nit6/Neurc, *N. crassa* nitrite reductase; Nasa/Klepn, *K. pneumoniae* nitrite reductase A; Nife_Braja, *B. japonicum* Nife protein; Yhea_Ecoli, *E. coli* hypothetical protein encoded by the *bfr* 3'-region. The newly identified domains show significant sequence similarity to known NifU proteins as follows: Nasa/Klepn, 19-30%; Nirb_Ecoli domain 1 (as an example of nitrite reductases from prokaryotes and fungi), 21-38%; Yhea_Ecoli, 19-26%; Nife_Braja, 22-38%. The similarities were verified as in (b).

- 12 Peakman, T. et al. (1990) *Eur. J. Biochem.* 191, 315-323
- 13 Johnstone, I. L. et al. (1990) *Gene* 90, 181-192
- 14 Exley, G. E., Colandene, J. D. and Garrett, R. H. (1993) *J. Bacteriol.* 175, 2379-2392
- 15 Aguilar, O. M. et al. (1990) *Mol. Gen. Genet.* 224, 413-420

- 16 Andrews, S. C., Harrison, P. M. and Guest, J. R. (1989) *J. Bacteriol.* 171, 3940-3947
- 17 Bork, P. (1993) *Proteins* 17, 363-374
- 18 Doolittle, R. F. and Bork, P. (1993) *Sci. Am.* 269(4), 50-56
- 19 Altschul, S. F. et al. (1990) *J. Mol. Biol.* 215, 403-410

- 20 Bork, P. et al. (1992) *Protein Sci.* 1, 1677-1690

CHRISTOS OUZOUNIS, PEER BORK AND CHRIS SANDER

European Molecular Biology Laboratory, Meyerhofstrasse 1, D-69012 Heidelberg, Germany.