

Assignment 3

Satya Sangram Mishra

Download all python codes from

<https://github.com/satyasm45/Summer-Internship/tree/main/Assignment-3/Codes>

and latex-tikz codes from

<https://github.com/satyasm45/Summer-Internship/tree/main/Assignment-3>

1 QUESTION No. 2.55

Let **A** and **B** be the centres of two circles of equal radii 3 such that each one of them passes through the centre of the other. Let them intersect at **C** and **D**. Is $AB \perp CD$?

2 SOLUTION

The centers and radii of the two circles without any loss of generality are given in table 2.1

	Circle 1	Circle 2
Centre	$\mathbf{A} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}$	$\mathbf{B} = \begin{pmatrix} 3 \\ 0 \end{pmatrix}$
Radius	$r_1 = 3$	$r_2 = 3$

TABLE 2.1: Input values

The choice for **A** and **B** is valid as:

$$\|\mathbf{B} - \mathbf{A}\| = \|\mathbf{A} - \mathbf{B}\| = \|\mathbf{B}\| = 3 \quad (\because \mathbf{A} = 0) \quad (2.0.1)$$

Let us define:

$$\mathbf{u} = \begin{pmatrix} \cos \theta \\ \sin \theta \end{pmatrix}, \theta \in [0, 2\pi]. \quad (2.0.2)$$

Then any point $\begin{pmatrix} x_1 \\ y_1 \end{pmatrix}$ on Circle 1 is given by :

$$\begin{pmatrix} x_1 \\ y_1 \end{pmatrix} = \mathbf{A} + r_1 \mathbf{u} = 3 \begin{pmatrix} \cos \theta \\ \sin \theta \end{pmatrix} \quad (\because \mathbf{A} = 0) \quad (2.0.3)$$

This is the locus of Circle 1.

Similarly, locus of Circle 2 is given by:

$$\begin{pmatrix} x_2 \\ y_2 \end{pmatrix} = \mathbf{B} + r_2 \mathbf{u} = \begin{pmatrix} 3 \\ 0 \end{pmatrix} + 3 \begin{pmatrix} \cos \theta \\ \sin \theta \end{pmatrix} \quad (2.0.4)$$

Using the locus of the circles Fig. 2.1 was plotted.




Fig. 2.1: Circles with their points of intersection

We have **C** and **D** as points of intersection and $r_1 = r_2$. So,

$$\|\mathbf{C} - \mathbf{A}\|^2 = \|\mathbf{C} - \mathbf{B}\|^2 \quad (2.0.5)$$

Also, $\|\mathbf{P}\|^2 = \mathbf{P}^T \mathbf{P}$. So,

$$(\mathbf{C} - \mathbf{A})^T (\mathbf{C} - \mathbf{A}) = (\mathbf{C} - \mathbf{B})^T (\mathbf{C} - \mathbf{B}) \quad (2.0.6)$$

$$\Rightarrow \mathbf{A}^T \mathbf{C} - \mathbf{B}^T \mathbf{C} = \mathbf{C}^T \mathbf{B} - \mathbf{C}^T \mathbf{A} + \|\mathbf{A}\|^2 - \|\mathbf{B}\|^2 \quad (2.0.7)$$

For any two vectors **P** and **Q** we have

$\mathbf{P}^T \mathbf{Q} = \mathbf{Q}^T \mathbf{P}$. So,

$$2 \times \mathbf{A}^T \mathbf{C} - 2 \times \mathbf{B}^T \mathbf{C} = \|\mathbf{A}\|^2 - \|\mathbf{B}\|^2 \quad (2.0.8)$$

Similarly, using:

$$\|\mathbf{D} - \mathbf{A}\|^2 = \|\mathbf{D} - \mathbf{B}\|^2 \quad (2.0.9)$$

We get:

$$2 \times \mathbf{A}^T \mathbf{D} - 2 \times \mathbf{B}^T \mathbf{D} = \|\mathbf{A}\|^2 - \|\mathbf{B}\|^2 \quad (2.0.10)$$

Subtracting equation 2.0.10 from equation 2.0.8:

$$2 \times (\mathbf{A}^T - \mathbf{B}^T)(\mathbf{C} - \mathbf{D}) = 0 \quad (2.0.11)$$

$$\Rightarrow (\mathbf{A} - \mathbf{B})^T (\mathbf{C} - \mathbf{D}) = 0 \quad (2.0.12)$$

$$\Rightarrow AB \perp CD \quad (2.0.13)$$