SANDIA REPORT SAND20XX-XXXX Printed November 2021



MANIPULATE-2020: An Interface Tool for Nuclear Data Processing

Patrick J. Griffin

Prepared by Sandia National Laboratories Albuquerque, New Mexico 87185 and Livermore, California 94550 Issued by Sandia National Laboratories, operated for the United States Department of Energy by National Technology & Engineering Solutions of Sandia, LLC.

NOTICE: This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government, nor any agency thereof, nor any of their employees, nor any of their contractors, subcontractors, or their employees, make any warranty, express or implied, or assume any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represent that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government, any agency thereof, or any of their contractors or subcontractors. The views and opinions expressed herein do not necessarily state or reflect those of the United States Government, any agency thereof, or any of their contractors.

Printed in the United States of America. This report has been reproduced directly from the best available copy.

Available to DOE and DOE contractors from

U.S. Department of Energy Office of Scientific and Technical Information P.O. Box 62 Oak Ridge, TN 37831

Telephone: (865) 576-8401 Facsimile: (865) 576-5728 E-Mail: reports@osti.gov

Online ordering: http://www.osti.gov/scitech

Available to the public from

U.S. Department of Commerce National Technical Information Service 5301 Shawnee Rd Alexandria, VA 22312

Telephone: (800) 553-6847 Facsimile: (703) 605-6900 E-Mail: orders@ntis.gov

Online order: https://classic.ntis.gov/help/order-methods/



ABSTRACT

This report provides basic background data on the Manipulate-2020 code. This code is used for processing and "manipulation" of nuclear data in support of radiation metrology applications. The code is made available on the open GitHub repository and is available to the general nuclear data community.

CONTENTS

| 1. | Purp | ose | 8 |
|----|-------|--|----|
| 2. | Code | e Release Documentation | 9 |
| | 2.1. | Software Name | 9 |
| | 2.2. | Software Version | 9 |
| | 2.3. | Author(s) | 9 |
| | 2.4. | Software Completion Date | 9 |
| | 2.5. | Brief Description | 9 |
| | 2.6. | Practical Application | 9 |
| | 2.7. | Method of Solution | |
| | 2.8. | Computer(s) and Operating System for Which Software is Written | 9 |
| | | 2.8.1. Programing Language | |
| | 2.9. | Compiler Version | 10 |
| | 2.10. | Software Limitations | 10 |
| | 2.11. | Related/Auxiliary/Co-Requisite Software | 10 |
| | 2.12. | Hardware Requirements | 10 |
| | 2.13. | References | 10 |
| | 2.14. | Funding Information | 11 |
| | 2.15. | Treaties/Agreements/IP Management Plans | 11 |
| | 2.16. | Sensitivities | 11 |
| | 2.17. | Commercialization/Distribution Plans | 11 |
| | 2.18. | Further Development | 11 |
| 3. | Insta | llation | 12 |
| 4. | Desc | ription of Structure of Input File | 13 |
| 5. | | Execution Instructions | |
| | | | |
| 6. | | ble Application Cases | |
| | 6.1. | NJOY/GROUPR Response Extraction | |
| | | 6.1.1. Description | |
| | | 6.1.2. Output | |
| | 6.2. | Weight and Combine Response Functions | |
| | | 6.2.1. Description | |
| | | 6.2.2. Output | |
| | 6.3. | NJOY/GROUPR Cross Section Extraction | |
| | | 6.3.1. Description | |
| | | 6.3.2. Output | |
| | 6.4. | NJOY/GROUPR Spectrum Extraction | |
| | | 6.4.1. Description | |
| | | 6.4.2. Output | |
| | 6.5. | Folding of Cross Section with Neutron Spectrum | |
| | | 6.5.1. Description | |
| | | 6.5.2. Output | |
| | 6.6. | Validate Positive Definite Attribute for the Cross Section Covariance Matrix | |
| | | 6.6.1. Description | |
| | | 6.6.2. Output | |
| | 6.7. | Cross Section Correlation Matrix Conversion for Plotting Interface | |
| | | 6.7.1. Description | 24 |

| | | 6.7.2. Output | 24 |
|--------------------------|---|---|----------------|
| | 6.8. | Enforce Positive Definite Attribute and Unit Normalization for the Neutron Spectrum | |
| | | Covariance Matrix | |
| | | 6.8.1. Description | 26 |
| | | 6.8.2. Output | |
| | 6.9. | Spectrum Correlation Matrix Conversion for Plotting Interface | |
| | | 6.9.1. Description | |
| | | 6.9.2. Output | |
| | 6.10. | Spectrum Uncertainty in Spectrum-Averaged Response | |
| | | 6.10.1. Description | |
| | | 6.10.2. Output | |
| | 6.11. | Cross Section Uncertainty in Spectrum-Averaged Response | |
| | | 6.11.1. Description | |
| | | 6.11.2. Output | |
| 7. | Conc | usion | 31 |
| Fig Fig Fig Fig | Kernure 3: ure 4: ure 5: ure 6: NJO | 75 As Neutron Ionizing Kerma Compared to the Neutron Total and Displacement as | 21 22 22 |
| _ | | Standard Deviation for Example Dosimetry Cross Sections | |
| | ure 8: | Standard Deviation for Benchmark Field Neutron Spectra | 2 |
| LIS | т О | TABLES | |
| | | . Control Flags for Manipulate-2020 Code | |
| | | . Summary Characteristic of Spectrum/Cross Section Folding | |
| | | . Eigenvalues for Representative Dosimetry Cross Sections | |
| | | . Summary of Spectrum Uncertainty Characteristic of Spectrum/Cross Section Folding. | |
| Tab | le 6-4 | . Summary of Response Uncertainty Characteristic of Spectrum/Cross Section Folding. | 30 |

This page left blank

ACRONYMS AND DEFINITIONS

| Abbreviation | Definition | | |
|--------------|--|--|--|
| DOE | Department of energy | | |
| ENDF | Evaluated Nuclear Data File | | |
| ESTSC | Energy Science and Technology Software Center | | |
| NEA | Nuclear Energy Agency | | |
| NTESS | Nuclear Technology and Engineering Solutions of Sandia | | |
| SNL | Sandia National Laboratories | | |
| WSL | Windows Subsystem for Linux | | |

1. PURPOSE

The purpose of this document is to provide the basic information on then Manipulate-2020 code that is required to make it available to the general user community.

2. CODE RELEASE DOCUMENTATION

This section addresses the various aspects of the code and its release that are required for release by Sandia, with authorization from NTESS, through Energy Science and Technology Software Center (ESTSC), DOE's central software repository.

2.1. Software Name

MANIPULATE-2020

2.2. Software Version

Version 21-1

2.3. Author(s)

Patrick J. Griffin

2.4. Software Completion Date

June 15, 2021

2.5. Brief Description

Manipulate-2020 is a nuclear data manipulation code that interfaces with files output by the SNL-NJOY-2016 code and supports radiation metrology applications and material damage studies. Example applications include: a) the generation of tabular multigroup damage response functions from the SNL-NJOY-2016/GROUPR module; b) the weighting and combination of the NJOY-produced isotopic response functions to make elemental and material-specific damage response functions; c) the generation of uncertainty in spectrum-averaged metrics using either the spectrum or response covariance matrix; d) generation of correlation matrix data compatible with 2D plots using commercial software packages such as SigmaPlot; and e) testing and correction of correlation matrices to ensure that they are positive semidefinite and that spectrum correlation matrices obey the unity normalization constraint.

All of the modifications provided here are being made available to the GitHub-based repository for the Manipulate-2020 code. As additional useful enhancements are incorporated into the baseline Manipulate-2020 code, the radiation effects community will be able to use this expanded functionality in this nuclear data manipulation.

2.6. Practical Application

2.7. Method of Solution

This software interfaces the information found in publicly available nuclear data evaluations, i.e., ENDF/B-VIII-0, with basic physics-related models supported by NJOY code processing for material damage, e.g., arc-dpa formulation found in NEA/NSC/DOC(2015)9. So, the "method of solution" is simple data manipulation according to defined physics relationships.

2.8. Computer(s) and Operating System for Which Software is Written

The code was tested on a PC using the Windows 10 operating system and using Windows Subsystem for Linux (WSL) with the Ubuntu 18.04 LTS Linux system. Since the code does not use

any specialized features, it should operate correctly under any version of Linux. It should operate under a normal Linux system or on a PC using the WSL interface and other versions of Linux.

2.8.1. Programing Language

2.9. Compiler Version

The code uses the GNU f77 Fortran compiler. Compile flags are set in a Makefile and use the "dec" feature compatibility option.

The code is invoked through the C-shell interface.

The C-shell interface calls a Perl script to run the code and to move the output files to various locations within the directory structure.

2.10. Software Limitations

There are no significant software limitations related to either data storage or computational time. The code is designed for a single processor and does not use any computational algorithms that would have merit in a parallel or multi-core/cpu implementation.

The code uses some dated legacy Fortran features that should be updated to a more modern Fortran syntax in the future.

The code has some hard-coded directory locations that, if changed, will need to be modified within the source code to match the user-selected implementation location. Some of these hard-coded locations are currently configured to reach into parallel implementations of the SNL-NJOY-2016 code and the SNL-LSL code.

2.11. Related/Auxiliary/Co-Requisite Software

The code is invoked in a script that has been tested using Perl 5, version 26.

2.12. Hardware Requirements

There are no unusual hardware requirements. The hardware should have the memory and processor speed needed to support use of the Windows 10 operating system.

2.13. References

The relevant references consist of the NJOY-2016 manual, the ENDF/B-VIII.0 cross section library, and the IRDFF-II dosimetry cross section library.

- [1] Trkov, A., Griffin, P.J., Simakov, S.P., Greenwood, L.R., Zolotarev, K.I., Capote, R., et al., "IRDFF-II: A New Neutron Metrology Library", Nuclear Data Sheets, Vol. 163, pp. 1-108, 2020.
- [2] "ENDF-6 Formats Manual: Data Formats and Procedures for the Evaluated Nuclear Data Files ENDF/B-VI, ENDF/B-VII and ENDF/B-VIII", edited by A. Trkov, M. Herman, D.A. Brown, report CSEWG Document ENDF-102, report BNL-203218-2018-INRE, February 1, 2018. Available at: https://www.nndc.bnl.gov/csewg/docs/endf-manual.pdf
- [3] D.A. Brown, M.B. Chadwick, R. Capote, et al., "ENDF/B-VIII.0: The 8th Major Release of the Nuclear Reaction Data Library with CIELO-project Cross Sections, New Standards and Thermal Scattering Data," Nuclear Data Sheets, Vol. 148, pp. 1-142, February 2018.

2.14. Funding Information

C7 (NNSA Nuclear Survivability Engineering Campaign), Project 188897 \$50,000, (2010-2020) Susan Norwood, NA-115 (202-586-4779), susan.norwood@nnsa.doe.gov

2.15. Treaties/Agreements/IP Management Plans

None.

2.16. Sensitivities

None. This code is unlimited release with no distribution limitations. The source code is made available on a public Git-based code repository.

2.17. Commercialization/Distribution Plans

There are no plans for commercialization of the software. The software is to be provided to the general radiation effects community in order to improve our basic physics understanding of neutron-induced material damage.

The software will be distributed via GitHub. The tentative link for the project is:

https://github.com/pigriff/Manipulate-2020

While pending approval for use of the open code repository, a version of the software is currently available at:

https://gitlab-ex.sandia.gov/bdhehr/Manipulate-2020

The specific license software is: BSD-3 since this provides the best permissive license for our intent.

I would like to release the code because it is of value to the general radiation effects community. Many parties are already making their own code patches to implement this data manipulation capability, e.g., the ASTM standards community is results from this code to support surrent standards. A general release like this is one way to focus the efforts of the international community and prevent duplicative efforts. This release will provide a framework where others can contribute their efforts to verify the software, provide validation evidence for various classes of radiation-induced damage, and expand the damage modeling.

2.18. Further Development

The data manipulation capabilities of this code are expected to be expanded to match the needs of the application community. Further development is expected and will be reflected in updates to the GitHub repository.

3. INSTALLATION

The following steps will install the code:

- 1. The first step in the installation is to clone the code from GitHub repository.
- 2. If the default directory structure is not used, then the user needs to change the hard-coded directory locations within the source code.
- 3. The Makefile should be used to recompile the code. This Makefile will place the executable in the proper location under the "bin" subdirectory.

4. DESCRIPTION OF STRUCTURE OF INPUT FILE

The input file consists of four input segments.

The first segment is a title card in a80 format.

The second segment is a series of control integers in 19I2 format. Table 4-1 provides a description of the input control flags.

The third segment is a set of input descriptors that will vary based upon the control flags.

The fourth segment is a termination line that has the format "9000000000000000". The "9" flags the end of the calculation.

Table 4-1. Control Flags for Manipulate-2020 Code

| | | Table 4-1. Control Flags for Manipulate-2020 Code | | | | |
|----------|----------------------------|--|--|--|--|--|
| Contro | Control Flags: format 19I2 | | | | | |
| | | | | | | |
| С | icon(1) | type of action | | | | |
| С | = 9 | end | | | | |
| С | = 1 | combine cross section elements - logic unclear, PJG 10/17/2012 | | | | |
| С | = 2 | prepare component plot file format output | | | | |
| С | = 3 | display a matxsr file from njoy | | | | |
| С | =-3 | same as 3 for xxx.plt 89 file | | | | |
| С | = 4 | multiple plot file option | | | | |
| С | = 5 | sum and plot components | | | | |
| С | =-5 | perform statistical ananlysis on a series of functions | | | | |
| С | = 6 | fold source and response | | | | |
| С | = 7 | weight and plot function | | | | |
| С | =-7 | reciprocal weight and plot function | | | | |
| С | = 8 | file difference | | | | |
| С | =-8 | covariance manipulation options | | | | |
| С | =10 | interpolate matxsr file | | | | |
| С | =11 | interpolate tabulated file | | | | |
| С | =12 | expand by interpolation a tabulated file | | | | |
| С | =13 | expand an analytic function in 640 group structure | | | | |
| C NEW | =14 | combine covariance files in LSL format (common energy structure) - | | | | |
| С | =15 | extract PKA recoil spectrum for given reaction/energy - NEW | | | | |
| C NEW | =16 | extract integrated PKA recoil spectrum for as a function of energy - | | | | |
| С | | | | | | |
| С | icon(2) | input energy grid/data format | | | | |
| С | = 0 | njoy sand-ii 620/640/725/770 point neutron cross section | | | | |
| С | | (energies in ev, response data to be multiplied | | | | |
| С | | by 10**-3 to yield kerma units of mev-mb) | | | | |
| С | = 1 | j. kelly sand-ii pun spectrum format | | | | |
| С | | (default quantity differential number spectrum) | | | | |
| С | | compute iselect material = 1 number fraction | | | | |
| С | | 2 energy fraction | | | | |
| | | | | | | |

| Control Flags: | rmat 1912 |
|----------------|---|
| Control Flags: | uniat 1912 |
| С | 3 differential number fraction |
| С | 4 differential energy fraction |
| С | 5 integral number |
| С | 2 same as 1 |
| С | 3 flash x-ray curr spectral input |
| С | 4 list directed (plot output) |
| С | 5 njoy sand-ii /770 point neutron pka recoil spectrum |
| С | (energies in ev, response data to be multiplied |
| С | by 10**-3 to kev) |
| С | 6 njoy matxsr interface file - arbitrary structure |
| С | 9 user defined - xxx.flux file |
| С | 10 640 group response structure = xxx.res |
| С | arbitrary response structure = xxx.res |
| С | |
| c icon(| display/output special option |
| С | 0 no action |
| С | 1 punch in damout format |
| С | 2 punch in tdam format |
| С | |
| c icon(| print energy order |
| С | 0 punch high to low |
| С | 1 punch low to high |
| С | |
| c icon(| sand structure |
| С | 0 640 groups - SAND-II |
| С | 1 770 groups - SAND-II extended |
| С | 2 89 groups - NuGET neutron |
| С | 3 48 groups - NuGET gamma |
| С | 5 175 group - Vitamin-J |
| С | 6 725 group - IAEA |
| С | |
| c icon(| inhibit renormalization |
| С | 0 normal |
| С | 1 response, inhibit renormalization in filein |
| С | |
| c icon(| NJOY interface location |
| С | 0 sync Projects-Linux/NJOY-2016 default |
| С | 1 sync Linux/NJOY-2016 selection |
| С | |
| c icon(| print inhibit |
| c = | default print |
| c = | inhibit print by this level |

| Control | Flags: forma | at 19l2 |
|---------|--------------|---|
| С | < 0 | enhance print (debug option) |
| С | | |
| С | icon(10) | label replacement |
| С | = 0 | no action |
| С | = 1 | read new legend information |
| С | | |
| С | icon(11) | special action |
| С | = 0 | not used |
| С | = 1 | on pka plot, divide entry 1 by entry 3 |
| С | | to produce average recoil energy - |
| С | | when $icon(2) = 5$ |
| С | = 2 | on icon(1)=7 combine, punch plt-format |
| С | | interface file on indicated dir:filename |
| С | | (tbd) |
| С | icon(12) | reverse order of data read - only partially implemented |
| С | = 0 | no effect |
| С | = 1 | reverse |
| С | icon(13) | attenuation function |
| С | = 1 | if equal to 1 and icon(1)=6, then |
| С | | use new function for exponential attenuation |
| С | | with coefficient alpha |
| С | = 2 | if equal to 2 and $icon(1)=6$, then |
| С | | use sensor self-shielding correction factor |
| С | icon(14) | new title |
| С | = 0 | use first card in input stream |
| С | = 1 | read-in new title |
| С | | |
| С | icon(15) | flip order of data for punch |
| С | = 0 | do nothing |
| С | + 1 | invert order |
| С | | |
| С | icon(16) | re-direct to look for pka files |
| С | = 0 | do nothing |
| С | = 1 | look in njoy/output-pka directory |
| С | = 2 | look in NJOY9-2012 for groupr/pka files |
| С | | |
| С | icon(17) | point interpolation for grprin option |
| С | = 0 | no, not used |
| С | = 1 | yes, interpolate |
| С | | |
| С | icon(18) | precision control for sum/difference |
| С | = 0 | no action |

| Control Flags: format 19I2 | | | | | |
|----------------------------|----------|---|--|--|--|
| С | = 1 | force 0.0 response for differences < 1.e-4 of maximum element | | | |
| С | | | | | |
| С | icon(19) | inversion of output energy order | | | |
| С | = 0 | no change | | | |
| С | = 1 | invert order | | | |
| С | = 6 | njoy matxsr interface file - arbitrary structure | | | |

5. CODE EXECUTION INSTRUCTIONS

After installation, the code should be executed from the ../Manipulate-2020/input directory by invoking the C-shell script "manipulate", followed by the name of the input file. For example, for the example input file "example_NJOY_groupr_convert" the user should type "./manipulate-2020 example_NJOY_groupr_convert" in the WSL window.

6. SAMPLE APPLICATION CASES

The example input files are found in the ../Manipulate-2020/inp subdirectory. Supporting files from SNL-NJOY-2016 calculations are found in the ../Manipulate-2020/use_case_support_files subdirectory. The support files, generally, need to appear in the associated NJOY-2016 file system subdirectory during the code execution, but they are included in the Manipulate-2020 subdirectory for archival purposes.

6.1. NJOY/GROUPR Response Extraction

Sample input file: example_NJOY_groupr_response_convert

6.1.1. Description

This "Use Case" extracts a response function from the GROUPR module output of the SNL-NJOY-2016 code. For this case to run, the input file:

must have been run. This file examines the ⁷⁵As nuclear data file from the ENDF/B-VIII.0 library [ENDF8a, ENDF8b], uses the NJOY/HEATR module to calculate the various kermas for the isotope, and used the NJOY/GROUPR module rebin these response functions into the SNL-SAND-IV 770-group energy structure. This energy structure is a superset of the commonly used SAND-II fine-group energy structure.

This case will produce two files within the ../Manipulate-2020/response directory. The files will reflect the naming found in the input file and uses the reaction identifiers, 301 and 444, as the file suffix. The 301 file corresponds to the total kerma. The 444 file corresponds to the damage energy, which, for the NJOY input file, in this case corresponds to the displacement kerma. Note, this calculation uses the SNL-NJOY-2016 code and not the original NJOY-2016 code. The damage energy in the original NJOY code does not generate the displacement kerma, but, rather, produces a sharp threshold Kinchin-Pease damage energy.

In addition to the response files, a summary of the processing, and any error conditions, will be reflected in a "*.out" file found in the ../Manipulate-2020/output subdirectory.

6.1.2. Output

The only output here is the presence of the tabulated response functions in the ../Manipulate-2020/response subdirectory. Figure 1 shows a plot of the ⁷⁵As damage metrics, i.e., the total neutron kerma and the displacement kerma. The neutron ionizing kerma represents the difference between the total kerma and the displacement kerma. Most of the deposited energy from the neutron interactions less than ~100 keV is from low energy recoil atoms and this deposited energy goes into the displacement kerma. At higher incident neutron energies, a higher energy recoil atom is produced and more of the energy goes into ionization.




Figure 1: 75 As Neutron Total Kerma and Displacement Kerma Limit

6.2. Weight and Combine Response Functions

Sample input file: example_NJOY_response_combination

6.2.1. Description

This "Use Case" illustrates the weighting and combination of response files. In this simple example, we subtract the displacement kerma from the total kerma in the previous example to produce a neutron ionizing kerma.

This case will produce one file within the ../Manipulate-2020/response directory. The files will reflect the naming found in the input file and uses the suffix "IK" to indicate that is is an ionizing kerma. The input file reflects the "combination" of two responses. The first component, the total kerma, has a weighting factor of 1. The second component, the displacement kerma, has a weighting factor of -1. The result is a simple subtraction of the responses.

As in the previous example, a summary of the processing, and any error conditions, will be reflected in a "*.out" file found in the ../Manipulate-2020/output subdirectory.

6.2.2. Output

The output here is the tabular presentation of the neutron ionizing kerma that is placed in the response subdirectory. Figure 2 shows this ionizing kerma and compares it to the total and displacement

.




Figure 2: 75 As Neutron Ionizing Kerma Compared to the Neutron Total and Displacement Kermas

6.3. NJOY/GROUPR Cross Section Extraction

Sample input file: example_NJOY_groupr_xsec_convert

6.3.1. Description

This "Use Case" extracts a cross section from the GROUPR module output of the SNL-NIOY-2016 code. For this case to run, you must have previously run three SNL-NJOY-2016 input files:

Co59g_IRDFF-II_640grp_xsec.inp

Ag109g_IRDFF-II_640grp_xsec.inp

S32p_IRDFF-II_640grp_xsec.inp

The first file generated for the 59 Co(n, γ) 60 Co reaction using the IRDFF-II dosimetry cross section [IRDFF]. In this case, the cross section is found in the MF=3/MT=102 entry. The second input file generated a cross section for 109 Ag(n, γ) 110m Ag reaction using the IRDFF-II dosimetry cross section library. In this case, the cross section is found in the MF=10/MT=102 entry since the residual nucleus is a meta-stable isotope and this represents an activation product and not a MF=3 cross section. The third input file generates a cross section for the ³²S(n,p)³²P reaction using the MF=3/MT=103 component found in the IRDFF-II library.

These input files used the NJOY/GROUPR module to rebin these response functions into the SNL-SAND-II 640-group energy structure. This case will produce three files within the ../Manipulate-2020/response directory. The files will reflect the naming found in the input file and uses the reaction identifiers, 102, as the file suffix. The 102 represents the MT=102 (n, y) reaction. The 103 represents the MT=103 (n,p) reaction.

In addition to the response files, a summary of the processing, and any error conditions, will be reflected in a "*.out" file found in the ../Manipulate-2020/output subdirectory.

6.3.2. Output

The output here is the tabular cross section that appears in the response subdirectory. Figure 3 shows the cross section for the three reactions addressed in this example.




Figure 3: Cross Section for Example NJOY/GROUPR Extraction

6.4. NJOY/GROUPR Spectrum Extraction

Sample input file: example_NJOY_groupr_spectrum_conversion

6.4.1. Description

This "Use Case" extracts neutron spectra from the GROUPR module output of the SNL-NJOY-2016 code. For this case to run, you must have previously run two SNL-NJOY-2016 input files:

Cf252_IRDFF-II_640grp_spectrum_validation.inp

U235_IRDFF-II_640grp_spectrum_validation.inp

These files convert the NJOY/GROUPR 640-group characterization of the neutron spectra as they are found in the IRDFF-II library.

There are four segments for the Manipulate-2020 input file. Portions one and two address the ²⁵²Cf(s.f.) spontaneous fission neutron spectrum. Portions three and four address the ²³⁵U(nth) thermal fission spectrum. While the first and third segment do the NJOY/GROUPR extraction, the second and fourth portion of the input stream processes the spectrum and outputs the spectrum in various spectral representations. The spectrum-averaged energy is also output for inspection.

This case will produce a summary of the processing with the spectral characterization, and any error conditions, in a "*.out" file found in the ../Manipulate-2020/output subdirectory.

6.4.2. Output

The output here is the tabulation of the spectrum in multiple formats. Figure 4 shows the ²⁵²Cf(s.f.) spontaneous fission neutron spectrum depicted as a differential number, differential energy, and lethargy representation. The only difference between the differential energy and lethargy representation is the use of a linear y-axis. In a lethargy representation, the use of a linear y-axis and a logarithmic energy axis means that equal areas under the curve correspond to equal probabilities for the neutron.

The 252 Cf(s.f.) and 235 U(n_{th}) fission neutron spectra have a very similar in shape. Figure 5 shows that, in a differential energy representation, the spectra can not be clearly differentiated. In the lethargy representation a small change is seen. This small change is consistent with the average neutron energy in these two spectral representations reported in the output: $\left\langle E_n^{^{252}Cf(s.f.)} \right\rangle = 2.123615$ MeV and $\left\langle E_n^{^{235}U(n_{th})} \right\rangle = 2.000431$ MeV.




Figure 4: Different Representations of the ²⁵²Cf(s.f.) Spontaneous Neutron Fission Spectrum




Figure 5: Comparison of the ²³⁵U(n_{th}) and ²⁵²Cf(s.f.) Neutron Spectra

6.5. Folding of Cross Section with Neutron Spectrum

Sample input file: example_response_fold

6.5.1. Description

This "Use Case" folds a response or cross section in with a neutron spectrum. This case uses cross sections and neutron spectra produced in the previous "use cases". The example has six segments of the calculation. The example case uses the 252 Cf(s.f.) spontaneous fission and the 235 U(n_{th}) thermal fission spectrum. These spectra are folded in with the 109 Ag(n, γ) 110m Ag, 59 Co(n, γ) 60 Co, and 32 S(n,p) 32 P cross sections.

This case will produce a summary of the processing with the spectral characterization, and any error conditions, in a "*.out" file found in the ../Manipulate-2020/output subdirectory.

6.5.2. Output

The output file reports spectrum-averaged cross section and the energy sensitivity regions. Table 6-1 shows a summary of the main characteristics for the two neutron spectra and three cross sections addresses in this example. These spectrum-averaged cross sections and energy response regions are similar to what is reported in ASTM E261, Standard Practice for Determining Neutron Fluence, Fluence Rate, and Spectra by Radioactivation Techniques [ASTM261], for the ³²S(n,p)³²P reactions. This information is also reported in the precision and bias section of ASTM E265 Standard Test Method for Measuring Reaction Rates and Fast-Neutron Fluences by Radioactivation of Sulfur-32 [ASTM265].

| | | _ | | | | |
|---|------------------------------------|----------|------------|---|----------|--|
| Cross Section | Neutron | | | Response Sensitivity Energy Limits (MeV) | | |
| | Spectrum (mb) | 5% | 50% | 95% | | |
| ⁵⁹ Co(n,γ) ⁶⁰ Co | ²⁵² Cf(s.f.) | 4.864277 | 0.04768145 | 0.8829955 | 3.446157 | |
| | ²³⁵ U(n _{th}) | 5.026129 | 0.04544459 | 0.8577079 | 3.286839 | |
| 109 A g/p\110m A g | ²⁵² Cf(s.f.) | 9.34751 | 0.07278118 | 0.7176283 | 2.615176 | |
| ¹⁰⁹ Ag(n,γ) ^{110m} Ag | ²³⁵ U(n _{th}) | 9.739893 | 0.07204609 | 0.7047761 | 2.556543 | |
| 32 C (n.n)32D | ²⁵² Cf(s.f.) | 73.99146 | 2.260848 | 4.024577 | 7.734848 | |
| ³² S(n,p) ³² P | ²³⁵ U(n _{th}) | 66.82598 | 2.235097 | 3.917196 | 7.298408 | |

Table 6-1. Summary Characteristic of Spectrum/Cross Section Folding

6.6. Validate Positive Definite Attribute for the Cross Section Covariance Matrix

Sample input file: example_cross_section_covariance_verification

6.6.1. Description

This "Use Case" verifies that the cross section covariance matrix is positive semi-definite, i.e., that it does not have negative eigenvalues. Since the Manipulate-2010 code routinely checks the input file for the required mathematical properties, the easiest way to verify this is to examine the eigenvalues and to visually inspect the correlation matrix.

This case depends having run the SNL-NJOY-2016 code to extract the lsl-format covariance files. This requires that you have previously run the following NJOY-2016 cases:

This case will produce a summary of the processing with the spectral characterization, and any error conditions, in a "*.out" file found in the ../Manipulate-2020/output subdirectory.

6.6.2. Output

The output file report provides the eigenvalues and eigenvectors for the covariance matrix. Table 6-2 summarizes the eigenvalues obtained for the three dosimetry reactions addressed in this example set.

Table 6-2. Eigenvalues for Representative Dosimetry Cross Sections

| Cross Section | Eigenvalues | | | | |
|---|---|--|--|--|------------------------------|
| ⁵⁹ Co(n,γ) ⁶⁰ Co | 3.188861 0.500000 | 0.2093912 0.7999997 | 0.6967719 0.9729697 | 0.8649985 0.9000000 | 0.9670087 0.8999999 |
| ¹⁰⁹ Ag(n,γ) ^{110m} Ag | 1.650150 0.9821049 0.9920523 | 0.3700236 0.9854667 1.147018 | 1.097767 0.9950126 0.8893262 | 0.9657941 0.9937788 0.9636558 | 0.9778988 0.9899527 |
| ³² S(n,p) ³² P | 13.95432 1.302666 0.9223984 0.1028905 0.2960616 0.2880561E-2 0.2472797E-2 0.3343157E-6 0.2651053E-3 0.2119063E-3 | 0.2558193E-2 0.3516820E-3 0.1056020E-3 | 6.911028 1.169561 0.5475745 0.2553883 0.7237343E-2 0.2106898E-2 0.9133368E-3 0.5826211E-4 0.2343758E-3 0.1313290E-3 | 0.2367189E-2 0.5724487E-3 0.2939752E-3 0.1228880E-3 | 0.4116856E-3 0.2508267E-3 |

6.7. Cross Section Correlation Matrix Conversion for Plotting Interface


Sample input file: example_cross_section_covariance_plot_interface

6.7.1. Description

This "Use Case" verifies that examines the cross section covariance matrix in more detail by breaking it into the energy-dependent standard deviations and the energy-dependent correlation matrix. While the covariance matrix is hard to evaluate, these two derivative quantities can be plotted and visually inspected for consistency with expectations.

6.7.2. Output

This calculation places into the output directory files that extract the energy-dependent standard deviations and the energy-dependent correlation matrix. These extracted components can be graphically depicted using commercial plotting codes. Figure 6 shows the energy-dependent correlation matrices for these reactions and compares them to the direct NJOY output for these reactions. Figure 7 shows the energy-dependent standard deviations for the three example dosimetry reactions, in the form of a percentage of the cross section.






Figure 6: Correlation Matrix for Example Dosimetry Cross Sections and a Comparison with the NJOY Output




Figure 7: Standard Deviation for Example Dosimetry Cross Sections

6.8. Enforce Positive Definite Attribute and Unit Normalization for the Neutron Spectrum Covariance Matrix

Sample input file: example_benchmark_spectra_covariance_verification

6.8.1. Description

This "Use Case" verifies that

6.8.2. Output

The output

6.9. Spectrum Correlation Matrix Conversion for Plotting Interface

Sample input file: example_benchmark_spectra_covariance_plot_interface

6.9.1. Description

This "Use Case" verifies that

6.9.2. Output

The output





Figure 8: Standard Deviation for Benchmark Field Neutron Spectra



6.10. Spectrum Uncertainty in Spectrum-Averaged Response

Sample input file: example_spct_unc_spectrum_averaged_response

6.10.1. Description

This "Use Case" provides the uncertainty component due to the knowledge of the neutron spectrum as propagated into the spectrum-averaged response. This calculation repeats the spectrum folding using the fine 640-group representation of the response function and neutron spectrum before it does the uncertainty propagation. That initial step is performed so that the user can inspect the variation between the fine-grouped spectrum-averaged response and the spectrum-averaged response that is computed using the energy grid used for the characterization of the neutron spectrum covariance. Any big differences here may flag an issue related to the adequacy of the energy grid used in the covariance matrix for the purpose of this assessment of the spectral uncertainty contribution.

6.10.2. Output

The output file has 12 segments. There is data for the two neutron benchmark spectra; the three dosimetry cross sections which function as response functions; and two different analysis modes. As detailed in Section 6.10.1, the output listing provides the spectrum-averaged response when the fine group representation is used. The second portion of each segment provides the spectrum contribution to the uncertainty. Table 6.3 summarizes the results. In this case, the covariance representation is adequate for the intended uncertainty calculations, but the fine-group spectrum-averaged response should be used in any analysis.

| Cross Section | Neutron Spectrum | <σ> (mb) [640-grp] | <\sigma> (mb) [cov-grp] | Spct. Unc. % |
|---|------------------------------------|--------------------------|-------------------------------|--------------------|
| ⁵⁹ Co(n,γ) ⁶⁰ Co | ²⁵² Cf(s.f.) | 4.864277 | 5.171 | 2.471 |
| | ²³⁵ U(n _{th}) | 5.026129 | 4.9889 | 0.7529 |
| 109 A g/p 3/\110m A g | ²⁵² Cf(s.f.) | 9.34751 | 9.280 | 0.5645 |
| ¹⁰⁹ Ag(n,γ) ^{110m} Ag | ²³⁵ U(n _{th}) | 9.739893 | 9.6348 | 0.7875 |
| 32 S (n n)32D | ²⁵² Cf(s.f.) | 73.99146 | 75.171 | 0.7412 |
| ³² S(n,p) ³² P | ²³⁵ U(n _{th}) | 66.82598 | 67.909 | 1.156 |

6.11. Cross Section Uncertainty in Spectrum-Averaged Response

Sample input file: example_xsec_unc_spectrum_averaged_response

6.11.1. Description

This "Use Case" provides the uncertainty component due to the knowledge of the response function as propagated into the spectrum-averaged response. This calculation repeats the spectrum folding using the fine 640-group representation of the response and neutron spectrum before it does the uncertainty propagation. That initial step is performed so that the user can inspect the variation between the fine-grouped spectrum-averaged response and the spectrum-averaged response that is

computed using the energy grid used for the characterization of the response covariance. Any big differences here may flag an issue related to the adequacy of the energy grid used in the response covariance matrix for the purpose of this assessment of the uncertainty contribution.

6.11.2. Output

The output file has 12 segments. There is data for the two neutron benchmark spectra; the three dosimetry cross sections; and two different analysis modes. As detailed in Section 6.11.1, the output listing provides the spectrum-averaged response when the fine group representation is used. The second portion of each segment provides the response contribution to the uncertainty. Table 6.3 summarizes the results. In this case, the response function was also identified in a fine 640-group representation, so the fine-group spectrum-averaged response should be identical to the spectrum-averaged response in the second part of each calculation segment.

For these well characterized benchmark neutron fields, comparison with the data in Table 6-3 indicates that the response uncertainty is seen to larger than is the spectral uncertainty component.

Table 6-4. Summary of Response Uncertainty Characteristic of Spectrum/Cross Section Folding

| Response / Cross Section | Neutron Spectrum | <σ> [640-grp] | Resp. Unc. % |
|---|------------------------------------|------------------|--------------------|
| ⁵⁹ Co(n,γ) ⁶⁰ Co | ²⁵² Cf(s.f.) | 4.864277 | 4.078 |
| | ²³⁵ U(n _{th}) | 5.026129 | 3.996 |
| 109 A a/p\110m A a | ²⁵² Cf(s.f.) | 9.34751 | 7.554 |
| ¹⁰⁹ Ag(n,γ) ^{110m} Ag | ²³⁵ U(n _{th}) | 9.739893 | 7.476 |
| 32C/n n\32D | ²⁵² Cf(s.f.) | 73.99146 | 2.484 |
| ³² S(n,p) ³² P | ²³⁵ U(n _{th}) | 66.82598 | 2.534 |

7. CONCLUSION

Xxx

REFERENCES

- [ASTM1855] ASTM E1855-20, Standard Test Method for Use of 2N2222A Silicon Bipolar Transistors as Neutron Spectrum Sensors and Displacement Damage Monitors, Annual Book of ASTM Standards, Vol. 12.02, ASTM International, West Conshohocken, PA, 2020.
- [ASTM261] ASTM E261-20, Standard Practice for Determining Neutron Fluence, Fluence Rate, and Spectra by Radioactivation Techniques, Annual Book of ASTM Standards, Vol. 12.02, ASTM International, West Conshohocken, PA, 2020.
- [ASTM265] ASTM E265-20, Standard Test Method for Measuring Reaction Rates and Fast-Neutron Fluences by Radioactivation of Sulfur-32, Annual Book of ASTM Standards, Vol. 12.02, ASTM International, West Conshohocken, PA, 2020.
- [IRDFF] Trkov, A., Griffin, P.J., Simakov, S.P., Greenwood, L.R., Zolotarev, K.I., Capote, R., et al., "IRDFF-II: A New Neutron Metrology Library", Nuclear Data Sheets, Vol. 163, pp. 1-108, 2020.
- [ENDF] "ENDF-6 Formats Manual: Data Formats and Procedures for the Evaluated Nuclear Data Files ENDF/B-VI, ENDF/B-VII and ENDF/B-VIII", edited by A. Trkov, M. Herman, D.A. Brown, report CSEWG Document ENDF-102, report BNL-203218-2018-INRE, February 1, 2018. Available at: https://www.nndc.bnl.gov/csewg/docs/endf-manual.pdf
- [ENDF8a] D.A. Brown, M.B. Chadwick, R. Capote, et al., "ENDF/B-VIII.0: The 8th Major Release of the Nuclear Reaction Data Library with CIELO-project Cross Sections, New Standards and Thermal Scattering Data," Nuclear Data Sheets, Vol. 148, pp. 1-142, February 2018.
- [ENDFB8b] ENDF/VI Release 8 files are available at: http;://www.nndc.bnl.gov/csewg/summary8.html

DISTRIBUTION

Email—Internal

| Name | Org. | Sandia Email Address |
|----------------------|-------|----------------------|
| Michael Gregson | 00512 | mwgregs@sandia.gov |
| Daniel Dorsey | 00513 | djdorse@sandia.gov |
| Patrick Griffin | 01000 | pjgriff@sandia.gov |
| J. Charles Barbour | 01300 | jcbarbo@sandia.gov |
| E. Frederick Hartman | 01340 | efhart@sandia.gov |
| Bryan Oliver | 01340 | bvolive@sandia.gov |
| Joseph Castro | 01341 | jpcastr@sandia.gov |
| Donald Bruss | 01341 | dbruss@sandia.gov |
| K. Russell DePriest | 01341 | krdepris@sandia.gov |
| Michael McLain | 01343 | mlmclai@sandia.gov |
| Jeffrey Black | 01343 | jefblac@sandia.gov |
| James Bryson | 01344 | jwbryso@sandia.gov |
| Dolores Black | 01344 | dablack@sandia.gov |
| J. Kyle McDonald | 01344 | jkmcdon@sandia.gov |
| Arlen Heger | 01354 | aheger@sandia.gov |
| Biliana Paskaleva | 01356 | bspaska@sandia.gov |
| Matthew Burger | 01380 | mjburge@sandia.gov |
| William Martin | 01384 | wjmarti@sandia.gov |
| Andrew Tonigan | 01384 | amtonig@sandia.gov |
| David Epp | 01390 | dsepp@sandia.gov |
| Edward Bielejec | 01866 | esbiele@sandia.gov |
| Rodney Keith | 08841 | rlkeithsandia.gov |
| Technical Library | 01177 | libref@sandia.gov |

This page left blank

This page left blank



Sandia National Laboratories is a multimission laboratory managed and operated by National Technology & Engineering Solutions of Sandia LLC, a wholly owned subsidiary of Honeywell International Inc. for the U.S. Department of Energy's National Nuclear Security Administration under contract DE-NA0003525.