

← Back

Module 2 Quiz

Graded Quiz • 20 min

- ≡ Hide menu
- Two Sample Binomial Tests
- Relative Risks & Odds Ratios
- Delta Method
- Review
- Practice Quiz: Module 2
- Homework
- Started
- Quiz: Module 2 Quiz
- Submitted

✔ Congratulations! You passed!

Grade received 100%

Latent Submission

To pass 80% or higher

Go to next item

Module 2 Quiz

coach

1. What is the delta method any good for? 1 / 1 point

✔ Correct

Help me practice

Let's chat

✔ Submit your assignment

Try again

2. You are a bookie taking 15 bets of the NFL Ravens game. The odds you give betters determines their payout. If you give 4 to 1 odds against the Ravens winning, then if a person bets on the Ravens winning and they win, you owe them their original dollar, plus an additional \$4 (a total of \$5). If a person bets on the Ravens losing and they lose, then you owe them their dollar back, plus \$0.25 (a total of \$1.25). 1 / 1 point

Due Aug 2, 11:59 PM ISTAttempts 1, sorry 18 hours left

✔ Correct

100%

Your grade

View Feedback

We keep your highest score

3. In a randomly sampled survey of self-reported stress levels from two occupations, the following data were obtained: 1 / 1 point

High Stress Low Stress

Professor 70

Loan Tamer 15

What is the P-value for a Z score test of equality of the proportions of high stress?

(Note the notation $1e-5$, for example, is 1×10^{-5}).

✔ Correct

4. Consider the following data recording case status relative to an environmental exposure 1 / 1 point

Case Control

Exposed 45 21

Unexposed 15 52

What would be the estimated asymptotic standard error for the log relative risk for this data? Consider case status as the outcome. That is, we're interested in evaluating the ratio of the proportion of cases comparing exposed to unexposed groups.

✔ Correct

5. If $x_1 \sim \text{Binomial}(n_1, p_1)$ and $x_2 \sim \text{Binomial}(n_2, p_2)$ and independent $\text{Beta}(2, 2)$ priors are placed on p_1 and p_2 , what is the posterior mean for $p_1 - p_2$? 1 / 1 point

✔ Correct

6. Researchers conducted a blind taste test of Coke versus Pepsi. Each of four people was asked which of two blinded drinks given in random order that they preferred. The data was such that 3 of the 4 people chose Coke. Assuming that this sample is representative, report a P-value for a test of the hypothesis that Coke is preferred to Pepsi using a ""two sided"" exact test. 1 / 1 point

✔ Correct