Hide menu

Summative Course Assessment

Graded Assignment: Summative Course Assessment
3h

Summative Course Assessment

Review Learning Objectives

You ran out of attempts. If you'd like to try again, you will have attempts available in 7h 56m

Assignment details

Due
Feb 11, 11:59 PM IST

Attempts
0 left (1 attempts every 8 hours)

Submitted
Jan 23, 3:35 AM IST

Your grade
To pass you need at least 80%. We keep your highest score.

100%

Your grade: 100%

Your latest: **100%** • Your highest: **100%** • To pass you need at least 80%. We keep your highest score.

Instructions

1. Which of the following is the link function you should use when using the glm() function in R? **1 / 1 point**

(I) binomial, link = "logit"

(II) gaussian, link = "indentity"

(III) Poisson, link = "log"

- ○ I only
- ○ II only
- ○ III only
- ○ I and II only
- ◉ I, II, and III

Correct

Try again **View submission** **See feedback**

Like Dislike Report an issue

2. Which of the following R-command is/are related to setting up a robust regression using the MASS library? **1 / 1 point**

(I) $\mathrm{rlm(y \sim x, data = df, psi = psi.bisquare)}$

(II) $\mathrm{rlm(y \sim x1 + x2, data = df, psi = psi.huber)}$

(III) $\mathrm{glm(y \sim x, data = df, family = binomial)}$

- ○ I only
- ○ II only
- ○ III only
- ◉ I and II only
- ○ I, II, and III

Correct

3. Suppose we have the following Poisson regression model **1 / 1 point**

$$\ln(\mathrm{E(Y|X = x)}) = \ln(\lambda(x)) = 0.23 + 0.05x$$

and consider x increases its value from x_0 to $x_0 + 1$. Compute the value of

$$\frac{\lambda_1 - \lambda_0}{\lambda_0} \times 100\%$$

where $\lambda(x_0) = \lambda_0$ and $\lambda(x_0 + 1) = \lambda_1$.

- ○ 0.23%
- ○ 23%
- ○ 0.05%
- ◉ 5.127%
- ○ 25.9%

Correct

4. Which of the following can be used in the variable selection process? **1 / 1 point**

(I) Ridge regression

(II) Forward stepwise regression

(III) Backward stepwise regression

- ○ I only
- ○ II only
- ○ III only
- ◉ II and III only
- ○ I, II, and III

Correct

5. Let Y denote whether a student get admitted to attend Illinois Tech, with $Y = 1$ as successfully being admitted, and $Y = 0$, otherwise. Also, let **1 / 1 point**

- X_1 denote the math scores (0-100)
- X_2 denote the reading scores (0-100)

of a particular standardize exam that a student need to take to gain admission. We collected n-data and run a logistic regression to obtain

$$\ln\left(\frac{\pi}{1-\pi}\right) = 0.13 + 0.05X_1 + 0.04X_2$$

where

$$\pi = \Pr(Y = 1|X_1 = x_1, X_2 = x_2).$$

Suppose a student has a math scores of 78% and the reading scores of 82%. If the student retake the math test and scores 83%. Compute the following change of odds in being admitted to Illinois Tech.

$$\frac{\mathrm{Odds}(x_0 + 1) - \mathrm{Odds}(x_0)}{\mathrm{Odds}(x_0)} \times 100\%$$

- ◉ 5.127%
- ○ 13.883%
- ○ 4.081%
- ○ 0.05%
- ○ 0.04%

Correct

6. Which of the following is an advantage of using the M-estimation technique? **1 / 1 point**

- ○ To detect and remediate multicollinearity issue in the data.
- ○ To obtain estimations of the regression parameters using the least square method.
- ○ To detect and delete outliers in the data.
- ○ M-estimation is used as a variance stabilizing technique.
- ◉ M-estimation is a robust regression technique used when the data has outliers.

Correct

7. Which of the following can be used to measure how good a model can predict the actual observations. **1 / 1 point**

(I) A high R^2 score.

(II) A high MAE score.

(III) A high RMSE score.

- ◉ I only
- ○ II only
- ○ III only
- ○ II and III only
- ○ I, II, and III

Correct

8. Which of the following statement(s) is/are true about a binary logistic regression? **1 / 1 point**

(I) The predictor variables (z) are continuous, with values between 0 and

(II) The response variable (y) is continuous, with values between 0 and 1.

(III) The outputs of the sigmoid function are continuous, with values between 0 and 1.

- ○ I only
- ○ II only
- ◉ III only
- ○ II and III only
- ○ I, II, and III

Correct

9. In the k-fold cross-validation technique, the data are shuffled randomly and then divided into k equal sub-samples. Which of the following statement(s) is/are true? **1 / 1 point**

(I) We then use all the samples as training samples to be learned from.

(II) We then use $k - 1$ samples as training samples and the k-th sample as a test sample.

(III) We then use $k - 2$ samples as training samples, and 2 samples as test samples.

- ○ I only
- ◉ II only
- ○ III only
- ○ II and III only
- ○ I, II, and III

Correct

10. Suppose we want to model the profitability of a movie (y) based on the opening-weekend revenue (x, in millions of dollar). We let $y = 1$ if the movie is profitable, that is, the revenue is larger than the budget, and not profitable as $y = 0$. Based on the data from 100 movies in the past 50 years, we obtain **1 / 1 point**

$$\log(\mathrm{odds}) = -2.27 + 1.276x$$

Which of the following is/are correct?

(I) The odds ratio is given by $e^{1.276}$.

(II) For a one-unit increase in x, we expect a 258.2% increase in the odds.

(III) For a one-unit increase in x, we expect the odds that the movie will be profitable increase by a factor of 3.58.

- ○ I only
- ○ II only
- ○ III only
- ○ II and III only
- ◉ I, II, and III

Correct

11. Which of the following component(s) is/are involved in a generalized linear model? **1 / 1 point**

(I) An exponential family model for the response variable $\mathbf{y}$.

(II) A link function that connects the means of the response to the linear predictors.

(III) Predictor variables having a normal distribution.

- ○ I only
- ○ II only
- ○ III only
- ◉ I and II only
- ○ I, II, and III

Correct

12. Suppose Y is a binary response variable, taking values either 0 or 1, while X_1 and X_2 are predictor variables. Predicting the probability **1 / 1 point**

$$\Pr(Y = 1|X_1, X_2)$$

involves setting up a

- ○ Multiple linear regression model
- ○ Simple linear regression model
- ○ Poisson regression model
- ◉ Logistic regression model
- ○ Gamma regression model

Correct

13. Which of the following statement(s) is/are true? **1 / 1 point**

(I) We say that a random variable Y follows the Poisson distribution if Y takes values in $[0, 1, 2, \ldots]$ and

$$\Pr(Y = k) = \frac{e^{-\lambda}\lambda^k}{k!}$$

where $k = 0, 1, 2, 3, \cdots$, for some $\lambda > 0$.

(II) If Y is a Poisson random variable, then the mean of Y is the same as the variance of Y .

(III) In Poisson regression, we assume that the mean of a Poisson random variable Y can be written as

$$E(Y|X_1, X_2, \cdots, X_n) = \beta_0 + \beta_1 X_1 + \cdots + \beta_n X_n.$$

- ○ I only
- ○ II only
- ○ III only
- ◉ I and II only
- ○ I, II, and III

Correct

14. Suppose that the probability of an event is 0.2, what is the odds of the event occurring? **1 / 1 point**

- ○ 0.2
- ○ 0.8
- ○ 0.12
- ◉ 0.25
- ○ 0.3

Correct