

[Courseware \(/courses/HumanitiesScience/StatLearning/Winter2014/courseware\)](/courses/HumanitiesScience/StatLearning/Winter2014/courseware)[Course Info \(/courses/HumanitiesScience/StatLearning/Winter2014/info\)](/courses/HumanitiesScience/StatLearning/Winter2014/info)[Discussion \(/courses/HumanitiesScience/StatLearning/Winter2014/discussion/forum\)](/courses/HumanitiesScience/StatLearning/Winter2014/discussion/forum)[Wiki \(/courses/HumanitiesScience/StatLearning/Winter2014/course_wiki\)](/courses/HumanitiesScience/StatLearning/Winter2014/course_wiki)[Progress \(/courses/HumanitiesScience/StatLearning/Winter2014/progress\)](/courses/HumanitiesScience/StatLearning/Winter2014/progress)[Lecture Slides \(pdf\) \(/courses/HumanitiesScience/StatLearning/Winter2014/ba1951b8f66c4cdca2fcaabdc91b792/\)](/courses/HumanitiesScience/StatLearning/Winter2014/ba1951b8f66c4cdca2fcaabdc91b792/)[R Sessions \(/courses/HumanitiesScience/StatLearning/Winter2014/0d68641c19484ef9aa6aeea03426dc68/\)](/courses/HumanitiesScience/StatLearning/Winter2014/0d68641c19484ef9aa6aeea03426dc68/)

9.Q.1 (1/1 point)

Suppose that after our computer works for an hour to fit an SVM on a large data set, we notice that x_4 , the feature vector for the fourth example, was recorded incorrectly (say, one of the decimal points is obviously in the wrong place).

However, your co-worker notices that the pair (x_4, y_4) did not turn out to be a support point in the original fit. He says there is no need to re-fit the SVM on the corrected data set, because changing the value of a non-support point can't possibly change the fit.

Is your co-worker correct?

☐ Yes☒ No**EXPLANATION**

When we change x_4 , the fourth example might become a support point; if so, the fit may change. However, we could check whether x_4, y_4 is still a support point even after correcting the value. If so, then we really don't need to re-fit the model.

[Hide Answer](#)

You have used 2 of 5 submissions



