

[Courseware \(/courses/HumanitiesScience/StatLearning/Winter2014/courseware/\)](/courses/HumanitiesScience/StatLearning/Winter2014/courseware/)[Course Info \(/courses/HumanitiesScience/StatLearning/Winter2014/info/\)](/courses/HumanitiesScience/StatLearning/Winter2014/info/)[Discussion \(/courses/HumanitiesScience/StatLearning/Winter2014/discussion/forum/\)](/courses/HumanitiesScience/StatLearning/Winter2014/discussion/forum/)[Wiki \(/courses/HumanitiesScience/StatLearning/Winter2014/course_wiki/\)](/courses/HumanitiesScience/StatLearning/Winter2014/course_wiki/)[Progress \(/courses/HumanitiesScience/StatLearning/Winter2014/progress/\)](/courses/HumanitiesScience/StatLearning/Winter2014/progress/)[Lecture Slides \(pdf\) \(/courses/HumanitiesScience/StatLearning/Winter2014/ba1951b8f66c4cdca2fcaabdc91b792/\)](/courses/HumanitiesScience/StatLearning/Winter2014/ba1951b8f66c4cdca2fcaabdc91b792/)[R Sessions \(/courses/HumanitiesScience/StatLearning/Winter2014/0d68641c19484ef9aa6aeea03426dc68/\)](/courses/HumanitiesScience/StatLearning/Winter2014/0d68641c19484ef9aa6aeea03426dc68/)

2.3 R1 (1/1 points)

True or False: A fitted model with more predictors will necessarily have a lower Training Set Error than a model with fewer predictors.

False ▾

 False**EXPLANATION**

False. While we typically expect a model with more predictors to have lower Training Set Error, it is not necessarily the case. An extreme counterexample would be a case where you have a model with one predictor that is always equal to the response, compared to a model with many predictors that are random.

Save

Submit

Hide Answer(s)

You have used 2 of 5 submissions



