

Your grade: 100%

Your latest: 100% • Your highest: 100% • To pass you need at least 70%. We keep your highest score.

Next item →

1. When is the Interquartile Range (IQR) useful?

1 / 1 point

- ☐ Analyzing data skewness
- ☐ Applying stratified sampling
- ☒ Identifying outliers

👍 Nice work

IQR is used for identifying outliers and describing the spread of the middle 50% of the data

- ☒ Describing the spread of the middle 50% of data

👍 Nice work

2. How is the Interquartile Range (IQR) defined?

1 / 1 point

- ☐ The difference between the maximum and minimum values
- ☒ The range covered by the middle 50% of the data
- ☐ The standard deviation of the data
- ☐ The average of the data

👍 Nice work

3. What does the z-score represent in statistics?

1 / 1 point

- ☐ The sum of squared errors in a regression model
- ☐ The probability density function of a random variable
- ☒ The number of standard deviations a data point is away from the average
- ☐ The correlation coefficient between two variables

👍 Nice work

4. What are the effects of normalization and standardization on the dataset?

1 / 1 point

- ☒ Scaling data to a specific range

👍 Nice work

- ☐ Reducing outliers
- ☐ Increasing skewness
- ☒ Making data comparable

👍 Nice work

5. What is the primary effect of a power transform on a dataset?

1 / 1 point

- ☐ Increases skewness
- ☐ Reduces dimensionality
- ☒ Stabilizes variance
- ☐ Makes data normally distributed

👍 Nice work