

- ≡ Hide menu
- Week 4 Intro
- Lesson 1: Confidence Intervals
- Lesson 2: Point Estimation, Part I
- Video: Point Estimation4 min
- Video: Variance Estimation: Biased vs Unbiased, Part I11 min
- Graded Assignment: Learning Check - Point Estimation, Part I**Scheduled
- Lesson 3: Point Estimation, Part II
- Weekly Assignment

Your grade: 100%

Your latest: 100%

Your highest: 100%

Next item →

Learning Check - Point Estimation, Part I

Instructions

Review Learning Objectives

1. True or false: A point estimate for a parameter θ gives the range of values that can be regarded as sensible values for θ .1 / 1 point

True

False

Correct

Assignment details

A point estimate is a single number that can be regarded as a sensible value for θ .

DueOct 7, 11:59 PM IST

Attempts4 left (5 attempts every 24 hours)

Try again

Help me practice

Let's chat

2. Statistically speaking, "knowledge" is... (Select all that apply):1 / 1 point

Learning some general pattern from data

Correct

Your grade

Review "Point Estimation: Biased vs Unbiased, Part I" for a full derivation.

100%

View submission

See feedback

Correct

Review "Point Estimation: Biased vs Unbiased, Part I" for a full derivation.

Report an issue

Facts that we know prior to acquiring and analyzing data.

Data that we acquire by analyzing a real world phenomenon

3. Given samples $x_1, \dots, x_n$ from an iid distribution, which of the following is a good estimator for the variance of the distribution (δ^2)?1 / 1 point

$\frac{x_1 + x_2 + \dots + x_n}{n}$

We do not have enough information to estimate the variance.

$\sqrt{\frac{(x_1 - \mu)^2 + (x_2 - \mu)^2 + \dots + (x_n - \mu)^2}{n}}$

$\frac{(x_1 - \mu)^2 + (x_2 - \mu)^2 + \dots + (x_n - \mu)^2}{n}$

Correct

Review "Variance Estimation: Biased vs Unbiased, Part II" for a full derivation.

https://www.coursera.org/learn/statistics-for-data-science-essentials/assignment/submission/2CmL2learning-check-point-estimation-part-i/view-feedback

1/1