

Two dice

$$\mathbf{P(H)} = \mathbf{P(H \mid A)} \mathbf{P(A)} + \mathbf{P(H \mid A^c)} \mathbf{P(A^c)}$$

One die has five red faces and one white face. The other die has five white faces and one red face. A die is chosen at random and thrown twice. What is the chance that the second throw shows red given that the first throw shows red?

$$\mathbf{P}(\mathbf{H}) = \mathbf{P}(\mathbf{H} \mid \mathbf{A}) \mathbf{P}(\mathbf{A}) + \mathbf{P}(\mathbf{H} \mid \mathbf{A}^c) \mathbf{P}(\mathbf{A}^c)$$

$$\mathbf{P}(H) = \mathbf{P}(H \mid A) \mathbf{P}(A) + \mathbf{P}(H \mid A^c) \mathbf{P}(A^c)$$

🟡 *Sample space* Ω : set of triples (coin toss; first face, second face);
sample points $\omega = (x; y_1, y_2)$, $x \in \{1, 2\}$, $y_1, y_2 \in \{\text{red}, \text{white}\}$.

$$\mathbf{P}(H) = \mathbf{P}(H \mid A) \mathbf{P}(A) + \mathbf{P}(H \mid A^c) \mathbf{P}(A^c)$$

🟡 *Sample space* Ω : set of triples (coin toss; first face, second face);
sample points $\omega = (x; y_1, y_2)$, $x \in \{1, 2\}$, $y_1, y_2 \in \{\text{red}, \text{white}\}$.

🟡 *The events of interest:*

$R_1 := \text{first throw shows red} = \{ (x; y_1, y_2) : y_1 = \text{red} \}$.

$R_2 := \text{second throw shows red} = \{ (x; y_1, y_2) : y_2 = \text{red} \}$.

$$\mathbf{P}(H) = \mathbf{P}(H \mid A) \mathbf{P}(A) + \mathbf{P}(H \mid A^c) \mathbf{P}(A^c)$$

🟡 *Sample space* Ω : set of triples (coin toss; first face, second face);
sample points $\omega = (x; y_1, y_2)$, $x \in \{1, 2\}$, $y_1, y_2 \in \{\text{red}, \text{white}\}$.

🟡 *The events of interest:*

$R_1 :=$ first throw shows red $= \{ (x; y_1, y_2) : y_1 = \text{red} \}$.

$R_2 :=$ second throw shows red $= \{ (x; y_1, y_2) : y_2 = \text{red} \}$.

$A :=$ first die is chosen $= \{ (x; y_1, y_2) : x = 1 \}$.

$$\mathbf{P}(H) = \mathbf{P}(H \mid A) \mathbf{P}(A) + \mathbf{P}(H \mid A^c) \mathbf{P}(A^c)$$

🟡 *Sample space* Ω : set of triples (coin toss; first face, second face);
sample points $\omega = (x; y_1, y_2)$, $x \in \{1, 2\}$, $y_1, y_2 \in \{\text{red}, \text{white}\}$.

🟡 The *events of interest*:

$R_1 :=$ first throw shows red $= \{ (x; y_1, y_2) : y_1 = \text{red} \}$.

$R_2 :=$ second throw shows red $= \{ (x; y_1, y_2) : y_2 = \text{red} \}$.

$A :=$ first die is chosen $= \{ (x; y_1, y_2) : x = 1 \}$.

🟡 *Implicit probability measure* \mathbf{P} :

$$\mathbf{P}(H) = \mathbf{P}(H \mid A) \mathbf{P}(A) + \mathbf{P}(H \mid A^c) \mathbf{P}(A^c)$$

🟡 *Sample space* Ω : set of triples (coin toss; first face, second face);
sample points $\omega = (x; y_1, y_2)$, $x \in \{1, 2\}$, $y_1, y_2 \in \{\text{red}, \text{white}\}$.

🟡 The *events of interest*:

$R_1 :=$ first throw shows red $= \{ (x; y_1, y_2) : y_1 = \text{red} \}$.

$R_2 :=$ second throw shows red $= \{ (x; y_1, y_2) : y_2 = \text{red} \}$.

$A :=$ first die is chosen $= \{ (x; y_1, y_2) : x = 1 \}$.

🟡 Implicit *probability measure* \mathbf{P} :

- Random selection of die: $\mathbf{P}(A) = \mathbf{P}(A^c) = 1/2$.

$$\mathbf{P}(H) = \mathbf{P}(H \mid A) \mathbf{P}(A) + \mathbf{P}(H \mid A^c) \mathbf{P}(A^c)$$

🟡 *Sample space* Ω : set of triples (coin toss; first face, second face);
sample points $\omega = (x; y_1, y_2)$, $x \in \{1, 2\}$, $y_1, y_2 \in \{\text{red}, \text{white}\}$.

🟡 The *events of interest*:

$R_1 :=$ first throw shows red $= \{ (x; y_1, y_2) : y_1 = \text{red} \}$.

$R_2 :=$ second throw shows red $= \{ (x; y_1, y_2) : y_2 = \text{red} \}$.

$A :=$ first die is chosen $= \{ (x; y_1, y_2) : x = 1 \}$.

🟡 Implicit *probability measure* \mathbf{P} :

- Random selection of die: $\mathbf{P}(A) = \mathbf{P}(A^c) = 1/2$.
- Conditional probabilities for a given die:

$$\mathbf{P}(H) = \mathbf{P}(H \mid A) \mathbf{P}(A) + \mathbf{P}(H \mid A^c) \mathbf{P}(A^c)$$

🏆 *Sample space* Ω : set of triples (coin toss; first face, second face);
sample points $\omega = (x; y_1, y_2)$, $x \in \{1, 2\}$, $y_1, y_2 \in \{\text{red}, \text{white}\}$.

🏆 *The events of interest:*

$R_1 :=$ first throw shows red $= \{ (x; y_1, y_2) : y_1 = \text{red} \}$.

$R_2 :=$ second throw shows red $= \{ (x; y_1, y_2) : y_2 = \text{red} \}$.

$A :=$ first die is chosen $= \{ (x; y_1, y_2) : x = 1 \}$.

🏆 *Implicit probability measure* \mathbf{P} :

- Random selection of die: $\mathbf{P}(A) = \mathbf{P}(A^c) = 1/2$.

- Conditional probabilities for a given die:

$$\mathbf{P}(R_1 \mid A) = 5/6,$$

$$\mathbf{P}(R_1 \mid A^c) = 1/6,$$

$$\mathbf{P}(H) = \mathbf{P}(H \mid A) \mathbf{P}(A) + \mathbf{P}(H \mid A^c) \mathbf{P}(A^c)$$

🟡 *Sample space* Ω : set of triples (coin toss; first face, second face);
sample points $\omega = (x; y_1, y_2)$, $x \in \{1, 2\}$, $y_1, y_2 \in \{\text{red}, \text{white}\}$.

🟡 *The events of interest:*

$R_1 :=$ first throw shows red $= \{ (x; y_1, y_2) : y_1 = \text{red} \}$.

$R_2 :=$ second throw shows red $= \{ (x; y_1, y_2) : y_2 = \text{red} \}$.

$A :=$ first die is chosen $= \{ (x; y_1, y_2) : x = 1 \}$.

🟡 *Implicit probability measure* \mathbf{P} :

- Random selection of die: $\mathbf{P}(A) = \mathbf{P}(A^c) = 1/2$.

- Conditional probabilities for a given die:

$$\mathbf{P}(R_1 \mid A) = 5/6,$$

$$\mathbf{P}(R_1 \mid A^c) = 1/6,$$

$$\mathbf{P}(R_1 \cap R_2 \mid A) = 5^2/6^2 = 25/36, \quad \mathbf{P}(R_1 \cap R_2 \mid A^c) = 1^2/6^2 = 1/36.$$

$$\mathbf{P}(H) = \mathbf{P}(H \mid A) \mathbf{P}(A) + \mathbf{P}(H \mid A^c) \mathbf{P}(A^c)$$

$$\begin{aligned} \mathbf{P}(R_1) &= \mathbf{P}(R_1 \mid A) \mathbf{P}(A) + \mathbf{P}(R_1 \mid A^c) \mathbf{P}(A^c) \\ &= \frac{5}{6} \cdot \frac{1}{2} + \frac{1}{6} \cdot \frac{1}{2} = \frac{1}{2} = 0.5 \end{aligned}$$

• *Sample space* Ω : set of triples (coin toss; first face, second face); sample points $\omega = (x; y_1, y_2)$, $x \in \{1, 2\}$, $y_1, y_2 \in \{\text{red}, \text{white}\}$.

• The *events of interest*:

$R_1 :=$ first throw shows red $= \{ (x; y_1, y_2) : y_1 = \text{red} \}$.

$R_2 :=$ second throw shows red $= \{ (x; y_1, y_2) : y_2 = \text{red} \}$.

$A :=$ first die is chosen $= \{ (x; y_1, y_2) : x = 1 \}$.

• Implicit *probability measure* \mathbf{P} :

- Random selection of die: $\mathbf{P}(A) = \mathbf{P}(A^c) = 1/2$.

- Conditional probabilities for a given die:

$$\mathbf{P}(R_1 \mid A) = 5/6,$$

$$\mathbf{P}(R_1 \mid A^c) = 1/6,$$

$$\mathbf{P}(R_1 \cap R_2 \mid A) = 5^2/6^2 = 25/36, \quad \mathbf{P}(R_1 \cap R_2 \mid A^c) = 1^2/6^2 = 1/36.$$

$$\mathbf{P}(H) = \mathbf{P}(H \mid A) \mathbf{P}(A) + \mathbf{P}(H \mid A^c) \mathbf{P}(A^c)$$

$$\begin{aligned} \mathbf{P}(R_1) &= \mathbf{P}(R_1 \mid A) \mathbf{P}(A) + \mathbf{P}(R_1 \mid A^c) \mathbf{P}(A^c) \\ &= \frac{5}{6} \cdot \frac{1}{2} + \frac{1}{6} \cdot \frac{1}{2} = \frac{1}{2} = 0.5 \end{aligned}$$

$$\begin{aligned} \mathbf{P}(R_1 \cap R_2) &= \mathbf{P}(R_1 \cap R_2 \mid A) \mathbf{P}(A) + \mathbf{P}(R_1 \cap R_2 \mid A^c) \mathbf{P}(A^c) \\ &= \frac{25}{36} \cdot \frac{1}{2} + \frac{1}{36} \cdot \frac{1}{2} = \frac{13}{18} \cdot \frac{1}{2} = 0.36\bar{1} \end{aligned}$$

👉 *Sample space Ω* : set of triples (coin toss; first face, second face); sample points $\omega = (x; y_1, y_2)$, $x \in \{1, 2\}$, $y_1, y_2 \in \{\text{red, white}\}$.

👉 *The events of interest*:

$R_1 :=$ first throw shows red $= \{ (x; y_1, y_2) : y_1 = \text{red} \}$.

$R_2 :=$ second throw shows red $= \{ (x; y_1, y_2) : y_2 = \text{red} \}$.

$A :=$ first die is chosen $= \{ (x; y_1, y_2) : x = 1 \}$.

👉 *Implicit probability measure \mathbf{P}* :

- Random selection of die: $\mathbf{P}(A) = \mathbf{P}(A^c) = 1/2$.

- Conditional probabilities for a given die:

$$\mathbf{P}(R_1 \mid A) = 5/6,$$

$$\mathbf{P}(R_1 \mid A^c) = 1/6,$$

$$\mathbf{P}(R_1 \cap R_2 \mid A) = 5^2/6^2 = 25/36, \quad \mathbf{P}(R_1 \cap R_2 \mid A^c) = 1^2/6^2 = 1/36.$$

$$\mathbf{P}(H) = \mathbf{P}(H \mid A) \mathbf{P}(A) + \mathbf{P}(H \mid A^c) \mathbf{P}(A^c)$$

$$\begin{aligned} \mathbf{P}(R_1) &= \mathbf{P}(R_1 \mid A) \mathbf{P}(A) + \mathbf{P}(R_1 \mid A^c) \mathbf{P}(A^c) \\ &= \frac{5}{6} \cdot \frac{1}{2} + \frac{1}{6} \cdot \frac{1}{2} = \frac{1}{2} = 0.5 \end{aligned}$$

$$\begin{aligned} \mathbf{P}(R_1 \cap R_2) &= \mathbf{P}(R_1 \cap R_2 \mid A) \mathbf{P}(A) + \mathbf{P}(R_1 \cap R_2 \mid A^c) \mathbf{P}(A^c) \\ &= \frac{25}{36} \cdot \frac{1}{2} + \frac{1}{36} \cdot \frac{1}{2} = \frac{13}{18} \cdot \frac{1}{2} = 0.36\bar{1} \end{aligned}$$

$$\mathbf{P}(R_2 \mid R_1) = \frac{\mathbf{P}(R_1 \cap R_2)}{\mathbf{P}(R_1)} = \frac{\frac{13}{18} \cdot \frac{1}{2}}{\frac{1}{2}} = \frac{13}{18} = 0.72\bar{2}$$

👉 *Sample space Ω* : set of triples (coin toss; first face, second face); sample points $\omega = (x; y_1, y_2)$, $x \in \{1, 2\}$, $y_1, y_2 \in \{\text{red, white}\}$.

👉 *The events of interest*:

$R_1 :=$ first throw shows red $= \{ (x; y_1, y_2) : y_1 = \text{red} \}$.

$R_2 :=$ second throw shows red $= \{ (x; y_1, y_2) : y_2 = \text{red} \}$.

$A :=$ first die is chosen $= \{ (x; y_1, y_2) : x = 1 \}$.

👉 *Implicit probability measure \mathbf{P}* :

- Random selection of die: $\mathbf{P}(A) = \mathbf{P}(A^c) = 1/2$.

- Conditional probabilities for a given die:

$$\mathbf{P}(R_1 \mid A) = 5/6,$$

$$\mathbf{P}(R_1 \mid A^c) = 1/6,$$

$$\mathbf{P}(R_1 \cap R_2 \mid A) = 5^2/6^2 = 25/36, \quad \mathbf{P}(R_1 \cap R_2 \mid A^c) = 1^2/6^2 = 1/36.$$

$$\mathbf{P}(H) = \mathbf{P}(H \mid A) \mathbf{P}(A) + \mathbf{P}(H \mid A^c) \mathbf{P}(A^c)$$

$$\begin{aligned} \mathbf{P}(R_1) &= \mathbf{P}(R_1 \mid A) \mathbf{P}(A) + \mathbf{P}(R_1 \mid A^c) \mathbf{P}(A^c) \\ &= \frac{5}{6} \cdot \frac{1}{2} + \frac{1}{6} \cdot \frac{1}{2} = \frac{1}{2} = 0.5 \end{aligned}$$

$$\begin{aligned} \mathbf{P}(R_1 \cap R_2) &= \mathbf{P}(R_1 \cap R_2 \mid A) \mathbf{P}(A) + \mathbf{P}(R_1 \cap R_2 \mid A^c) \mathbf{P}(A^c) \\ &= \frac{25}{36} \cdot \frac{1}{2} + \frac{1}{36} \cdot \frac{1}{2} = \frac{13}{18} \cdot \frac{1}{2} = 0.36\bar{1} \end{aligned}$$

$$\mathbf{P}(R_2 \mid R_1) = \frac{\mathbf{P}(R_1 \cap R_2)}{\mathbf{P}(R_1)} = \frac{\frac{13}{18} \cdot \frac{1}{2}}{\frac{1}{2}} = \frac{13}{18} = 0.72\bar{2}$$

👉 *Sample space* Ω : set of triples (coin toss; first face, second face); sample points $\omega = (x; y_1, y_2)$, $x \in \{1, 2\}$, $y_1, y_2 \in \{\text{red, white}\}$.

👉 *The events of interest*:

$R_1 :=$ first throw shows red $= \{ (x; y_1, y_2) : y_1 = \text{red} \}$.

$R_2 :=$ second throw shows red $= \{ (x; y_1, y_2) : y_2 = \text{red} \}$.

$A :=$ first die is chosen $= \{ (x; y_1, y_2) : x = 1 \}$.

👉 *Implicit probability measure* \mathbf{P} :

- Random selection of die: $\mathbf{P}(A) = \mathbf{P}(A^c) = 1/2$.

- Conditional probabilities for a given die:

$$\mathbf{P}(R_1 \mid A) = 5/6,$$

$$\mathbf{P}(R_1 \mid A^c) = 1/6,$$

$$\mathbf{P}(R_1 \cap R_2 \mid A) = 5^2/6^2 = 25/36, \quad \mathbf{P}(R_1 \cap R_2 \mid A^c) = 1^2/6^2 = 1/36.$$

What is the mass function for this problem?