

Additivity, third time pays for all: milking independent trials

$$s_n = f_1 + f_2 + \cdots + f_n$$

$$f_n = u_n - f_1 u_{n-1} - f_2 u_{n-2} - \cdots - f_j u_{n-j} - \cdots - f_{n-1} u_1$$

---



Additivity, third time pays for all: milking independent trials

$$s_n = f_1 + f_2 + \cdots + f_n$$

$$f_n = u_n - f_1 u_{n-1} - f_2 u_{n-2} - \cdots - f_j u_{n-j} - \cdots - f_{n-1} u_1$$

---

**Key observation:** if there is a run of five consecutive successes from trial  $n - 4$  through  $n$ , then a success run must have occurred at one of the trials  $n - 4$ ,  $n - 3$ ,  $n - 2$ ,  $n - 1$ , or  $n$ .



# Additivity, third time pays for all: milking independent trials

$$s_n = f_1 + f_2 + \cdots + f_n$$

$$f_n = u_n - f_1 u_{n-1} - f_2 u_{n-2} - \cdots - f_j u_{n-j} - \cdots - f_{n-1} u_1$$

**Key observation:** if there is a run of five consecutive successes from trial  $n - 4$  through  $n$ , then a success run must have occurred at one of the trials  $n - 4$ ,  $n - 3$ ,  $n - 2$ ,  $n - 1$ , or  $n$ .

																								1	1	1	1	1	
																								1	1	1	1	1	
																							1	1	1	1	1	1	
																						1	1	1	1	1	1	1	
																					1	1	1	1	1	1	1	1	
																				1	1	1	1	1	1	1	1	1	
1	2	3	4	5	6																1	1	1	1	1	1	1	1	
																								n-5	n-4	n-3	n-2	n-1	n



## Additivity, third time pays for all: milking independent trials

$$s_n = f_1 + f_2 + \cdots + f_n$$

$$f_n = u_n - f_1 u_{n-1} - f_2 u_{n-2} - \cdots - f_j u_{n-j} - \cdots - f_{n-1} u_1$$

**Key observation:** if there is a run of five consecutive successes from trial  $n - 4$  through  $n$ , then a success run must have occurred at one of the trials  $n - 4$ ,  $n - 3$ ,  $n - 2$ ,  $n - 1$ , or  $n$ .

																									1	1	1	1	1			
																								1	1	1	1	1				
																								1	1	1	1	1				
																								1	1	1	1	1				
																								1	1	1	1	1				
																								1	1	1	1	1				
																								1	1	1	1	1				
1	2	3	4	5	6																						n-5	n-4	n-3	n-2	n-1	n



# Additivity, third time pays for all: milking independent trials

$$s_n = f_1 + f_2 + \cdots + f_n$$

$$f_n = u_n - f_1 u_{n-1} - f_2 u_{n-2} - \cdots - f_j u_{n-j} - \cdots - f_{n-1} u_1$$

**Key observation:** if there is a run of five consecutive successes from trial  $n - 4$  through  $n$ , then a success run must have occurred at one of the trials  $n - 4$ ,  $n - 3$ ,  $n - 2$ ,  $n - 1$ , or  $n$ .

																									1	1	1	1	1		
																								1	1	1	1	1			
																							1	1	1	1	1	1			
																							1	1	1	1	1	1			
																						1	1	1	1	1	1	1			
																					1	1	1	1	1	1	1	1			
1	2	3	4	5	6																					n-5	n-4	n-3	n-2	n-1	n

**Additivity!**  $2^{-5} = u_n + u_{n-1}2^{-1} + u_{n-2}2^{-2} + u_{n-3}2^{-3} + u_{n-4}2^{-4}$



## Additivity, third time pays for all: milking independent trials

$$s_n = f_1 + f_2 + \cdots + f_n$$

$$f_n = u_n - f_1 u_{n-1} - f_2 u_{n-2} - \cdots - f_j u_{n-j} - \cdots - f_{n-1} u_1$$

**Key observation:** if there is a run of five consecutive successes from trial  $n - 4$  through  $n$ , then a success run must have occurred at one of the trials  $n - 4$ ,  $n - 3$ ,  $n - 2$ ,  $n - 1$ , or  $n$ .

																									1	1	1	1	1			
																								1	1	1	1	1				
																								1	1	1	1	1				
																								1	1	1	1	1				
																								1	1	1	1	1				
																								1	1	1	1	1				
																								1	1	1	1	1				
																								1	1	1	1	1				
1	2	3	4	5	6																						n-5	n-4	n-3	n-2	n-1	n

**Additivity!**  $2^{-5} = u_n + u_{n-1}2^{-1} + u_{n-2}2^{-2} + u_{n-3}2^{-3} + u_{n-4}2^{-4}$

— or equivalently —



# Additivity, third time pays for all: milking independent trials

$$s_n = f_1 + f_2 + \cdots + f_n$$

$$f_n = u_n - f_1 u_{n-1} - f_2 u_{n-2} - \cdots - f_j u_{n-j} - \cdots - f_{n-1} u_1$$

**Key observation:** if there is a run of five consecutive successes from trial  $n - 4$  through  $n$ , then a success run must have occurred at one of the trials  $n - 4$ ,  $n - 3$ ,  $n - 2$ ,  $n - 1$ , or  $n$ .

																									1	1	1	1	1
																								1	1	1	1	1	
																								1	1	1	1	1	
																								1	1	1	1	1	
																								1	1	1	1	1	
																								1	1	1	1	1	
																								1	1	1	1	1	
																								1	1	1	1	1	
1	2	3	4	5	6																								

**Additivity!**  $2^{-5} = u_n + u_{n-1}2^{-1} + u_{n-2}2^{-2} + u_{n-3}2^{-3} + u_{n-4}2^{-4}$

— or equivalently —

$$u_n = 2^{-5} - u_{n-1}2^{-1} - u_{n-2}2^{-2} - u_{n-3}2^{-3} - u_{n-4}2^{-4}$$



## Additivity, third time pays for all: milking independent trials

$$s_n = f_1 + f_2 + \cdots + f_n$$

$$f_n = u_n - f_1 u_{n-1} - f_2 u_{n-2} - \cdots - f_j u_{n-j} - \cdots - f_{n-1} u_1$$

**Key observation:** if there is a run of five consecutive successes from trial  $n - 4$  through  $n$ , then a success run must have occurred at one of the trials  $n - 4$ ,  $n - 3$ ,  $n - 2$ ,  $n - 1$ , or  $n$ .

																										1	1	1	1	1
																									1	1	1	1	1	1
																								1	1	1	1	1	1	1
																								1	1	1	1	1	1	1
																								1	1	1	1	1	1	1
																								1	1	1	1	1	1	1
1	2	3	4	5	6																									

**Additivity!**  $2^{-5} = u_n + u_{n-1}2^{-1} + u_{n-2}2^{-2} + u_{n-3}2^{-3} + u_{n-4}2^{-4}$

— or equivalently —

$$u_n = 2^{-5} - u_{n-1}2^{-1} - u_{n-2}2^{-2} - u_{n-3}2^{-3} - u_{n-4}2^{-4}$$

Slogan: If you know  $u_1, u_2, \dots, u_{n-1}$  then you know  $u_n$ .