

## Lesson 4

[Back to Week 2](#)

**6/6** points  
earned (100%)

Quiz passed!



**Be Recognized for Your Achievements.** "Course Certificates give you the recognition you need to get the job, the material gives you the skills to do the job. It makes you look more valuable because you are more valuable." - Peter B., USA, Software Developer

**Showcase Your Accomplishment! Earn Your Course Certificate! ₹1,953 ➔**



1 / 1  
points

**For Questions 1-3, consider the following scenario:**

In the example from Lesson 4.1 of flipping a coin 100 times, suppose instead that you observe 47 heads and 53 tails.

1. • Report the value of  $\hat{p}$ , the MLE (Maximum Likelihood Estimate) of the probability of obtaining heads.

**Correct Response**

This is simply  $47/100$ , the number of successes divided by the number of trials.

---



1 / 1  
points

2.

Coin flip:

Using the central limit theorem as an approximation, and following the example of Lesson 4.1, construct a 95% confidence interval for  $p$ , the probability of obtaining heads.

- Report the lower end of this interval and round your answer to two decimal places.

0.37

**Correct Response**

We have  $\hat{p} - 1.96\sqrt{\hat{p}(1-\hat{p})/n} = .47 - 1.96\sqrt{(.47)(.53)/100} = .372$ , which is the lower end of a 95% confidence interval for  $p$ .

---



1 / 1  
points

Coin flip:

3. • Report the upper end of this interval and round your answer to two decimal places.

0.57

▲

**Correct Response**

We have  $\hat{p} + 1.96\sqrt{\hat{p}(1-\hat{p})/n} = .47 + 1.96\sqrt{(.47)(.53)/100} = .568$ , which is the upper end of a 95% confidence interval for  $p$ .

---



1 / 1  
points

4.

The likelihood function for parameter  $\theta$  with data  $\mathbf{y}$  is based on which of the following?

☐  $P(\theta \mid \mathbf{y})$

☒  $P(\mathbf{y} \mid \theta)$

▲

**Correct Response**

The likelihood is based on the sampling distribution of the data, given the parameter. Note that although the likelihood has the same functional form as  $P(\mathbf{y} \mid \theta)$ , it is considered a function of  $\theta$ .

☐  $P(\theta)$

☐  $P(\mathbf{y})$

☐ None of the above.

---

1 / 1  
points

5.

Recall from Lesson 4.4 that if  $X_1, \dots, X_n \stackrel{\text{iid}}{\sim} \text{Exponential}(\lambda)$  (iid means independent and identically distributed), then the MLE for  $\lambda$  is  $1/\bar{x}$  where  $\bar{x}$  is the sample mean. Suppose we observe the following data:

$X_1 = 2.0, X_2 = 2.5, X_3 = 4.1, X_4 = 1.8, X_5 = 4.0$ .

Calculate the MLE for  $\lambda$ . Round your answer to two decimal places.

**Correct Response**

The sample mean is  $\bar{x} = 2.88$ .

1 / 1  
points

6.

It turns out that the sample mean  $\bar{x}$  is involved in the MLE calculation for several models. In fact, if the data are independent and identically distributed from a Bernoulli( $p$ ), Poisson( $\lambda$ ), or Normal( $\mu, \sigma^2$ ), then  $\bar{x}$  is the MLE for  $p$ ,  $\lambda$ , and  $\mu$  respectively.

Suppose we observe  $n = 4$  data points from a normal distribution with unknown mean  $\mu$ . The data are  $\mathbf{x} = \{-1.2, 0.5, 0.8, -0.3\}$ .

What is the MLE for  $\mu$ ? Round your answer to two decimal places.

**Correct Response**

This is  $(-1.2 + 0.5 + 0.8 - 0.3)/4$ .

---

---

