

Siméon Denis Poisson



Siméon Denis Poisson

❖ θ : *a priori* probability that the accused is guilty.

Indicator of guilt: $Z \sim \text{Bernoulli}(\theta)$.



Siméon Denis Poisson

- ❖ θ : *a priori* probability that the accused is guilty.
Indicator of guilt: $Z \sim \text{Bernoulli}(\theta)$.
- ❖ p : probability that a juror makes an error in decision.
Error indicator: $X \sim \text{Bernoulli}(p)$.



Siméon Denis Poisson

- ❖ θ : *a priori* probability that the accused is guilty.
Indicator of guilt: $Z \sim \text{Bernoulli}(\theta)$.
- ❖ p : probability that a juror makes an error in decision.
Error indicator: $X \sim \text{Bernoulli}(p)$.



In the years 1825–1830 in France, juries were comprised of twelve individuals with conviction requiring a vote of seven or more jurists voting to convict.

A probability space for jury verdicts

A probability space for jury verdicts

Sample points built from independent trials: List of guilt and juror error indicators ($Z; X_1, \dots, X_{12}$).

A probability space for jury verdicts

Sample points built from independent trials: List of guilt and juror error indicators ($Z; X_1, \dots, X_{12}$).

Probability measure: $Z \sim \text{Bernoulli}(\theta); X_1, \dots, X_{12} \sim \text{Bernoulli}(p)$.

A probability space for jury verdicts

Sample points built from independent trials: List of guilt and juror error indicators $(Z; X_1, \dots, X_{12})$.

Probability measure: $Z \sim \text{Bernoulli}(\theta)$; $X_1, \dots, X_{12} \sim \text{Bernoulli}(p)$.

✦ Number of juror errors: $S_{12} = X_1 + \dots + X_{12} \sim \text{Binomial}(12, p)$.

A probability space for jury verdicts

Sample points built from independent trials: List of guilt and juror error indicators $(Z; X_1, \dots, X_{12})$.

Probability measure: $Z \sim \text{Bernoulli}(\theta)$; $X_1, \dots, X_{12} \sim \text{Bernoulli}(p)$.

* Number of juror errors: $S_{12} = X_1 + \dots + X_{12} \sim \text{Binomial}(12, p)$.

Guilt indicator: $Z = \begin{cases} 1 & \text{with probability } \theta, \\ 0 & \text{with probability } 1 - \theta. \end{cases}$

A probability space for jury verdicts

Sample points built from independent trials: List of guilt and juror error indicators ($Z; X_1, \dots, X_{12}$).

Probability measure: $Z \sim \text{Bernoulli}(\theta)$; $X_1, \dots, X_{12} \sim \text{Bernoulli}(p)$.

* Number of juror errors: $S_{12} = X_1 + \dots + X_{12} \sim \text{Binomial}(12, p)$.

Guilt indicator: $Z = \begin{cases} 1 & \text{with probability } \theta, \\ 0 & \text{with probability } 1 - \theta. \end{cases}$

Juror error indicators: $X_1, \dots, X_{12} = \begin{cases} 1 & \text{with probability } p, \\ 0 & \text{with probability } 1 - p. \end{cases}$

A probability space for jury verdicts

Sample points built from independent trials: List of guilt and juror error indicators $(Z; X_1, \dots, X_{12})$.

Probability measure: $Z \sim \text{Bernoulli}(\theta)$; $X_1, \dots, X_{12} \sim \text{Bernoulli}(p)$.

* Number of juror errors: $S_{12} = X_1 + \dots + X_{12} \sim \text{Binomial}(12, p)$.

Guilt indicator: $Z = \begin{cases} 1 & \text{with probability } \theta, \\ 0 & \text{with probability } 1 - \theta. \end{cases}$

Juror error indicators: $X_1, \dots, X_{12} = \begin{cases} 1 & \text{with probability } p, \\ 0 & \text{with probability } 1 - p. \end{cases}$

$$\mathbf{P}\{S_{12} = k\} = b_{12}(k; p) = \binom{12}{k} p^k (1 - p)^{12-k} \quad (k = 0, 1, \dots, 12)$$

A probability space for jury verdicts

Sample points built from independent trials: List of guilt and juror error indicators $(Z; X_1, \dots, X_{12})$.

Probability measure: $Z \sim \text{Bernoulli}(\theta)$; $X_1, \dots, X_{12} \sim \text{Bernoulli}(p)$.

* Number of juror errors: $S_{12} = X_1 + \dots + X_{12} \sim \text{Binomial}(12, p)$.

Guilt indicator: $Z = \begin{cases} 1 & \text{with probability } \theta, \\ 0 & \text{with probability } 1 - \theta. \end{cases}$

Juror error indicators: $X_1, \dots, X_{12} = \begin{cases} 1 & \text{with probability } p, \\ 0 & \text{with probability } 1 - p. \end{cases}$

$$\mathbf{P}\{S_{12} = k\} = b_{12}(k; p) = \binom{12}{k} p^k (1 - p)^{12-k} \quad (k = 0, 1, \dots, 12)$$

Relevant events:

A probability space for jury verdicts

Sample points built from independent trials: List of guilt and juror error indicators $(Z; X_1, \dots, X_{12})$.

Probability measure: $Z \sim \text{Bernoulli}(\theta)$; $X_1, \dots, X_{12} \sim \text{Bernoulli}(p)$.

* Number of juror errors: $S_{12} = X_1 + \dots + X_{12} \sim \text{Binomial}(12, p)$.

Guilt indicator: $Z = \begin{cases} 1 & \text{with probability } \theta, \\ 0 & \text{with probability } 1 - \theta. \end{cases}$

Juror error indicators: $X_1, \dots, X_{12} = \begin{cases} 1 & \text{with probability } p, \\ 0 & \text{with probability } 1 - p. \end{cases}$

$$\mathbf{P}\{S_{12} = k\} = b_{12}(k; p) = \binom{12}{k} p^k (1 - p)^{12-k} \quad (k = 0, 1, \dots, 12)$$

Relevant events:

* $G :=$ the accused is guilty. $I = G^c :=$ the accused is innocent.

A probability space for jury verdicts

Sample points built from independent trials: List of guilt and juror error indicators ($Z; X_1, \dots, X_{12}$).

Probability measure: $Z \sim \text{Bernoulli}(\theta)$; $X_1, \dots, X_{12} \sim \text{Bernoulli}(p)$.

* Number of juror errors: $S_{12} = X_1 + \dots + X_{12} \sim \text{Binomial}(12, p)$.

Guilt indicator: $Z = \begin{cases} 1 & \text{with probability } \theta, \\ 0 & \text{with probability } 1 - \theta. \end{cases}$

Juror error indicators: $X_1, \dots, X_{12} = \begin{cases} 1 & \text{with probability } p, \\ 0 & \text{with probability } 1 - p. \end{cases}$

$$\mathbf{P}\{S_{12} = k\} = b_{12}(k; p) = \binom{12}{k} p^k (1 - p)^{12-k} \quad (k = 0, 1, \dots, 12)$$

Relevant events:

* $G :=$ the accused is guilty. $I = G^c :=$ the accused is innocent.

* $C :=$ the accused is convicted. $A = C^c :=$ the accused is acquitted.

A probability space for jury verdicts

Sample points built from independent trials: List of guilt and juror error indicators ($Z; X_1, \dots, X_{12}$).

Probability measure: $Z \sim \text{Bernoulli}(\theta)$; $X_1, \dots, X_{12} \sim \text{Bernoulli}(p)$.

* Number of juror errors: $S_{12} = X_1 + \dots + X_{12} \sim \text{Binomial}(12, p)$.

Guilt indicator: $Z = \begin{cases} 1 & \text{with probability } \theta, \\ 0 & \text{with probability } 1 - \theta. \end{cases}$

Juror error indicators: $X_1, \dots, X_{12} = \begin{cases} 1 & \text{with probability } p, \\ 0 & \text{with probability } 1 - p. \end{cases}$

$$\mathbf{P}\{S_{12} = k\} = b_{12}(k; p) = \binom{12}{k} p^k (1 - p)^{12-k} \quad (k = 0, 1, \dots, 12)$$

Relevant events:

* $G :=$ the accused is guilty. $I = G^c :=$ the accused is innocent.

G occurs if $Z = 1$

* $C :=$ the accused is convicted. $A = C^c :=$ the accused is acquitted.

A probability space for jury verdicts

Sample points built from independent trials: List of guilt and juror error indicators $(Z; X_1, \dots, X_{12})$.

Probability measure: $Z \sim \text{Bernoulli}(\theta)$; $X_1, \dots, X_{12} \sim \text{Bernoulli}(p)$.

* Number of juror errors: $S_{12} = X_1 + \dots + X_{12} \sim \text{Binomial}(12, p)$.

Guilt indicator: $Z = \begin{cases} 1 & \text{with probability } \theta, \\ 0 & \text{with probability } 1 - \theta. \end{cases}$

Juror error indicators: $X_1, \dots, X_{12} = \begin{cases} 1 & \text{with probability } p, \\ 0 & \text{with probability } 1 - p. \end{cases}$

$$\mathbf{P}\{S_{12} = k\} = b_{12}(k; p) = \binom{12}{k} p^k (1 - p)^{12-k} \quad (k = 0, 1, \dots, 12)$$

Relevant events:

* $G :=$ the accused is guilty. $I = G^c :=$ the accused is innocent.

G occurs if $Z = 1$

* $C :=$ the accused is convicted. $A = C^c :=$ the accused is acquitted.

C occurs if $\begin{cases} S_{12} \leq 5 & \text{when } Z = 1, \\ S_{12} \geq 7 & \text{when } Z = 0. \end{cases}$

The probability of conviction

G occurs if $Z = 1$

C occurs if $\begin{cases} S_{12} \leq 5 & \text{when } Z = 1, \\ S_{12} \geq 7 & \text{when } Z = 0. \end{cases}$