

# Probability densities in a continuum sample space


---



A generic probability **density** function




A generic probability **density** function






A generic probability **density** function



positivity:  $p(x) \geq 0$



A generic probability **density** function




positivity:  $p(x) \geq 0$

normalisation:  $\int_{-\infty}^{\infty} p(x) dx = 1$



A generic probability **density** function




positivity:  $p(x) \geq 0$

normalisation:  $\int_{-\infty}^{\infty} p(x) dx = 1$

mass **density** per unit length at the point  $x$



A generic probability **density** function



positivity:  $p(x) \geq 0$


normalisation:  $\int_{-\infty}^{\infty} p(x) dx = 1$

mass **density** per unit length at the point  $x$

infinitesimal length  $dx$



A generic probability **density** function



positivity:  $p(x) \geq 0$

normalisation:  $\int_{-\infty}^{\infty} p(x) dx = 1$

mass **density** per unit length at the point  $x$


infinitesimal length  $dx$

probability of infinitesimal interval  $[x, x + dx]$

$$\frac{\text{mass}}{\text{length}} \times \text{length} = \text{mass}$$



A generic probability **density** function



positivity:  $p(x) \geq 0$

normalisation:  $\int_{-\infty}^{\infty} p(x) dx = 1$

mass **density** per unit length at the point  $x$


infinitesimal length  $dx$

probability of infinitesimal interval  $[x, x + dx]$

$$\frac{\text{mass}}{\text{length}} \times \text{length} = \text{mass}$$

a density is any non-negative function with unit area under the curve






$$p(x) \geq 0$$

$$\int_{-\infty}^{\infty} p(x) dx = 1$$






$$p(x) \geq 0$$

$$\int_{-\infty}^{\infty} p(x) dx = 1$$

The continuum sample space:

$$\Omega = \mathbb{R} = (-\infty, \infty)$$





$$p(x) \geq 0$$


$$\int_{-\infty}^{\infty} p(x) dx = 1$$

The continuum sample space:

$$\Omega = \mathbb{R} = (-\infty, \infty)$$

**Convention:** upper case symbols  $R, \dots, Z$  at the end of the alphabet represent the sample points;  
generic sample point (**random variable**)  $\omega = X$ .





$$p(x) \geq 0$$

$$\int_{-\infty}^{\infty} p(x) dx = 1$$

The continuum sample space:


$$\Omega = \mathbb{R} = (-\infty, \infty)$$

**Convention:** upper case symbols  $R, \dots, Z$  at the end of the alphabet represent the sample points;  
generic sample point (**random variable**)  $\omega = X$ .

The basic events are intervals:

$$A = [a, b]$$





$$p(x) \geq 0$$

$$\int_{-\infty}^{\infty} p(x) dx = 1$$

The continuum sample space:


$$\Omega = \mathbb{R} = (-\infty, \infty)$$

**Convention:** upper case symbols  $R, \dots, Z$  at the end of the alphabet represent the sample points;  
generic sample point (**random variable**)  $\omega = X$ .

The basic events are intervals:

$$A = [a, b] =: \{a \leq X \leq b\}$$





$$p(x) \geq 0$$

$$\int_{-\infty}^{\infty} p(x) dx = 1$$

The continuum sample space:

$$\Omega = \mathbb{R} = (-\infty, \infty)$$

**Convention:** upper case symbols  $R, \dots, Z$  at the end of the alphabet represent the sample points;  
generic sample point (**random variable**)  $\omega = X$ .


The basic events are intervals:

$$A = [a, b] =: \{a \leq X \leq b\}$$

Probabilities identified with areas:

$$\mathbf{P}\{a \leq X \leq b\} := \mathbf{P}([a, b])$$





$$p(x) \geq 0$$

$$\int_{-\infty}^{\infty} p(x) dx = 1$$

The continuum sample space:

$$\Omega = \mathbb{R} = (-\infty, \infty)$$

**Convention:** upper case symbols  $R, \dots, Z$  at the end of the alphabet represent the sample points;  
generic sample point (**random variable**)  $\omega = X$ .


The basic events are intervals:

$$A = [a, b] =: \{a \leq X \leq b\}$$

Probabilities identified with areas:

$$\mathbf{P}\{a \leq X \leq b\} := \mathbf{P}([a, b]) = \int_a^b p(x) dx$$





$$p(x) \geq 0$$

$$\int_{-\infty}^{\infty} p(x) dx = 1$$

The continuum sample space:

$$\Omega = \mathbb{R} = (-\infty, \infty)$$

**Convention:** upper case symbols  $R, \dots, Z$  at the end of the alphabet represent the sample points;  
generic sample point (**random variable**)  $\omega = X$ .


Events are subsets of the real line:

$$A = \{X \in A\}$$

Probabilities identified with areas:

$$\mathbf{P}\{X \in A\} := \mathbf{P}(A) = \int_A p(x) dx$$





$$p(x) \geq 0$$

$$\int_{-\infty}^{\infty} p(x) dx = 1$$

The continuum sample space:

$$\Omega = \mathbb{R} = (-\infty, \infty)$$

**Convention:** upper case symbols  $R, \dots, Z$  at the end of the alphabet represent the sample points;  
generic sample point (**random variable**)  $\omega = X$ .

Events are subsets of the real line:

$$A = \{X \in A\}$$

Probabilities identified with areas:

$$\mathbf{P}\{X \in A\} := \mathbf{P}(A) = \int_A p(x) dx$$

$$\frac{\text{mass}}{\text{length}} \times \text{length} = \text{mass}$$