

Additivity once: deconstructing the occurrence of success runs

Additivity once: deconstructing the occurrence of success runs

Key observation: a success run will occur by trial n if, and only if, there is a *first* success run occurring at or before trial n .

Additivity once: deconstructing the occurrence of success runs

Key observation: a success run will occur by trial n if, and only if, there is a *first* success run occurring at or before trial n .

A partition:

$$S_n = F_1 \cup F_2 \cup \cdots \cup F_n$$

Additivity once: deconstructing the occurrence of success runs

Key observation: a success run will occur by trial n if, and only if, there is a *first* success run occurring at or before trial n .

A partition:

$$S_n = F_1 \cup F_2 \cup \cdots \cup F_n$$

Additivity!

$$\mathbf{P}(S_n) = \mathbf{P}(F_1) + \mathbf{P}(F_2) + \cdots + \mathbf{P}(F_n)$$

Additivity once: deconstructing the occurrence of success runs

Key observation: a success run will occur by trial n if, and only if, there is a *first* success run occurring at or before trial n .

A partition:

$$S_n = F_1 \cup F_2 \cup \cdots \cup F_n$$

Additivity!

$$\mathbf{P}(S_n) = \mathbf{P}(F_1) + \mathbf{P}(F_2) + \cdots + \mathbf{P}(F_n)$$

$$s_n = f_1 + f_2 + \cdots + f_n$$

Additivity once: deconstructing the occurrence of success runs

Key observation: a success run will occur by trial n if, and only if, there is a *first* success run occurring at or before trial n .

A partition:

$$S_n = F_1 \cup F_2 \cup \cdots \cup F_n$$

Additivity!

$$\mathbf{P}(S_n) = \mathbf{P}(F_1) + \mathbf{P}(F_2) + \cdots + \mathbf{P}(F_n)$$

$$s_n = f_1 + f_2 + \cdots + f_n$$

Slogan: If you know $\{f_1, f_2, \dots, f_n, \dots\}$ then you know $\{s_1, s_2, \dots, s_n, \dots\}$!