

Typical events: visions of centrality

Random sample (repeated independent trials): $X, X_1, X_2, \dots, X_n, \dots$

$$\mathbf{E}(X) = \mu, \quad \text{Var}(X) = \sigma^2$$

Random sample (repeated independent trials): $X, X_1, X_2, \dots, X_n, \dots$

$$\mathbf{E}(X) = \mu, \quad \text{Var}(X) = \sigma^2$$

The partial sums: $S_n = X_1 + X_2 + \dots + X_n$

Random sample (repeated independent trials): $X, X_1, X_2, \dots, X_n, \dots$
 $\mathbf{E}(X) = \mu, \quad \text{Var}(X) = \sigma^2$

The partial sums: $S_n = X_1 + X_2 + \dots + X_n$

Standardisation to a common centre and scale: $S_n^* = \frac{S_n - n\mu}{\sqrt{n} \sigma}$

Random sample (repeated independent trials): $X, X_1, X_2, \dots, X_n, \dots$
 $\mathbf{E}(X) = \mu, \quad \text{Var}(X) = \sigma^2$

The partial sums: $S_n = X_1 + X_2 + \dots + X_n$

Standardisation to a common centre and scale: $S_n^* = \frac{S_n - n\mu}{\sqrt{n} \sigma}$

The law of large numbers $\frac{S_n}{n}$ is asymptotically concentrated at μ

$$\mathbf{P}\left\{\left|\frac{S_n}{n} - \mu\right| > \epsilon\right\} \rightarrow 0$$

or, even, stronger (Lecture 11.2: n)

$$\mathbf{P}\left\{\left|\frac{S_n}{n} - \mu\right| > \epsilon \text{ i.o.}\right\} = 0$$

infinitely often

Random sample (repeated independent trials): $X, X_1, X_2, \dots, X_n, \dots$

$$\mathbf{E}(X) = \mu, \quad \text{Var}(X) = \sigma^2$$

The partial sums: $S_n = X_1 + X_2 + \dots + X_n$

Standardisation to a common centre and scale: $S_n^* = \frac{S_n - n\mu}{\sqrt{n} \sigma}$

The law of large numbers

$\frac{S_n}{n}$ is asymptotically concentrated at μ

$$\mathbf{P}\left\{\left|\frac{S_n}{n} - \mu\right| > \epsilon\right\} \rightarrow 0$$

or, even, stronger (Lecture 11.2: n)

$$\mathbf{P}\left\{\left|\frac{S_n}{n} - \mu\right| > \epsilon \text{ i.o.}\right\} = 0$$

infinitely often

The central limit theorem

S_n^* is asymptotically normally distributed

$$\mathbf{P}\{a < S_n^* \leq b\} \rightarrow \int_a^b \phi(x) dx \quad (n \rightarrow \infty)$$

