

Is the normal density properly normalised?

Is the normal density properly normalised?

$$\int_{-\infty}^{\infty} \frac{e^{-x^2/2}}{\sqrt{2\pi}} dx$$

Is the normal density properly normalised?

$$\left(\int_{-\infty}^{\infty} \frac{e^{-x^2/2}}{\sqrt{2\pi}} dx \right)^2 = \int_{-\infty}^{\infty} \frac{e^{-x^2/2}}{\sqrt{2\pi}} dx \cdot \int_{-\infty}^{\infty} \frac{e^{-y^2/2}}{\sqrt{2\pi}} dy$$

Is the normal density properly normalised?

$$\begin{aligned}\left(\int_{-\infty}^{\infty} \frac{e^{-x^2/2}}{\sqrt{2\pi}} dx\right)^2 &= \int_{-\infty}^{\infty} \frac{e^{-x^2/2}}{\sqrt{2\pi}} dx \cdot \int_{-\infty}^{\infty} \frac{e^{-y^2/2}}{\sqrt{2\pi}} dy \\ &= \frac{1}{2\pi} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} e^{-(x^2+y^2)/2} dy dx\end{aligned}$$

Is the normal density properly normalised?

$$\begin{aligned} \left(\int_{-\infty}^{\infty} \frac{e^{-x^2/2}}{\sqrt{2\pi}} dx \right)^2 &= \int_{-\infty}^{\infty} \frac{e^{-x^2/2}}{\sqrt{2\pi}} dx \cdot \int_{-\infty}^{\infty} \frac{e^{-y^2/2}}{\sqrt{2\pi}} dy \\ &= \frac{1}{2\pi} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} e^{-(x^2+y^2)/2} dy dx \end{aligned}$$

rectangular-to-polar transformation
 $x = r \cos \theta, \quad y = r \sin \theta, \quad dy dx = r d\theta dr$

Is the normal density properly normalised?

$$\begin{aligned} \left(\int_{-\infty}^{\infty} \frac{e^{-x^2/2}}{\sqrt{2\pi}} dx \right)^2 &= \int_{-\infty}^{\infty} \frac{e^{-x^2/2}}{\sqrt{2\pi}} dx \cdot \int_{-\infty}^{\infty} \frac{e^{-y^2/2}}{\sqrt{2\pi}} dy \\ &= \frac{1}{2\pi} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} e^{-(x^2+y^2)/2} dy dx \\ &= \frac{1}{2\pi} \int_0^{\infty} \int_0^{2\pi} e^{-r^2/2} r d\theta dr \end{aligned}$$

rectangular-to-polar transformation
 $x = r \cos \theta, \quad y = r \sin \theta, \quad dy dx = r d\theta dr$

Is the normal density properly normalised?

$$\begin{aligned} \left(\int_{-\infty}^{\infty} \frac{e^{-x^2/2}}{\sqrt{2\pi}} dx \right)^2 &= \int_{-\infty}^{\infty} \frac{e^{-x^2/2}}{\sqrt{2\pi}} dx \cdot \int_{-\infty}^{\infty} \frac{e^{-y^2/2}}{\sqrt{2\pi}} dy \\ &= \frac{1}{2\pi} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} e^{-(x^2+y^2)/2} dy dx \\ &= \frac{1}{2\pi} \int_0^{\infty} \int_0^{2\pi} e^{-r^2/2} r d\theta dr \\ &= \frac{1}{2\pi} \int_0^{\infty} e^{-r^2/2} r \int_0^{2\pi} d\theta dr \end{aligned}$$

rectangular-to-polar transformation
 $x = r \cos \theta, \quad y = r \sin \theta, \quad dy dx = r d\theta dr$

Is the normal density properly normalised?

$$\begin{aligned}\left(\int_{-\infty}^{\infty} \frac{e^{-x^2/2}}{\sqrt{2\pi}} dx\right)^2 &= \int_{-\infty}^{\infty} \frac{e^{-x^2/2}}{\sqrt{2\pi}} dx \cdot \int_{-\infty}^{\infty} \frac{e^{-y^2/2}}{\sqrt{2\pi}} dy \\&= \frac{1}{2\pi} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} e^{-(x^2+y^2)/2} dy dx \\&= \frac{1}{2\pi} \int_0^{\infty} \int_0^{2\pi} e^{-r^2/2} r d\theta dr \\&= \frac{1}{2\pi} \int_0^{\infty} e^{-r^2/2} r \int_0^{2\pi} d\theta dr \\&= \frac{1}{2\pi} \int_0^{\infty} e^{-r^2/2} r dr \cdot 2\pi\end{aligned}$$

rectangular-to-polar transformation
 $x = r \cos \theta, \quad y = r \sin \theta, \quad dy dx = r d\theta dr$

Is the normal density properly normalised?

$$\begin{aligned}\left(\int_{-\infty}^{\infty} \frac{e^{-x^2/2}}{\sqrt{2\pi}} dx\right)^2 &= \int_{-\infty}^{\infty} \frac{e^{-x^2/2}}{\sqrt{2\pi}} dx \cdot \int_{-\infty}^{\infty} \frac{e^{-y^2/2}}{\sqrt{2\pi}} dy \\&= \frac{1}{2\pi} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} e^{-(x^2+y^2)/2} dy dx \\&= \frac{1}{2\pi} \int_0^{\infty} \int_0^{2\pi} e^{-r^2/2} r d\theta dr \\&= \frac{1}{2\pi} \int_0^{\infty} e^{-r^2/2} r \int_0^{2\pi} d\theta dr \\&= \frac{1}{2\pi} \int_0^{\infty} e^{-r^2/2} r dr \cdot 2\pi \\&= \int_0^{\infty} e^{-r^2/2} r dr\end{aligned}$$

rectangular-to-polar transformation
 $x = r \cos \theta, \quad y = r \sin \theta, \quad dy dx = r d\theta dr$

Is the normal density properly normalised?

$$\begin{aligned}\left(\int_{-\infty}^{\infty} \frac{e^{-x^2/2}}{\sqrt{2\pi}} dx\right)^2 &= \int_{-\infty}^{\infty} \frac{e^{-x^2/2}}{\sqrt{2\pi}} dx \cdot \int_{-\infty}^{\infty} \frac{e^{-y^2/2}}{\sqrt{2\pi}} dy \\&= \frac{1}{2\pi} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} e^{-(x^2+y^2)/2} dy dx \\&= \frac{1}{2\pi} \int_0^{\infty} \int_0^{2\pi} e^{-r^2/2} r d\theta dr \\&= \frac{1}{2\pi} \int_0^{\infty} e^{-r^2/2} r \int_0^{2\pi} d\theta dr \\&= \frac{1}{2\pi} \int_0^{\infty} e^{-r^2/2} r dr \cdot 2\pi \\&= \int_0^{\infty} e^{-r^2/2} r dr\end{aligned}$$

rectangular-to-polar transformation
 $x = r \cos \theta, \quad y = r \sin \theta, \quad dy dx = r d\theta dr$

change of variable
 $t = r^2/2, \quad dt = r dr$

Is the normal density properly normalised?

$$\begin{aligned}\left(\int_{-\infty}^{\infty} \frac{e^{-x^2/2}}{\sqrt{2\pi}} dx\right)^2 &= \int_{-\infty}^{\infty} \frac{e^{-x^2/2}}{\sqrt{2\pi}} dx \cdot \int_{-\infty}^{\infty} \frac{e^{-y^2/2}}{\sqrt{2\pi}} dy \\&= \frac{1}{2\pi} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} e^{-(x^2+y^2)/2} dy dx \\&= \frac{1}{2\pi} \int_0^{\infty} \int_0^{2\pi} e^{-r^2/2} r d\theta dr \\&= \frac{1}{2\pi} \int_0^{\infty} e^{-r^2/2} r \int_0^{2\pi} d\theta dr \\&= \frac{1}{2\pi} \int_0^{\infty} e^{-r^2/2} r dr \cdot 2\pi \\&= \int_0^{\infty} e^{-r^2/2} r dr \\&= \int_0^{\infty} e^{-t} dt\end{aligned}$$

rectangular-to-polar transformation
 $x = r \cos \theta, \quad y = r \sin \theta, \quad dy dx = r d\theta dr$

change of variable
 $t = r^2/2, \quad dt = r dr$

Is the normal density properly normalised?

$$\begin{aligned} \left(\int_{-\infty}^{\infty} \frac{e^{-x^2/2}}{\sqrt{2\pi}} dx \right)^2 &= \int_{-\infty}^{\infty} \frac{e^{-x^2/2}}{\sqrt{2\pi}} dx \cdot \int_{-\infty}^{\infty} \frac{e^{-y^2/2}}{\sqrt{2\pi}} dy \\ &= \frac{1}{2\pi} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} e^{-(x^2+y^2)/2} dy dx \\ &= \frac{1}{2\pi} \int_0^{\infty} \int_0^{2\pi} e^{-r^2/2} r d\theta dr \\ &= \frac{1}{2\pi} \int_0^{\infty} e^{-r^2/2} r \int_0^{2\pi} d\theta dr \\ &= \frac{1}{2\pi} \int_0^{\infty} e^{-r^2/2} r dr \cdot 2\pi \\ &= \int_0^{\infty} e^{-r^2/2} r dr \\ &= \int_0^{\infty} e^{-t} dt \\ &= -e^{-t} \Big|_0^{\infty} = -0 + 1 = 1 \end{aligned}$$

rectangular-to-polar transformation
 $x = r \cos \theta, \quad y = r \sin \theta, \quad dy dx = r d\theta dr$

change of variable
 $t = r^2/2, \quad dt = r dr$