

Product space and measure

Product space and measure

- ❖ *Individual chance experiments, independent trials:*

Product space and measure

- ❖ *Individual chance experiments, independent trials:*
 - * Trial 1: $\mathfrak{A} = \{\mathfrak{a}_j, j \geq 1\}$. Atomic mass function $\{\mathfrak{a}_j\} \mapsto \mathfrak{p}_j$.

Product space and measure

- ❖ *Individual chance experiments, independent trials:*
 - * Trial 1: $\mathfrak{A} = \{\mathfrak{a}_j, j \geq 1\}$. Atomic mass function $\{\mathfrak{a}_j\} \mapsto \mathfrak{p}_j$.
 - * Trial 2: $\mathfrak{B} = \{\mathfrak{b}_k, k \geq 1\}$. Atomic mass function $\{\mathfrak{b}_k\} \mapsto \mathfrak{q}_k$.

Product space and measure

- ❖ *Individual chance experiments, independent trials:*
 - * Trial 1: $\mathfrak{A} = \{\mathfrak{a}_j, j \geq 1\}$. Atomic mass function $\{\mathfrak{a}_j\} \mapsto \mathfrak{p}_j$.
 - * Trial 2: $\mathfrak{B} = \{\mathfrak{b}_k, k \geq 1\}$. Atomic mass function $\{\mathfrak{b}_k\} \mapsto \mathfrak{q}_k$.
- ❖ *Compound chance experiment, product space and measure:*

Product space and measure

- ❖ *Individual chance experiments, independent trials:*
 - * Trial 1: $\mathfrak{A} = \{\mathfrak{a}_j, j \geq 1\}$. Atomic mass function $\{\mathfrak{a}_j\} \mapsto \mathfrak{p}_j$.
 - * Trial 2: $\mathfrak{B} = \{\mathfrak{b}_k, k \geq 1\}$. Atomic mass function $\{\mathfrak{b}_k\} \mapsto \mathfrak{q}_k$.
- ❖ *Compound chance experiment, product space and measure:*
 - * $\Omega = \mathfrak{A} \times \mathfrak{B} = \{(\mathfrak{a}_j, \mathfrak{b}_k): j \geq 1, k \geq 1\}$. Atomic measure $\mathbf{P}\{(\mathfrak{a}_j, \mathfrak{b}_k)\} := \mathfrak{p}_j \mathfrak{q}_k$.

Product space and measure

- ❖ *Individual chance experiments, independent trials:*
 - * Trial 1: $\mathfrak{A} = \{\mathfrak{a}_j, j \geq 1\}$. Atomic mass function $\{\mathfrak{a}_j\} \mapsto \mathfrak{p}_j$.
 - * Trial 2: $\mathfrak{B} = \{\mathfrak{b}_k, k \geq 1\}$. Atomic mass function $\{\mathfrak{b}_k\} \mapsto \mathfrak{q}_k$.
- ❖ *Compound chance experiment, product space and measure:*
 - * $\Omega = \mathfrak{A} \times \mathfrak{B} = \{(\mathfrak{a}_j, \mathfrak{b}_k): j \geq 1, k \geq 1\}$. Atomic measure $\mathbf{P}\{(\mathfrak{a}_j, \mathfrak{b}_k)\} := \mathfrak{p}_j \mathfrak{q}_k$.
- ❖ Suppose that \mathbb{J} and \mathbb{K} are any two subsets of indices. Then:

Product space and measure

- ❖ *Individual chance experiments, independent trials:*
 - * Trial 1: $\mathfrak{A} = \{a_j, j \geq 1\}$. Atomic mass function $\{a_j\} \mapsto p_j$.
 - * Trial 2: $\mathfrak{B} = \{b_k, k \geq 1\}$. Atomic mass function $\{b_k\} \mapsto q_k$.
- ❖ *Compound chance experiment, product space and measure:*
 - * $\Omega = \mathfrak{A} \times \mathfrak{B} = \{(a_j, b_k): j \geq 1, k \geq 1\}$. Atomic measure $P\{(a_j, b_k)\} := p_j q_k$.
- ❖ Suppose that \mathbb{J} and \mathbb{K} are any two subsets of indices. Then:
 - * The event $A := \{(a_j, b_k): j \in \mathbb{J}, k \geq 1\}$ is completely determined by the subset $\mathfrak{S} = \{a_j: j \in \mathbb{J}\}$ of \mathfrak{A} ;


Product space and measure

- ❖ *Individual chance experiments, independent trials:*
 - * Trial 1: $\mathfrak{A} = \{\mathfrak{a}_j, j \geq 1\}$. Atomic mass function $\{\mathfrak{a}_j\} \mapsto \mathfrak{p}_j$.
 - * Trial 2: $\mathfrak{B} = \{\mathfrak{b}_k, k \geq 1\}$. Atomic mass function $\{\mathfrak{b}_k\} \mapsto \mathfrak{q}_k$.
- ❖ *Compound chance experiment, product space and measure:*
 - * $\Omega = \mathfrak{A} \times \mathfrak{B} = \{(\mathfrak{a}_j, \mathfrak{b}_k): j \geq 1, k \geq 1\}$. Atomic measure $\mathbf{P}\{(\mathfrak{a}_j, \mathfrak{b}_k)\} := \mathfrak{p}_j \mathfrak{q}_k$.
- ❖ Suppose that \mathbb{J} and \mathbb{K} are any two subsets of indices. Then:
 - * The event $A := \{(\mathfrak{a}_j, \mathfrak{b}_k): j \in \mathbb{J}, k \geq 1\}$ is completely determined by the subset $\mathfrak{S} = \{\mathfrak{a}_j: j \in \mathbb{J}\}$ of \mathfrak{A} ;
 - * The event $B := \{(\mathfrak{a}_j, \mathfrak{b}_k): j \geq 1, k \in \mathbb{K}\}$ is completely determined by the subset $\mathfrak{T} = \{\mathfrak{b}_k: k \in \mathbb{K}\}$ of \mathfrak{B} .

Product space and measure


- ❖ *Individual chance experiments, independent trials:*
 - * Trial 1: $\mathfrak{A} = \{\mathfrak{a}_j, j \geq 1\}$. Atomic mass function $\{\mathfrak{a}_j\} \mapsto \mathfrak{p}_j$.
 - * Trial 2: $\mathfrak{B} = \{\mathfrak{b}_k, k \geq 1\}$. Atomic mass function $\{\mathfrak{b}_k\} \mapsto \mathfrak{q}_k$.
- ❖ *Compound chance experiment, product space and measure:*
 - * $\Omega = \mathfrak{A} \times \mathfrak{B} = \{(\mathfrak{a}_j, \mathfrak{b}_k): j \geq 1, k \geq 1\}$. Atomic measure $\mathbf{P}\{(\mathfrak{a}_j, \mathfrak{b}_k)\} := \mathfrak{p}_j \mathfrak{q}_k$.
- ❖ Suppose that \mathbb{J} and \mathbb{K} are any two subsets of indices. Then:
 - * The event $A := \{(\mathfrak{a}_j, \mathfrak{b}_k): j \in \mathbb{J}, k \geq 1\}$ is completely determined by the subset $\mathfrak{S} = \{\mathfrak{a}_j: j \in \mathbb{J}\}$ of \mathfrak{A} ;
 - * The event $B := \{(\mathfrak{a}_j, \mathfrak{b}_k): j \geq 1, k \in \mathbb{K}\}$ is completely determined by the subset $\mathfrak{T} = \{\mathfrak{b}_k: k \in \mathbb{K}\}$ of \mathfrak{B} .
 - * “Rectangular” intersection: $A \cap B = \{(\mathfrak{a}_j, \mathfrak{b}_k): j \in \mathbb{J}, k \in \mathbb{K}\} = \mathfrak{S} \times \mathfrak{T}$.

Product space and measure



- ❖ *Individual chance experiments, independent trials:*
 - * Trial 1: $\mathfrak{A} = \{\mathfrak{a}_j, j \geq 1\}$. Atomic mass function $\{\mathfrak{a}_j\} \mapsto \mathfrak{p}_j$.
 - * Trial 2: $\mathfrak{B} = \{\mathfrak{b}_k, k \geq 1\}$. Atomic mass function $\{\mathfrak{b}_k\} \mapsto \mathfrak{q}_k$.
- ❖ *Compound chance experiment, product space and measure:*
 - * $\Omega = \mathfrak{A} \times \mathfrak{B} = \{(\mathfrak{a}_j, \mathfrak{b}_k): j \geq 1, k \geq 1\}$. Atomic measure $\mathbf{P}\{(\mathfrak{a}_j, \mathfrak{b}_k)\} := \mathfrak{p}_j \mathfrak{q}_k$.
- ❖ Suppose that \mathbb{J} and \mathbb{K} are any two subsets of indices. Then:
 - * The event $A := \{(\mathfrak{a}_j, \mathfrak{b}_k): j \in \mathbb{J}, k \geq 1\}$ is completely determined by the subset $\mathfrak{S} = \{\mathfrak{a}_j: j \in \mathbb{J}\}$ of \mathfrak{A} ;
 - * The event $B := \{(\mathfrak{a}_j, \mathfrak{b}_k): j \geq 1, k \in \mathbb{K}\}$ is completely determined by the subset $\mathfrak{T} = \{\mathfrak{b}_k: k \in \mathbb{K}\}$ of \mathfrak{B} .
 - * “Rectangular” intersection: $A \cap B = \{(\mathfrak{a}_j, \mathfrak{b}_k): j \in \mathbb{J}, k \in \mathbb{K}\} = \mathfrak{S} \times \mathfrak{T}$.

Product space and measure



❖ *Individual chance experiments, independent trials:*

- * Trial 1: $\mathfrak{A} = \{\mathfrak{a}_j, j \geq 1\}$. Atomic mass function $\{\mathfrak{a}_j\} \mapsto \mathfrak{p}_j$.
- * Trial 2: $\mathfrak{B} = \{\mathfrak{b}_k, k \geq 1\}$. Atomic mass function $\{\mathfrak{b}_k\} \mapsto \mathfrak{q}_k$.

$$\mathfrak{p}_j \geq 0, \quad \sum_{j \geq 1} \mathfrak{p}_j = 1$$

$$\mathfrak{q}_k \geq 0, \quad \sum_{k \geq 1} \mathfrak{q}_k = 1$$


❖ *Compound chance experiment, product space and measure:*

- * $\Omega = \mathfrak{A} \times \mathfrak{B} = \{(\mathfrak{a}_j, \mathfrak{b}_k): j \geq 1, k \geq 1\}$. Atomic measure $\mathbf{P}\{(\mathfrak{a}_j, \mathfrak{b}_k)\} := \mathfrak{p}_j \mathfrak{q}_k$.

❖ Suppose that \mathbb{J} and \mathbb{K} are any two subsets of indices. Then:

- * The event $A := \{(\mathfrak{a}_j, \mathfrak{b}_k): j \in \mathbb{J}, k \geq 1\}$ is completely determined by the subset $\mathfrak{S} = \{\mathfrak{a}_j: j \in \mathbb{J}\}$ of \mathfrak{A} ;
- * The event $B := \{(\mathfrak{a}_j, \mathfrak{b}_k): j \geq 1, k \in \mathbb{K}\}$ is completely determined by the subset $\mathfrak{T} = \{\mathfrak{b}_k: k \in \mathbb{K}\}$ of \mathfrak{B} .
- * “Rectangular” intersection: $A \cap B = \{(\mathfrak{a}_j, \mathfrak{b}_k): j \in \mathbb{J}, k \in \mathbb{K}\} = \mathfrak{S} \times \mathfrak{T}$.

Product space and measure



❖ *Individual chance experiments, independent trials:*

- * Trial 1: $\mathfrak{A} = \{\mathfrak{a}_j, j \geq 1\}$. Atomic mass function $\{\mathfrak{a}_j\} \mapsto \mathfrak{p}_j$.
- * Trial 2: $\mathfrak{B} = \{\mathfrak{b}_k, k \geq 1\}$. Atomic mass function $\{\mathfrak{b}_k\} \mapsto \mathfrak{q}_k$.

$$\mathfrak{p}_j \geq 0, \quad \sum_{j \geq 1} \mathfrak{p}_j = 1$$

$$\mathfrak{q}_k \geq 0, \quad \sum_{k \geq 1} \mathfrak{q}_k = 1$$

❖ *Compound chance experiment, product space and measure:*


Is the product measure suitably normalised?

- * $\Omega = \mathfrak{A} \times \mathfrak{B} = \{(\mathfrak{a}_j, \mathfrak{b}_k): j \geq 1, k \geq 1\}$. Atomic measure $\mathbf{P}\{(\mathfrak{a}_j, \mathfrak{b}_k)\} := \mathfrak{p}_j \mathfrak{q}_k$.

❖ Suppose that \mathbb{J} and \mathbb{K} are any two subsets of indices. Then:

- * The event $A := \{(\mathfrak{a}_j, \mathfrak{b}_k): j \in \mathbb{J}, k \geq 1\}$ is completely determined by the subset $\mathfrak{S} = \{\mathfrak{a}_j: j \in \mathbb{J}\}$ of \mathfrak{A} ;
- * The event $B := \{(\mathfrak{a}_j, \mathfrak{b}_k): j \geq 1, k \in \mathbb{K}\}$ is completely determined by the subset $\mathfrak{T} = \{\mathfrak{b}_k: k \in \mathbb{K}\}$ of \mathfrak{B} .
- * “Rectangular” intersection: $A \cap B = \{(\mathfrak{a}_j, \mathfrak{b}_k): j \in \mathbb{J}, k \in \mathbb{K}\} = \mathfrak{S} \times \mathfrak{T}$.

Product space and measure



❖ *Individual chance experiments, independent trials:*

- * Trial 1: $\mathfrak{A} = \{\mathfrak{a}_j, j \geq 1\}$. Atomic mass function $\{\mathfrak{a}_j\} \mapsto p_j$.
- * Trial 2: $\mathfrak{B} = \{\mathfrak{b}_k, k \geq 1\}$. Atomic mass function $\{\mathfrak{b}_k\} \mapsto q_k$.

$$p_j \geq 0, \quad \sum_{j \geq 1} p_j = 1$$

$$q_k \geq 0, \quad \sum_{k \geq 1} q_k = 1$$

❖ *Compound chance experiment, product space and measure:*

- * $\Omega = \mathfrak{A} \times \mathfrak{B} = \{(\mathfrak{a}_j, \mathfrak{b}_k): j \geq 1, k \geq 1\}$. Atomic measure $P\{(\mathfrak{a}_j, \mathfrak{b}_k)\} := p_j q_k$.


Is the product measure suitably normalised?

$$\sum_{j \geq 1} \sum_{k \geq 1} p_j q_k = \left(\sum_{j \geq 1} p_j \right) \left(\sum_{k \geq 1} q_k \right) = 1$$

❖ Suppose that \mathbb{J} and \mathbb{K} are any two subsets of indices. Then:

- * The event $A := \{(\mathfrak{a}_j, \mathfrak{b}_k): j \in \mathbb{J}, k \geq 1\}$ is completely determined by the subset $\mathfrak{S} = \{\mathfrak{a}_j: j \in \mathbb{J}\}$ of \mathfrak{A} ;
- * The event $B := \{(\mathfrak{a}_j, \mathfrak{b}_k): j \geq 1, k \in \mathbb{K}\}$ is completely determined by the subset $\mathfrak{T} = \{\mathfrak{b}_k: k \in \mathbb{K}\}$ of \mathfrak{B} .
- * “Rectangular” intersection: $A \cap B = \{(\mathfrak{a}_j, \mathfrak{b}_k): j \in \mathbb{J}, k \in \mathbb{K}\} = \mathfrak{S} \times \mathfrak{T}$.

Product space and measure



❖ *Individual chance experiments, independent trials:*

- * Trial 1: $\mathfrak{A} = \{\mathfrak{a}_j, j \geq 1\}$. Atomic mass function $\{\mathfrak{a}_j\} \mapsto p_j$.
- * Trial 2: $\mathfrak{B} = \{\mathfrak{b}_k, k \geq 1\}$. Atomic mass function $\{\mathfrak{b}_k\} \mapsto q_k$.

$$p_j \geq 0, \quad \sum_{j \geq 1} p_j = 1$$

$$q_k \geq 0, \quad \sum_{k \geq 1} q_k = 1$$

❖ *Compound chance experiment, product space and measure:*

- * $\Omega = \mathfrak{A} \times \mathfrak{B} = \{(\mathfrak{a}_j, \mathfrak{b}_k): j \geq 1, k \geq 1\}$. Atomic measure $\mathbf{P}\{(\mathfrak{a}_j, \mathfrak{b}_k)\} := p_j q_k$.

Is the product measure suitably normalised?


$$\sum_{j \geq 1} \sum_{k \geq 1} p_j q_k = \left(\sum_{j \geq 1} p_j \right) \left(\sum_{k \geq 1} q_k \right) = 1$$

❖ Suppose that \mathbb{J} and \mathbb{K} are any two subsets of indices. Then:

- * The event $A := \{(\mathfrak{a}_j, \mathfrak{b}_k): j \in \mathbb{J}, k \geq 1\}$ is completely determined by the subset $\mathfrak{S} = \{\mathfrak{a}_j: j \in \mathbb{J}\}$ of \mathfrak{A} ;
- * The event $B := \{(\mathfrak{a}_j, \mathfrak{b}_k): j \geq 1, k \in \mathbb{K}\}$ is completely determined by the subset $\mathfrak{T} = \{\mathfrak{b}_k: k \in \mathbb{K}\}$ of \mathfrak{B} .
- * “Rectangular” intersection: $A \cap B = \{(\mathfrak{a}_j, \mathfrak{b}_k): j \in \mathbb{J}, k \in \mathbb{K}\} = \mathfrak{S} \times \mathfrak{T}$.

$$\mathbf{P}(A) = \sum_{j \in \mathbb{J}} \sum_{k \geq 1} p_j q_k = \left(\sum_{j \in \mathbb{J}} p_j \right) \left(\sum_{k \geq 1} q_k \right) = \sum_{j \in \mathbb{J}} p_j$$

Product space and measure



❖ *Individual chance experiments, independent trials:*

- * Trial 1: $\mathfrak{A} = \{\mathfrak{a}_j, j \geq 1\}$. Atomic mass function $\{\mathfrak{a}_j\} \mapsto p_j$.
- * Trial 2: $\mathfrak{B} = \{\mathfrak{b}_k, k \geq 1\}$. Atomic mass function $\{\mathfrak{b}_k\} \mapsto q_k$.

$$p_j \geq 0, \quad \sum_{j \geq 1} p_j = 1$$

$$q_k \geq 0, \quad \sum_{k \geq 1} q_k = 1$$

❖ *Compound chance experiment, product space and measure:*

- * $\Omega = \mathfrak{A} \times \mathfrak{B} = \{(\mathfrak{a}_j, \mathfrak{b}_k): j \geq 1, k \geq 1\}$. Atomic measure $\mathbf{P}\{(\mathfrak{a}_j, \mathfrak{b}_k)\} := p_j q_k$.

Is the product measure suitably normalised?

$$\sum_{j \geq 1} \sum_{k \geq 1} p_j q_k = \left(\sum_{j \geq 1} p_j \right) \left(\sum_{k \geq 1} q_k \right) = 1$$


❖ Suppose that \mathbb{J} and \mathbb{K} are any two subsets of indices. Then:

- * The event $A := \{(\mathfrak{a}_j, \mathfrak{b}_k): j \in \mathbb{J}, k \geq 1\}$ is completely determined by the subset $\mathfrak{G} = \{\mathfrak{a}_j: j \in \mathbb{J}\}$ of \mathfrak{A} ;
- * The event $B := \{(\mathfrak{a}_j, \mathfrak{b}_k): j \geq 1, k \in \mathbb{K}\}$ is completely determined by the subset $\mathfrak{T} = \{\mathfrak{b}_k: k \in \mathbb{K}\}$ of \mathfrak{B} .
- * “Rectangular” intersection: $A \cap B = \{(\mathfrak{a}_j, \mathfrak{b}_k): j \in \mathbb{J}, k \in \mathbb{K}\} = \mathfrak{G} \times \mathfrak{T}$.

$$\mathbf{P}(A) = \sum_{j \in \mathbb{J}} \sum_{k \geq 1} p_j q_k = \left(\sum_{j \in \mathbb{J}} p_j \right) \left(\sum_{k \geq 1} q_k \right) = \sum_{j \in \mathbb{J}} p_j$$

$$\mathbf{P}(B) = \sum_{j \geq 1} \sum_{k \in \mathbb{K}} p_j q_k = \left(\sum_{j \geq 1} p_j \right) \left(\sum_{k \in \mathbb{K}} q_k \right) = \sum_{k \in \mathbb{K}} q_k$$

Product space and measure



❖ *Individual chance experiments, independent trials:*

- * Trial 1: $\mathfrak{A} = \{\mathfrak{a}_j, j \geq 1\}$. Atomic mass function $\{\mathfrak{a}_j\} \mapsto p_j$.
- * Trial 2: $\mathfrak{B} = \{\mathfrak{b}_k, k \geq 1\}$. Atomic mass function $\{\mathfrak{b}_k\} \mapsto q_k$.

$$p_j \geq 0, \quad \sum_{j \geq 1} p_j = 1$$

$$q_k \geq 0, \quad \sum_{k \geq 1} q_k = 1$$

❖ *Compound chance experiment, product space and measure:*

- * $\Omega = \mathfrak{A} \times \mathfrak{B} = \{(\mathfrak{a}_j, \mathfrak{b}_k): j \geq 1, k \geq 1\}$. Atomic measure $\mathbf{P}\{(\mathfrak{a}_j, \mathfrak{b}_k)\} := p_j q_k$.

Is the product measure suitably normalised?

$$\sum_{j \geq 1} \sum_{k \geq 1} p_j q_k = \left(\sum_{j \geq 1} p_j \right) \left(\sum_{k \geq 1} q_k \right) = 1$$

❖ Suppose that \mathbb{J} and \mathbb{K} are any two subsets of indices. Then:


- * The event $A := \{(\mathfrak{a}_j, \mathfrak{b}_k): j \in \mathbb{J}, k \geq 1\}$ is completely determined by the subset $\mathfrak{S} = \{\mathfrak{a}_j: j \in \mathbb{J}\}$ of \mathfrak{A} ;
- * The event $B := \{(\mathfrak{a}_j, \mathfrak{b}_k): j \geq 1, k \in \mathbb{K}\}$ is completely determined by the subset $\mathfrak{T} = \{\mathfrak{b}_k: k \in \mathbb{K}\}$ of \mathfrak{B} .
- * “Rectangular” intersection: $A \cap B = \{(\mathfrak{a}_j, \mathfrak{b}_k): j \in \mathbb{J}, k \in \mathbb{K}\} = \mathfrak{S} \times \mathfrak{T}$.

$$\mathbf{P}(A) = \sum_{j \in \mathbb{J}} \sum_{k \geq 1} p_j q_k = \left(\sum_{j \in \mathbb{J}} p_j \right) \left(\sum_{k \geq 1} q_k \right) = \sum_{j \in \mathbb{J}} p_j$$

$$\mathbf{P}(B) = \sum_{j \geq 1} \sum_{k \in \mathbb{K}} p_j q_k = \left(\sum_{j \geq 1} p_j \right) \left(\sum_{k \in \mathbb{K}} q_k \right) = \sum_{k \in \mathbb{K}} q_k$$

$$\mathbf{P}(A \cap B) = \sum_{j \in \mathbb{J}} \sum_{k \in \mathbb{K}} p_j q_k = \left(\sum_{j \in \mathbb{J}} p_j \right) \left(\sum_{k \in \mathbb{K}} q_k \right) = \mathbf{P}(A) \mathbf{P}(B)$$

Product space and measure



❖ *Individual chance experiments, independent trials:*

- * Trial 1: $\mathfrak{A} = \{\mathfrak{a}_j, j \geq 1\}$. Atomic mass function $\{\mathfrak{a}_j\} \mapsto p_j$.
- * Trial 2: $\mathfrak{B} = \{\mathfrak{b}_k, k \geq 1\}$. Atomic mass function $\{\mathfrak{b}_k\} \mapsto q_k$.

$$p_j \geq 0, \quad \sum_{j \geq 1} p_j = 1$$

$$q_k \geq 0, \quad \sum_{k \geq 1} q_k = 1$$

❖ *Compound chance experiment, product space and measure:*

- * $\Omega = \mathfrak{A} \times \mathfrak{B} = \{(\mathfrak{a}_j, \mathfrak{b}_k): j \geq 1, k \geq 1\}$. Atomic measure $\mathbf{P}\{(\mathfrak{a}_j, \mathfrak{b}_k)\} := p_j q_k$.

Is the product measure suitably normalised?

$$\sum_{j \geq 1} \sum_{k \geq 1} p_j q_k = \left(\sum_{j \geq 1} p_j \right) \left(\sum_{k \geq 1} q_k \right) = 1$$

❖ Suppose that \mathbb{J} and \mathbb{K} are any two subsets of indices. Then:

- * The event $A := \{(\mathfrak{a}_j, \mathfrak{b}_k): j \in \mathbb{J}, k \geq 1\}$ is completely determined by the subset $\mathfrak{S} = \{\mathfrak{a}_j: j \in \mathbb{J}\}$ of \mathfrak{A} ;
- * The event $B := \{(\mathfrak{a}_j, \mathfrak{b}_k): j \geq 1, k \in \mathbb{K}\}$ is completely determined by the subset $\mathfrak{T} = \{\mathfrak{b}_k: k \in \mathbb{K}\}$ of \mathfrak{B} .
- * “Rectangular” intersection: $A \cap B = \{(\mathfrak{a}_j, \mathfrak{b}_k): j \in \mathbb{J}, k \in \mathbb{K}\} = \mathfrak{S} \times \mathfrak{T}$.

$$\mathbf{P}(A) = \sum_{j \in \mathbb{J}} \sum_{k \geq 1} p_j q_k = \left(\sum_{j \in \mathbb{J}} p_j \right) \left(\sum_{k \geq 1} q_k \right) = \sum_{j \in \mathbb{J}} p_j$$

$$\mathbf{P}(B) = \sum_{j \geq 1} \sum_{k \in \mathbb{K}} p_j q_k = \left(\sum_{j \geq 1} p_j \right) \left(\sum_{k \in \mathbb{K}} q_k \right) = \sum_{k \in \mathbb{K}} q_k$$

$$\mathbf{P}(A \cap B) = \sum_{j \in \mathbb{J}} \sum_{k \in \mathbb{K}} p_j q_k = \left(\sum_{j \in \mathbb{J}} p_j \right) \left(\sum_{k \in \mathbb{K}} q_k \right) = \mathbf{P}(A) \mathbf{P}(B)$$

Any such events A and B determined by separate trials are independent.