

One curve to rule them all

Random sample (repeated independent trials):

$$X, X_1, X_2, \dots, X_n, \dots$$

$$\mathbb{E}(X) = \mu, \quad \text{Var}(X) = \sigma^2$$

Random sample (repeated independent trials):

$$X, X_1, X_2, \dots, X_n, \dots$$

$$\mathbf{E}(X) = \mu, \quad \text{Var}(X) = \sigma^2$$

The partial sums: $S_n = X_1 + X_2 + \dots + X_n$

Random sample (repeated independent trials):

$$X, X_1, X_2, \dots, X_n, \dots$$

$$\mathbf{E}(X) = \mu, \quad \text{Var}(X) = \sigma^2$$

The partial sums: $S_n = X_1 + X_2 + \dots + X_n$

Standardisation to a common centre and scale:

$$S_n^* = \frac{S_n - n\mu}{\sqrt{n} \sigma}$$

Random sample (repeated independent trials):

$$X, X_1, X_2, \dots, X_n, \dots$$

$$\mathbf{E}(X) = \mu, \quad \text{Var}(X) = \sigma^2$$


The partial sums: $S_n = X_1 + X_2 + \dots + X_n$

Standardisation to a common centre and scale:

$$S_n^* = \frac{S_n - n\mu}{\sqrt{n} \sigma}$$

The central limit theorem

$$\mathbf{P}\{a < S_n^* \leq b\} \rightarrow \int_a^b \phi(x) dx \quad (n \rightarrow \infty)$$



Slogan

Sums of independent perturbations follow a normal law.