

Introduction to Week Five

Initial Value Problems

- ✓

Video: Euler Method | Lecture 48
7 min
- ✓

Reading: When the Euler Method is Exact
10 min
- ▶

Video: Modified Euler Method | Lecture 49
9 min
- Ⓜ

Reading: When the Modified Euler Method is Exact
10 min
- ▶

Video: Runge-Kutta Methods | Lecture 50
12 min
- ▶

Video: Second-Order Runge-Kutta Methods | Lecture 51
7 min
- Ⓜ

Reading: Ralston's Method
5 min
- Ⓜ

Reading: Runge-Kutta Methods and Quadrature Formulas
10 min
- ▶

Video: Higher-Order Runge-Kutta Methods | Lecture 52
10 min
- Ⓜ

Reading: Fourth-Order Runge-Kutta Method and Simpson's Rule
10 min

Systems of Differential Equations

Initial Value Problems in MATLAB

Boundary Value Problems

Quiz

Programming Assignment: The Two-Body Problem

When the Euler Method is Exact

Let $\dot{x} = b$, with initial condition $x(0) = x_0$ and b a constant. With $t = n\Delta t$, show that the Euler method results in the exact solution

$$x(t) = x_0 + bt.$$

✓ Completed

Go to next item

👍 Like

🗨 Dislike

🚩 Report an issue

