

MOOC Econometrics

Lecture 5.1 on Binary Choice: Motivation


Richard Paap

Examples of binary dependent variables


- Answers to “yes/no” questions
- Choice for private or public health care
- Vote decision for Democrat or Republican president (USA)
- Choice for private or public transport
- Choice to renew or cancel a mobile phone contract
- Business cycle indicator (expansion/recession)

and so forth.

Histogram of data



Scatter diagram



Linear regression model

Test

Suppose we model the binary variable using a regression model

$$\text{response} = \beta_1 + \beta_2 \text{price} + \varepsilon$$

Is it possible to estimate the β parameters using least squares?


Yes, this is possible. Least squares estimation results in:

$$\text{response} = 0.720 - 0.861 \text{price}/1000 + e$$





Lecture 5.1, Slide 5 of 8, Erasmus School of Economics

Scatter diagram with regression line



Lecture 5.1, Slide 6 of 8, Erasmus School of Economics

Model for binary dependent variable



Lecture 5.1, Slide 7 of 8, Erasmus School of Economics

Training Exercise 5.1

- Train yourself by making the training exercise (see the website).
- After making this exercise, check your answers by studying the webcast solution (also available on the website).



Lecture 5.1, Slide 8 of 8, Erasmus School of Economics

MOOC Econometrics

Lecture 5.2 on Binary Choice: Representation

Richard Paap

Logit model

Individual-specific probabilities:

$$\Pr[y_i = 1] = \pi_i,$$

where the value of π_i depends on the explanatory variable x_i .

Logit model:

$$\Pr[y_i = 1] = \frac{\exp(\beta_1 + \beta_2 x_i)}{1 + \exp(\beta_1 + \beta_2 x_i)}$$

and

$$\begin{aligned} \Pr[y_i = 0] &= 1 - \frac{\exp(\beta_1 + \beta_2 x_i)}{1 + \exp(\beta_1 + \beta_2 x_i)} \\ &= \frac{1}{1 + \exp(\beta_1 + \beta_2 x_i)} \end{aligned}$$

Introduction

Let y_i be a binary variable with value 0 or 1 and assume

$$y_i \sim \text{Bernoulli}(\pi),$$

such that

$$\pi = \Pr[y_i = 1] \text{ with } 0 < \pi < 1$$


and hence

$$\Pr[y_i = 0] = 1 - \pi.$$


Individual-specific probabilities:

$$\Pr[y_i = 1] = \pi_i$$

Graphical interpretation




Graphical interpretation




Lecture 5.2, Slide 5 of 12, Erasmus School of Economics

Graphical interpretation



Lecture 5.2, Slide 6 of 12, Erasmus School of Economics

Graphical interpretation



Lecture 5.2, Slide 7 of 12, Erasmus School of Economics

Test question

Test

What happens to the location and shape of the logit function

$$\frac{\exp(\beta_1 + x)}{1 + \exp(\beta_1 + x)}$$

if you change the β_1 parameter from $\beta_1 = 0$ to $\beta_1 = -2$?

The logit function only shifts 2 units to the right.

Lecture 5.2, Slide 8 of 12, Erasmus School of Economics

Odds ratio

Logit model:

$$\Pr[y_i = 1] = \frac{\exp(\beta_1 + \beta_2 x_i)}{1 + \exp(\beta_1 + \beta_2 x_i)}$$
$$\Pr[y_i = 0] = \frac{1}{1 + \exp(\beta_1 + \beta_2 x_i)}$$

Odds ratio:

$$\frac{\Pr[y_i = 1]}{\Pr[y_i = 0]} = \exp(\beta_1 + \beta_2 x_i)$$

Log odds ratio:

$$\log \left(\frac{\Pr[y_i = 1]}{\Pr[y_i = 0]} \right) = \beta_1 + \beta_2 x_i$$



Lecture 5.2, Slide 9 of 12, Erasmus School of Economics

More explanatory variables

Logit specification with x_{2i}, \dots, x_{ki} as explanatory variables:

$$\Pr[y_i = 1] = \frac{\exp(\beta_1 + \sum_{j=2}^k \beta_j x_{ji})}{1 + \exp(\beta_1 + \sum_{j=2}^k \beta_j x_{ji})}$$


Log odds ratio:

$$\log \left(\frac{\Pr[y_i = 1]}{\Pr[y_i = 0]} \right) = \beta_1 + \sum_{j=2}^k \beta_j x_{ji}$$

Marginal effect:

$$\frac{\partial \Pr[y_i = 1]}{\partial x_{ji}} = \Pr[y_i = 1] \Pr[y_i = 0] \beta_j \text{ for } j = 2, \dots, k.$$

Change in probability that $y_i = 1$ due to change in x_{ji} .



Lecture 5.2, Slide 11 of 12, Erasmus School of Economics

Marginal effect

Marginal effect:

$$\frac{d \Pr[y_i = 1]}{d x_i} = \Pr[y_i = 1] \Pr[y_i = 0] \beta_2$$

Change in probability that $y_i = 1$ due to change in x_i .

Average marginal effect:

$$\frac{1}{n} \sum_{i=1}^n \frac{d \Pr[y_i = 1]}{d x_i} = \left(\frac{1}{n} \sum_{i=1}^n \Pr[y_i = 1] \Pr[y_i = 0] \right) \beta_2$$



Lecture 5.2, Slide 10 of 12, Erasmus School of Economics

Training Exercise 5.2

- Train yourself by making the training exercise (see the website).
- After making this exercise, check your answers by studying the webcast solution (also available on the website).



Lecture 5.2, Slide 12 of 12, Erasmus School of Economics

MOOC Econometrics

Lecture 5.3 on Binary Choice: Estimation

Richard Paap

Construction of likelihood function

Likelihood contribution for

- observation $y_i = 1$:

$$\Pr[y_i = 1] = \frac{\exp(x_i' \beta)}{1 + \exp(x_i' \beta)}$$

- observation $y_i = 0$:

$$\Pr[y_i = 0] = \frac{1}{1 + \exp(x_i' \beta)}$$

Likelihood contribution for observation i :

$$\left(\frac{\exp(x_i' \beta)}{1 + \exp(x_i' \beta)} \right)^{y_i} \left(\frac{1}{1 + \exp(x_i' \beta)} \right)^{1-y_i}$$

Introduction

Logit model specification in vector notation:

$$\Pr[y_i = 1] = \frac{\exp(x_i' \beta)}{1 + \exp(x_i' \beta)},$$

where $x_i = (1, x_{2i}, \dots, x_{ki})'$ and $\beta = (\beta_1, \dots, \beta_k)'$

It is not possible to write this model in regression notation

$$y_i = x_i' \beta + \varepsilon_i$$

We use maximum likelihood for parameter estimation.

(Log)-likelihood function

Likelihood function of n independent observations:

$$L(\beta) = \prod_{i=1}^n \left(\frac{\exp(x_i' \beta)}{1 + \exp(x_i' \beta)} \right)^{y_i} \left(\frac{1}{1 + \exp(x_i' \beta)} \right)^{1-y_i}$$

Log-likelihood function:

$$\begin{aligned} \log(L(\beta)) &= \sum_{i=1}^n y_i \log \left(\frac{\exp(x_i' \beta)}{1 + \exp(x_i' \beta)} \right) + (1 - y_i) \log \left(\frac{1}{1 + \exp(x_i' \beta)} \right) \\ &= \sum_{i=1}^n y_i x_i' \beta - \log(1 + \exp(x_i' \beta)), \end{aligned}$$

where we use that $\log(ab) = \log(a) + \log(b)$, $\log(a^b) = b \log(a)$ and $\log(a/b) = \log(a) - \log(b)$.

Test question

Test

The maximum likelihood estimator [MLE] is the value of β that maximizes the log-likelihood function. Is the MLE also the value that maximizes the likelihood function?

As the log function is a monotonically increasing function in β , the answer is yes.



Lecture 5.3, Slide 5 of 1, Erasmus School of Economics

Test question


Test

Suppose that all observations on y_i are 0, that is, $y_i = 0$ for $i = 1, \dots, n$. What is the value of the maximum likelihood estimator in this case?

When all observations are 0, the first-order conditions imply that

$$\frac{1}{n} \sum_{i=1}^n \frac{\exp(x_i' b)}{1 + \exp(x_i' b)} = \frac{1}{n} \sum_{i=1}^n y_i = 0.$$

As the logit function is always larger than 0, there is no value of b for which the first-order conditions holds. The MLE does not exist.



Lecture 5.3, Slide 7 of 1, Erasmus School of Economics

Maximum likelihood estimation

The MLE b is obtained by maximizing $\log(L(\beta))$ with respect to β . First-order conditions:

$$\begin{aligned} \frac{\partial \log(L(\beta))}{\partial \beta} &= \frac{\partial \sum_{i=1}^n y_i x_i' \beta - \log(1 + \exp(x_i' \beta))}{\partial \beta} = 0 \\ &= \sum_{i=1}^n y_i x_i' - \frac{\exp(x_i' \beta) x_i'}{1 + \exp(x_i' \beta)} = 0 \end{aligned}$$

Use numerical methods to solve for β .

The first-order conditions imply that

$$\frac{1}{n} \sum_{i=1}^n \frac{\exp(x_i' b)}{1 + \exp(x_i' b)} = \frac{1}{n} \sum_{i=1}^n y_i$$



Lecture 5.3, Slide 6 of 1, Erasmus School of Economics

Properties of maximum likelihood estimator

It can be shown that under regularity conditions the maximum likelihood estimator [MLE] is

- 1 Consistent
- 2 Efficient for large n
- 3 Asymptotically normally distributed, and hence

$$b \approx N(\beta, V)$$

The (co)variance matrix V can be estimated by

$$\hat{V} = \left(\sum_{i=1}^n \left(\frac{\exp(x_i' b)}{1 + \exp(x_i' b)} \right) \left(\frac{1}{1 + \exp(x_i' b)} \right) x_i x_i' \right)^{-1}$$



Lecture 5.3, Slide 8 of 1, Erasmus School of Economics

Testing for single parameter restriction

We want to compare

- logit model without parameter restrictions
- logit model with a single $\beta_j = 0$

Hypothesis:

$$H_0: \beta_j = 0 \text{ versus } H_1: \beta_j \neq 0$$

You can use the t -test like in a linear regression: Test statistic:

$$z_j = \frac{b_j - 0}{SE(b_j)} \approx N(0, 1),$$

where $SE(b_j)$ is the standard error of b_j .



Lecture 5.3, Slide 9 of 1, Erasmus School of Economics

Testing for a set of parameter restrictions

We want to compare

- logit model without parameter restrictions and estimates b_1
- logit model with m parameter restrictions and estimates b_0

The null hypothesis is that the m parameter restrictions are correct.

To compute the test statistic we need

- $L(b_1)$: maximum likelihood value in full model
- $L(b_0)$: maximum likelihood value in restricted model

Test statistic:

$$LR = -2(\log(L(b_0)) - \log(L(b_1))) \approx \chi^2(m),$$


where m is the number of restrictions.



Lecture 5.3, Slide 10 of 1, Erasmus School of Economics

Training Exercise 5.3

- Train yourself by making the training exercise (see the website).
- After making this exercise, check your answers by studying the webcast solution (also available on the website).



Lecture 5.3, Slide 11 of 1, Erasmus School of Economics

MOOC Econometrics

Lecture 5.4 on Binary Choice: Evaluation

Richard Paap

Erasmus University Rotterdam



Test question

Test

Suppose that we have perfect fit for all n observations, that is,

$$y_i - \frac{\exp(x_i' \beta)}{1 + \exp(x_i' \beta)} \approx 0$$

for all i . What is the numerical value of the likelihood function

$$\prod_{i=1}^n \left(\frac{\exp(x_i' \beta)}{1 + \exp(x_i' \beta)} \right)^{y_i} \left(\frac{1}{1 + \exp(x_i' \beta)} \right)^{1-y_i} ?$$

For all observations equal to 1 (or 0) the likelihood contribution is very close to 1. Hence, the likelihood function equals about 1.



Residuals

Logit residuals:

$$\begin{aligned} y_i - E[y_i] &= y_i - (0 \times \Pr[y_i = 0] + 1 \times \Pr[y_i = 1]) \\ &= y_i - \Pr[y_i = 1] \\ &= y_i - \frac{\exp(x_i' b)}{1 + \exp(x_i' b)} \end{aligned}$$

Interesting cases:

- Lower bound: $y_i - E[y_i] \approx -1$
- Upper bound: $y_i - E[y_i] \approx 1$
- Perfect fit $y_i - E[y_i] \approx 0$



Measures of fit

Define

- $L(b)$: the maximum value of the likelihood function of the model under consideration
- $L(b_1)$: maximum value of the likelihood function in case the model only contains an intercept.

Perfect fit corresponds to $L(b) \approx 1$ or $\log(L(b)) \approx 0$.

Two popular pseudo R^2 measures are:

- McFadden R^2 :

$$R^2 = 1 - \frac{\log(L(b))}{\log(L(b_1))}$$

- Nagelkerke R^2 :

$$R^2 = 1 - \frac{1 - \left(\frac{L(b_1)}{L(b)} \right)^{2/n}}{1 - L(b_1)^{2/n}}$$



Prediction probability

If the value of x_{n+1} is available, one can predict the value of y_{n+1} using

$$\begin{aligned} E[y_{n+1}] &= 0 \times \Pr[y_{n+1} = 0] + 1 \times \Pr[y_{n+1} = 1] \\ &= \Pr[y_{n+1} = 1] \\ &= \frac{\exp(x'_{n+1}\beta)}{1 + \exp(x'_{n+1}\beta)} \end{aligned}$$

To estimate this probability we replace β by its estimate b and obtain $\hat{\Pr}[y_{n+1} = 1]$.




Lecture 5.4, Slide 5 of 12, Erasmus School of Economics

Test question

Test

Does a higher value of the cut-off value c generate more, the same or less predictions which are equal to 1?

A higher value of c means that less (or the same number of) prediction probabilities are above c and hence you forecast less (or the same number of) ones.



Lecture 5.4, Slide 7 of 12, Erasmus School of Economics

0/1 Prediction

The prediction is a probability and never exactly equal to 0 or 1.

Transform the prediction probability into 0/1 forecast \hat{y}_{n+1} by the rule:

$$\begin{aligned} \hat{y}_{n+1} &= 1 \text{ if } \hat{\Pr}[y_{n+1} = 1] > c \\ \hat{y}_{n+1} &= 0 \text{ if } \hat{\Pr}[y_{n+1} = 1] \leq c. \end{aligned}$$

Many statistical packages use $c = 0.5$. However, one may also consider

$$c = \frac{1}{n} \sum_{i=1}^n y_i,$$

that is, the fraction of observations in the sample equal to one.



Lecture 5.4, Slide 6 of 12, Erasmus School of Economics

Evaluation of predication accuracy

Suppose one has m out-of-sample predictions for y_i denoted by \hat{y}_i .

Count the number of correct and incorrect predictions:

$$m_{11} = \sum_{i=1}^m y_{n+i} \hat{y}_{n+i} \quad \text{data}=1 \text{ \& prediction}=1$$

$$m_{00} = \sum_{i=1}^m (1 - y_{n+i})(1 - \hat{y}_{n+i}) \quad \text{data}=0 \text{ \& prediction}=0$$

$$m_{10} = \sum_{i=1}^m y_{n+i}(1 - \hat{y}_{n+i}) \quad \text{data}=1 \text{ \& prediction}=0$$

$$m_{01} = \sum_{i=1}^m (1 - y_{n+i})\hat{y}_{n+i} \quad \text{data}=0 \text{ \& prediction}=1$$



Lecture 5.4, Slide 8 of 12, Erasmus School of Economics

Prediction-realization table

Classify predictions in right and wrong:

observed	predicted		sum
	$\hat{y} = 0$	$\hat{y} = 1$	
$y = 0$	m_{00}/m	m_{01}/m	$(m_{00} + m_{01})/m$
$y = 1$	m_{10}/m	m_{11}/m	$(m_{10} + m_{11})/m$
sum	$(m_{00} + m_{10})/m$	$(m_{01} + m_{11})/m$	1




Lecture 5.4, Slide 9 of 12, Erasmus School of Economics

Prediction-realization table

Classify predictions in right and wrong:

observed	predicted		sum
	$\hat{y} = 0$	$\hat{y} = 1$	
$y = 0$	m_{00}/m	m_{01}/m	$(m_{00} + m_{01})/m$
$y = 1$	m_{10}/m	m_{11}/m	$(m_{10} + m_{11})/m$
sum	$(m_{00} + m_{10})/m$	$(m_{01} + m_{11})/m$	1

$m_{01}/m + m_{10}/m$ denotes the fraction of incorrect forecasts.



Lecture 5.4, Slide 11 of 12, Erasmus School of Economics

Prediction-realization table

Classify predictions in right and wrong:

observed	predicted		sum
	$\hat{y} = 0$	$\hat{y} = 1$	
$y = 0$	m_{00}/m	m_{01}/m	$(m_{00} + m_{01})/m$
$y = 1$	m_{10}/m	m_{11}/m	$(m_{10} + m_{11})/m$
sum	$(m_{00} + m_{10})/m$	$(m_{01} + m_{11})/m$	1

The fraction $m_{00}/m + m_{11}/m$ is called the hit rate.



Lecture 5.4, Slide 10 of 12, Erasmus School of Economics

Training Exercise 5.4

- Train yourself by making the training exercise (see the website).
- After making this exercise, check your answers by studying the webcast solution (also available on the website).



Lecture 5.4, Slide 12 of 12, Erasmus School of Economics

MOOC Econometrics

Lecture 5.5 on Binary Choice: Application

Richard Paap

Response to direct mailing

Sample:

- 925 observations

Dependent variable:

- Resp: Response to direct mailing with 1 = yes and 0 = no

Potential explanatory variables:

- Male: 1 = Male and 0 = Female
- Age: Age of the customer in years
- Active: 1 = Active customer and 0 = Inactive customer

Data characteristics

Average values of the explanatory variables

Variable	resp = 0	resp = 1	all observations
Gender	0.624	0.823	0.725
Active	0.114	0.260	0.188
Age	50.813	50.553	50.681

Data characteristics

Average values of the explanatory variables

Variable	resp = 0	resp = 1	all observations
Gender	0.624	0.823	0.725
Active	0.114	0.260	0.188
Age	50.813	50.553	50.681


Model specification

Proposed logit model specification:

$$\Pr[\text{resp}_i = 1] =$$

$$\frac{\exp(\beta_0 + \beta_1 \text{male}_i + \beta_2 \text{active}_i + \beta_3 \text{age}_i + \beta_4 (\text{age}_i/10)^2)}{1 + \exp(\beta_0 + \beta_1 \text{male}_i + \beta_2 \text{active}_i + \beta_3 \text{age}_i + \beta_4 (\text{age}_i/10)^2)}$$

for $i = 1, \dots, 925$.



Lecture 5.5, Slide 5 of 15, Erasmus School of Economics

Estimation results logit model

Variable	Coefficient	Std. Error	t-value	p-value.
Intercept	-2.488	0.890	-2.796	0.005
Male	0.954	0.158	6.029	0.000
Active	0.914	0.185	4.945	0.000
Age	0.070	0.036	1.964	0.050
(Age/10) ²	-0.069	0.034	-2.015	0.044
McFadden R^2	0.061			
Nagelkerke R^2	0.892			
Log-likelihood	-601.862			



Lecture 5.5, Slide 6 of 15, Erasmus School of Economics


Odds ratio

$$\frac{\Pr[\text{resp}_i = 1]}{\Pr[\text{resp}_i = 0]}$$

$$= \exp(\beta_0 + \beta_1 \text{male}_i + \beta_2 \text{active}_i + \beta_3 \text{age}_i + \beta_4 (\text{age}_i/10)^2)$$

$$\approx \exp(-2.49 + 0.95 \text{male}_i + 0.91 \text{active}_i + 0.07 \text{age}_i - 0.07 (\text{age}_i/10)^2)$$

$$= 0.08 \times 2.57^{\text{male}_i} \times 2.50^{\text{active}_i} \times \exp(0.07 \text{age}_i - 0.07 (\text{age}_i/10)^2)$$



Lecture 5.5, Slide 7 of 15, Erasmus School of Economics


Odds ratio

$$\frac{\Pr[\text{resp}_i = 1]}{\Pr[\text{resp}_i = 0]}$$

$$= \exp(\beta_0 + \beta_1 \text{male}_i + \beta_2 \text{active}_i + \beta_3 \text{age}_i + \beta_4 (\text{age}_i/10)^2)$$

$$\approx \exp(-2.49 + 0.95 \text{male}_i + 0.91 \text{active}_i + 0.07 \text{age}_i - 0.07 (\text{age}_i/10)^2)$$


$$= 0.08 \times 2.57^{\text{male}_i} \times 2.50^{\text{active}_i} \times \exp(0.07 \text{age}_i - 0.07 (\text{age}_i/10)^2)$$



Lecture 5.5, Slide 8 of 15, Erasmus School of Economics

Odds ratio

$$\begin{aligned} \frac{\Pr[\text{resp}_i = 1]}{\Pr[\text{resp}_i = 0]} &= \exp(\beta_0 + \beta_1 \text{male}_i + \beta_2 \text{active}_i + \beta_3 \text{age}_i + \beta_4 (\text{age}_i/10)^2) \\ &\approx \exp(-2.49 + 0.95 \text{male}_i + 0.91 \text{active}_i + 0.07 \text{age}_i - 0.07 (\text{age}_i/10)^2) \\ &= 0.08 \times 2.57^{\text{male}_i} \times 2.50^{\text{active}_i} \times \exp(0.07 \text{age}_i - 0.07 (\text{age}_i/10)^2) \end{aligned}$$



Lecture 5.5, Slide 9 of 15, Erasmus School of Economics

Test question

Test


For which value of age do we have the highest value of the odds ratio

$$0.08 \times 2.57^{\text{male}_i} \times 2.50^{\text{active}_i} \times \exp(0.07 \text{age}_i - 0.07 (\text{age}_i/10)^2)?$$

The first-order condition is


$$\begin{aligned} [0.08 \times 2.57^{\text{male}_i} \times 2.50^{\text{active}_i} \times \exp(0.07 \text{age}_i - 0.07 (\text{age}_i/10)^2)] \\ \times (0.07 - 2 \times 0.07 (\text{age}_i/100)) = 0 \end{aligned}$$

The solution to this first-order condition is 50 years.



Lecture 5.5, Slide 10 of 15, Erasmus School of Economics

Odds ratio versus age



Lecture 5.5, Slide 11 of 15, Erasmus School of Economics

Marginal effect of age

$$\begin{aligned} \frac{\partial \Pr[\text{resp}_i = 1]}{\partial \text{age}_i} &= \Pr[\text{resp}_i = 1] \Pr[\text{resp}_i = 0] (\beta_3 + 2\beta_4 (\text{age}_i/10)^2) \\ &\approx \Pr[\text{resp}_i = 1] \Pr[\text{resp}_i = 0] (0.07 - 2 \times 0.07 \text{age}_i/100) \end{aligned}$$


Marginal effect depends on

- age_i
- $\Pr[\text{resp}_i = 1]$ and $\Pr[\text{resp}_i = 0]$ and hence also on male and active dummy.



Lecture 5.5, Slide 12 of 15, Erasmus School of Economics

Marginal effect of age



Lecture 5.5, Slide 13 of 15, Erasmus School of Economics

In-sample prediction-realisation table

Cut-off value: 0.5

observed	predicted		sum
	$\hat{y} = 0$	$\hat{y} = 1$	
$y = 0$	0.212	0.280	0.492
$y = 1$	0.104	0.404	0.508
sum	0.316	0.684	1

Hit rate: $0.212 + 0.404 = 0.616$.

Erasmus

Lecture 5.5, Slide 14 of 15, Erasmus School of Economics

Training Exercise 5.5

- Train yourself by making the training exercise (see the website).
- After making this exercise, check your answers by studying the webcast solution (also available on the website).

Erasmus

Lecture 5.5, Slide 15 of 15, Erasmus School of Economics