

Dice revisited, an example of S. N. Bernstein

Dice revisited, an example of S. N. Bernstein

- ❖ A die is thrown twice.

Dice revisited, an example of S. N. Bernstein

- ❖ A die is thrown twice.
 - ❖ *Sample space:* $\Omega = \{(j, k): 1 \leq j, k \leq 6\}$.

Dice revisited, an example of S. N. Bernstein

- ❖ A die is thrown twice.
 - ❖ *Sample space:* $\Omega = \{(j, k): 1 \leq j, k \leq 6\}$.
 - ❖ *Events:*

Dice revisited, an example of S. N. Bernstein

- ❖ A die is thrown twice.
 - ❖ *Sample space:* $\Omega = \{(j, k): 1 \leq j, k \leq 6\}$.
 - ❖ *Events:*
 - ❖ $A :=$ The first throw shows an odd number.

Dice revisited, an example of S. N. Bernstein

- ❖ A die is thrown twice.
 - ❖ *Sample space:* $\Omega = \{(j, k): 1 \leq j, k \leq 6\}$.
 - ❖ *Events:*
 - ❖ $A :=$ The first throw shows an odd number.
 - ❖ $B :=$ The second throw shows an even number.

Dice revisited, an example of S. N. Bernstein

- ❖ A die is thrown twice.
 - ❖ *Sample space:* $\Omega = \{(j, k): 1 \leq j, k \leq 6\}$.
 - ❖ *Events:*
 - ❖ $A :=$ The first throw shows an odd number.
 - ❖ $B :=$ The second throw shows an even number.
 - ❖ $C :=$ The sum of face values is odd.

Dice revisited, an example of S. N. Bernstein

- ✧ A die is thrown twice.
 - ✧ *Sample space:* $\Omega = \{(j, k): 1 \leq j, k \leq 6\}$.
 - ✧ *Events:*
 - ✧ $A :=$ The first throw shows an odd number.
 - ✧ $B :=$ The second throw shows an even number.
 - ✧ $C :=$ The sum of face values is odd.
 - ✧ *Probability measure:* Combinatorial setting with mass function assigning equal probability $1/36$ to each atom.

Dice revisited, an example of S. N. Bernstein

- ❖ A die is thrown twice.
 - ❖ *Sample space:* $\Omega = \{(j, k): 1 \leq j, k \leq 6\}$.
 - ❖ *Events:*
 - ❖ $A :=$ The first throw shows an odd number.
 - ❖ $B :=$ The second throw shows an even number.
 - ❖ $C :=$ The sum of face values is odd.
 - ❖ *Probability measure:* Combinatorial setting with mass function assigning equal probability $1/36$ to each atom.
- ❖ Are A,B, and C independent events?

Dice revisited, an example of S. N. Bernstein

- * A die is thrown twice.
 - * *Sample space:* $\Omega = \{(j, k): 1 \leq j, k \leq 6\}$.
 - * *Events:*
 - * $A :=$ The first throw shows an odd number.
 - * $B :=$ The second throw shows an even number.
 - * $C :=$ The sum of face values is odd.
 - * *Probability measure:* Combinatorial setting with mass function assigning equal probability $1/36$ to each atom.
- * Are A,B, and C independent events?

Event		Probability
A	$\{(j, k): j \text{ is odd}\}$	
B	$\{(j, k): k \text{ is even}\}$	
C	$\{(j, k): \text{either } (j \text{ is odd and } k \text{ is even}) \text{ or } (j \text{ is even and } k \text{ is odd})\}$	
$A \cap B$	$\{(j, k): j \text{ is odd and } k \text{ is even}\}$	
$A \cap C$	$\{(j, k): j \text{ is odd and } k \text{ is even}\}$	
$B \cap C$	$\{(j, k): j \text{ is odd and } k \text{ is even}\}$	
$A \cap B \cap C$	$\{(j, k): j \text{ is odd and } k \text{ is even}\}$	

Dice revisited, an example of S. N. Bernstein

- ❖ A die is thrown twice.
 - ❖ *Sample space:* $\Omega = \{(j, k): 1 \leq j, k \leq 6\}$.
 - ❖ *Events:*
 - ❖ $A :=$ The first throw shows an odd number.
 - ❖ $B :=$ The second throw shows an even number.
 - ❖ $C :=$ The sum of face values is odd.
 - ❖ *Probability measure:* Combinatorial setting with mass function assigning equal probability $1/36$ to each atom.
- ❖ Are A,B, and C independent events?

Event		Probability
A	$\{(j, k): j \text{ is odd}\}$	$\frac{3 \times 6}{36} = \frac{1}{2}$
B	$\{(j, k): k \text{ is even}\}$	
C	$\{(j, k): \text{either } (j \text{ is odd and } k \text{ is even}) \text{ or } (j \text{ is even and } k \text{ is odd})\}$	
$A \cap B$	$\{(j, k): j \text{ is odd and } k \text{ is even}\}$	
$A \cap C$	$\{(j, k): j \text{ is odd and } k \text{ is even}\}$	
$B \cap C$	$\{(j, k): j \text{ is odd and } k \text{ is even}\}$	
$A \cap B \cap C$	$\{(j, k): j \text{ is odd and } k \text{ is even}\}$	

Dice revisited, an example of S. N. Bernstein

- ❖ A die is thrown twice.
 - ❖ *Sample space:* $\Omega = \{(j, k): 1 \leq j, k \leq 6\}$.
 - ❖ *Events:*
 - ❖ $A :=$ The first throw shows an odd number.
 - ❖ $B :=$ The second throw shows an even number.
 - ❖ $C :=$ The sum of face values is odd.
 - ❖ *Probability measure:* Combinatorial setting with mass function assigning equal probability $1/36$ to each atom.
- ❖ Are A,B, and C independent events?

Event		Probability
A	$\{(j, k): j \text{ is odd}\}$	$\frac{3 \times 6}{36} = \frac{1}{2}$
B	$\{(j, k): k \text{ is even}\}$	$\frac{6 \times 3}{36} = \frac{1}{2}$
C	$\{(j, k): \text{either } (j \text{ is odd and } k \text{ is even}) \text{ or } (j \text{ is even and } k \text{ is odd})\}$	
$A \cap B$	$\{(j, k): j \text{ is odd and } k \text{ is even}\}$	
$A \cap C$	$\{(j, k): j \text{ is odd and } k \text{ is even}\}$	
$B \cap C$	$\{(j, k): j \text{ is odd and } k \text{ is even}\}$	
$A \cap B \cap C$	$\{(j, k): j \text{ is odd and } k \text{ is even}\}$	

Dice revisited, an example of S. N. Bernstein

- * A die is thrown twice.
 - * *Sample space:* $\Omega = \{(j, k): 1 \leq j, k \leq 6\}$.
 - * *Events:*
 - * $A :=$ The first throw shows an odd number.
 - * $B :=$ The second throw shows an even number.
 - * $C :=$ The sum of face values is odd.
 - * *Probability measure:* Combinatorial setting with mass function assigning equal probability $1/36$ to each atom.
- * Are A,B, and C independent events?

Event		Probability
A	$\{(j, k): j \text{ is odd}\}$	$\frac{3 \times 6}{36} = \frac{1}{2}$
B	$\{(j, k): k \text{ is even}\}$	$\frac{6 \times 3}{36} = \frac{1}{2}$
C	$\{(j, k): \text{either } (j \text{ is odd and } k \text{ is even}) \text{ or } (j \text{ is even and } k \text{ is odd})\}$	$\frac{(3 \times 3) + (3 \times 3)}{36} = \frac{1}{2}$
$A \cap B$	$\{(j, k): j \text{ is odd and } k \text{ is even}\}$	
$A \cap C$	$\{(j, k): j \text{ is odd and } k \text{ is even}\}$	
$B \cap C$	$\{(j, k): j \text{ is odd and } k \text{ is even}\}$	
$A \cap B \cap C$	$\{(j, k): j \text{ is odd and } k \text{ is even}\}$	

Dice revisited, an example of S. N. Bernstein

- * A die is thrown twice.
 - * *Sample space:* $\Omega = \{(j, k): 1 \leq j, k \leq 6\}$.
 - * *Events:*
 - * $A :=$ The first throw shows an odd number.
 - * $B :=$ The second throw shows an even number.
 - * $C :=$ The sum of face values is odd.
 - * *Probability measure:* Combinatorial setting with mass function assigning equal probability $1/36$ to each atom.
- * Are A,B, and C independent events?

Event		Probability
A	$\{(j, k): j \text{ is odd}\}$	$\frac{3 \times 6}{36} = \frac{1}{2}$
B	$\{(j, k): k \text{ is even}\}$	$\frac{6 \times 3}{36} = \frac{1}{2}$
C	$\{(j, k): \text{either } (j \text{ is odd and } k \text{ is even}) \text{ or } (j \text{ is even and } k \text{ is odd})\}$	$\frac{(3 \times 3) + (3 \times 3)}{36} = \frac{1}{2}$
$A \cap B$	$\{(j, k): j \text{ is odd and } k \text{ is even}\}$	$\frac{3 \times 3}{36} = \frac{1}{4}$
$A \cap C$	$\{(j, k): j \text{ is odd and } k \text{ is even}\}$	$\frac{3 \times 3}{36} = \frac{1}{4}$
$B \cap C$	$\{(j, k): j \text{ is odd and } k \text{ is even}\}$	$\frac{3 \times 3}{36} = \frac{1}{4}$
$A \cap B \cap C$	$\{(j, k): j \text{ is odd and } k \text{ is even}\}$	

Dice revisited, an example of S. N. Bernstein

- ❖ A die is thrown twice.
 - ❖ *Sample space:* $\Omega = \{(j, k): 1 \leq j, k \leq 6\}$.
 - ❖ *Events:*
 - ❖ $A :=$ The first throw shows an odd number.
 - ❖ $B :=$ The second throw shows an even number.
 - ❖ $C :=$ The sum of face values is odd.
 - ❖ *Probability measure:* Combinatorial setting with mass function assigning equal probability $1/36$ to each atom.
- ❖ Are A,B, and C independent events?

Event		Probability
A	$\{(j, k): j \text{ is odd}\}$	$\frac{3 \times 6}{36} = \frac{1}{2}$
B	$\{(j, k): k \text{ is even}\}$	$\frac{6 \times 3}{36} = \frac{1}{2}$
C	$\{(j, k): \text{either } (j \text{ is odd and } k \text{ is even}) \text{ or } (j \text{ is even and } k \text{ is odd})\}$	$\frac{(3 \times 3) + (3 \times 3)}{36} = \frac{1}{2}$
$A \cap B$	$\{(j, k): j \text{ is odd and } k \text{ is even}\}$	$\frac{3 \times 3}{36} = \frac{1}{4}$
$A \cap C$	$\{(j, k): j \text{ is odd and } k \text{ is even}\}$	$\frac{3 \times 3}{36} = \frac{1}{4}$
$B \cap C$	$\{(j, k): j \text{ is odd and } k \text{ is even}\}$	$\frac{3 \times 3}{36} = \frac{1}{4}$
$A \cap B \cap C$	$\{(j, k): j \text{ is odd and } k \text{ is even}\}$	$\frac{3 \times 3}{36} = \frac{1}{4}$

Dice revisited, an example of S. N. Bernstein

- ❖ A die is thrown twice.
 - ❖ *Sample space:* $\Omega = \{(j, k): 1 \leq j, k \leq 6\}$.
 - ❖ *Events:*
 - ❖ $A :=$ The first throw shows an odd number.
 - ❖ $B :=$ The second throw shows an even number.
 - ❖ $C :=$ The sum of face values is odd.
 - ❖ *Probability measure:* Combinatorial setting with mass function assigning equal probability $1/36$ to each atom.
- ❖ Are A,B, and C independent events?

Event		Probability
A	$\{(j, k): j \text{ is odd}\}$	$\frac{3 \times 6}{36} = \frac{1}{2}$
B	$\{(j, k): k \text{ is even}\}$	$\frac{6 \times 3}{36} = \frac{1}{2}$
C	$\{(j, k): \text{either } (j \text{ is odd and } k \text{ is even}) \text{ or } (j \text{ is even and } k \text{ is odd})\}$	$\frac{(3 \times 3) + (3 \times 3)}{36} = \frac{1}{2}$
$A \cap B$	$\{(j, k): j \text{ is odd and } k \text{ is even}\}$	$\frac{3 \times 3}{36} = \frac{1}{4} = \frac{1}{2} \times \frac{1}{2}$
$A \cap C$	$\{(j, k): j \text{ is odd and } k \text{ is even}\}$	$\frac{3 \times 3}{36} = \frac{1}{4} = \frac{1}{2} \times \frac{1}{2}$
$B \cap C$	$\{(j, k): j \text{ is odd and } k \text{ is even}\}$	$\frac{3 \times 3}{36} = \frac{1}{4} = \frac{1}{2} \times \frac{1}{2}$
$A \cap B \cap C$	$\{(j, k): j \text{ is odd and } k \text{ is even}\}$	$\frac{3 \times 3}{36} = \frac{1}{4} \neq \frac{1}{2} \times \frac{1}{2} \times \frac{1}{2}$

Dice revisited, an example of S. N. Bernstein

- ❖ A die is thrown twice.
 - ❖ *Sample space:* $\Omega = \{(j, k): 1 \leq j, k \leq 6\}$.
 - ❖ *Events:*
 - ❖ $A :=$ The first throw shows an odd number.
 - ❖ $B :=$ The second throw shows an even number.
 - ❖ $C :=$ The sum of face values is odd.
 - ❖ *Probability measure:* Combinatorial setting with mass function assigning equal probability $1/36$ to each atom.
- ❖ Are A,B, and C independent events?

Event		Probability
A	$\{(j, k): j \text{ is odd}\}$	$\frac{3 \times 6}{36} = \frac{1}{2}$
B	$\{(j, k): k \text{ is even}\}$	$\frac{6 \times 3}{36} = \frac{1}{2}$
C	$\{(j, k): \text{either } (j \text{ is odd and } k \text{ is even}) \text{ or } (j \text{ is even and } k \text{ is odd})\}$	$\frac{(3 \times 3) + (3 \times 3)}{36} = \frac{1}{2}$
$A \cap B$	$\{(j, k): j \text{ is odd and } k \text{ is even}\}$	$\frac{3 \times 3}{36} = \frac{1}{4} = \frac{1}{2} \times \frac{1}{2}$
$A \cap C$	$\{(j, k): j \text{ is odd and } k \text{ is even}\}$	$\frac{3 \times 3}{36} = \frac{1}{4} = \frac{1}{2} \times \frac{1}{2}$
$B \cap C$	$\{(j, k): j \text{ is odd and } k \text{ is even}\}$	$\frac{3 \times 3}{36} = \frac{1}{4} = \frac{1}{2} \times \frac{1}{2}$
$A \cap B \cap C$	$\{(j, k): j \text{ is odd and } k \text{ is even}\}$	$\frac{3 \times 3}{36} = \frac{1}{4} \neq \frac{1}{2} \times \frac{1}{2} \times \frac{1}{2}$

A, B, and C are pairwise independent but not (jointly) independent.

- 1) $\mathbf{P(A \cap B) = P(A) \times P(B),}$
- 2) $\mathbf{P(A \cap C) = P(A) \times P(C),}$
- 3) $\mathbf{P(B \cap C) = P(B) \times P(C),}$
- 4) $\mathbf{P(A \cap B \cap C) = P(A) \times P(B) \times P(C).}$

- 1) $\mathbf{P(A \cap B) = P(A) \times P(B),}$
- 2) $\mathbf{P(A \cap C) = P(A) \times P(C),}$
- 3) $\mathbf{P(B \cap C) = P(B) \times P(C),}$
- 4) $\mathbf{P(A \cap B \cap C) = P(A) \times P(B) \times P(C).}$

Independence implies pairwise independence but pairwise independence does not imply independence.