





Like this course? Become an expert by joining the Methods and Statistics in Social Sciences Specialization (/specializations/social-science).

Upgrade

(https://accounts.coursera.org/i/zendesk/courserahelp?return_to=https://learner.coursera.help/hc)

Course Forums

All Course Discussions (/learn/infer...
General Discussion (/learn/inferenti...
Meet and Greet (/learn/inferential-s...

Module Forums

Before we get started... (/learn/infer...
Comparing two groups (/learn/infer...
Categorical association (/learn/infer...
Simple regression (/learn/inferentia...
Multiple regression (/learn/inferenti...
Analysis of variance (/learn/inferent...
Non-parametric tests (/learn/infere...
Exam time! (/learn/inferential-statis...

← All Course Discussions

(/learn/inferential-statistics/discussions?sort=lastActivityAtDesc&page=1&q=)

DS

Exam Q22 - is the auto- ▼

Help Center

(/learn/inferential-statistics/discussions/CTbXUvgnEeWYIQ4BcV9YKQ)

statistics/profiles/1f7f04368de1cdf42d40ee185779133)

(/learn/inferential-statistics/module/jyCOE/discussions/CTbXL

There is more than one way to skin a cat for two-sided test of dependent proportions. They should all drive the same p-value, but the associated entry 0.xxx is not accepted.

Whether for a mcnemar (correct=FALSE) chi-squared or z-test as per course PDF, key values seem to be Pass/No Graduate (64) and Graduate/No Pass (81).

```
x <- matrix(data=c(97, 64, 81, 158), nrow=2, byrow=FALSE)
```

This sets 64 and 81 as the key values as they are the off-diagonal entries. It could be reformatted for Pass/Graduate as x[1,1] and/or swap x[2,1] with x[1,2] (or both), but those all drive the same two-sided p-value.

The data are paired and a test of independent

proportions seems inappropriate. The p-values using mcnemar chi-squared (correct=FALSE) or z (course PDF) are the same.

Am I reading or interpreting the question wrong? Does the auto-grader want a format other than 0.xxx? Does it want the p-value from a continuity corrected mcnemar chi-squared (correct=TRUE)? Something else?

Reply

▲ 0

Follow

to

Upvote

2

David

Scott

Earliest (/learn/inferential-statistics/discussions/CTbXUvgnEeWYIQ4BcV9YKQ?sort=createdAtAsc&page=1)

Top (/learn/inferential-statistics/discussions/CTbXUvgnEeWYIQ4BcV9YKQ?sort=upvotesDesc&page=1)

Most Recent (/learn/inferential-statistics/discussions/CTbXUvgnEeWYIQ4BcV9YKQ?sort=createdAtDesc&page=1) 4 days ago (/learn/inferential-statistics/discussions/CTbXUvgnEeWYIQ4BcV9YKQ/replies/8o7t8vpXEeWEAhLOZ9BW_w)

Most Recent (/learn/inferential-statistics/discussions/CTbXUvgnEeWYIQ4BcV9YKQ?sort=createdAtDesc&page=1)

(/learn/inferential-statistics/profiles/3cdb3ac214a020ebf6e64f1e50b03eac)

The autograder had an extra decimal place. It's changed now so your answer should be correct when you round to x.xxx. Very sorry about that!

▲ 0 Upvote · Reply

< 1 >

SD

(/learn/inferential-

Reply

statistics/profiles/f34069ce8df6de7dbebfedfb7e760d9f)

Reply