

Eigenvalues and eigenvectors

10/10 points (100%)

Quiz, 10 questions

✓ **Congratulations! You passed!**

Next Item


 1 / 1
points

1.

This assessment will test your ability to apply your knowledge of eigenvalues and eigenvectors to some special cases.

Use the following code blocks to assist you in this quiz. They calculate eigenvectors and eigenvalues respectively:

```
1 # Eigenvalues
2 M = np.array([[4, -5, 6],
3               [7, -8, 6],
4               [3/2., -1/2., -2]])
5 vals, vecs = np.linalg.eig(M)
6 print(vals)
7 vecs
```

Run

Reset

```
1 # Eigenvectors - Note, the eigenvectors are the columns of the output.
2 M = np.array([[0.1, 0.7, 0.1, 0.1],
3               [0.7, 0.1, 0.1, 0.1],
4               [0.1, 0.1, 0.1, 0.7],
5               [0.1, 0.1, 0.7, 0.1]])
6 vals, vecs = np.linalg.eig(M)
7 print(vals)
8 vecs
9
```

Run

Reset

To practice, select all eigenvectors of the matrix, $A = \begin{bmatrix} 4 & -5 & 6 \\ 7 & -8 & 6 \\ 3/2 & -1/2 & -2 \end{bmatrix}$.



$$\begin{bmatrix} 1/\sqrt{6} \\ -1/\sqrt{6} \\ 2/\sqrt{6} \end{bmatrix}$$



Un-selected is correct

Eigenvalues and eigenvectors

Quiz, 10 questions

10/10 points (100%)



$$\begin{bmatrix} -2/\sqrt{9} \\ 2/\sqrt{9} \\ 1/\sqrt{9} \end{bmatrix}$$

Correct

This is one of the eigenvectors. Note eigenvectors are only defined upto a scale factor.



$$\begin{bmatrix} -3 \\ -3 \\ -1 \end{bmatrix}$$

Correct

This is one of the eigenvectors.



$$\begin{bmatrix} -3 \\ -2 \\ 1 \end{bmatrix}$$

Un-selected is correct



None of the other options.

Un-selected is correct



$$\begin{bmatrix} -1 \\ 1 \\ -2 \end{bmatrix}$$

Un-selected is correct



$$\begin{bmatrix} 1/2 \\ -1/2 \\ -1 \end{bmatrix}$$

Correct

This is one of the eigenvectors. Note eigenvectors are only defined upto a scale factor.

Eigenvalues and eigenvectors

10/10 points (100%)


Quiz, 10 questions

1 / 1
points

2.

Recall from the *PageRank* notebook, that in PageRank, we care about the eigenvector of the link matrix, L , that has eigenvalue 1, and that we can find this using *power iteration method* as this will be the largest eigenvalue.

PageRank can sometimes get into trouble if closed-loop structures appear. A simplified example might look like this,



With link matrix, $L = \begin{bmatrix} 0 & 0 & 0 & 1 \\ 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \end{bmatrix}$.

Use the calculator in Q1 to check the eigenvalues and vectors for this system.

What might be going wrong? Select all that apply.

☐

Some of the eigenvectors are complex.

**Un-selected is correct**☐

The system is too small.

**Un-selected is correct**☐

Because of the loop, *Procrastinating Pats* that are browsing will go around in a cycle rather than settling on a webpage.



Eigenvalues and eigenvectors

Quiz, 10 questions

Correct

If all sites started out populated equally, then the incoming pats would equal the outgoing, but in general the system will not converge to this result by applying power iteration.

10/10 points (100%)

☐

Other eigenvalues are not small compared to 1, and so do not decay away with each power iteration.



Correct

The other eigenvectors have the same size as 1 (they are $-1, i, -i$)

☐

None of the other options.



Un-selected is correct



1 / 1
points

3.

The loop in the previous question is a situation that can be remedied by damping.

If we replace the link matrix with the damped,

$$L' = \begin{bmatrix} 0.1 & 0.1 & 0.1 & 0.7 \\ 0.7 & 0.1 & 0.1 & 0.1 \\ 0.1 & 0.7 & 0.1 & 0.1 \\ 0.1 & 0.1 & 0.7 & 0.1 \end{bmatrix}, \text{ how does this help?}$$

☐

The complex number disappear.



Un-selected is correct

☐

It makes the eigenvalue we want bigger.



Un-selected is correct

☐

None of the other options.



Un-selected is correct

Eigenvalues and eigenvectors

10/10 points (100%)

Quiz, 10 questions



The other eigenvalues get smaller.

**Correct**

So their eigenvectors will decay away on power iteration.



There is now a probability to move to any website.

**Correct**

This helps the power iteration settle down as it will spread out the distribution of Pats

1 / 1
points


4.

Another issue that may come up, is if there are disconnected parts to the internet. Take this example,

Eigenvalues and eigenvectors

10/10 points (100%)

Quiz, 10 questions



with link matrix, $L = \begin{bmatrix} 0 & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \end{bmatrix}$.

This form is known as block diagonal, as it can be split into square blocks along the main diagonal, i.e., $L = \begin{bmatrix} A & 0 \\ 0 & B \end{bmatrix}$, with $A = B = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}$ in this case.

What is happening in this system?

☒ There are loops in the system.



Correct

There are two loops of size 2. ($A \rightleftarrows B$) and ($C \rightleftarrows D$)

☐ There isn't a unique PageRank.



Correct

The power iteration algorithm could settle to multiple values, depending on its starting conditions.

☐ None of the other options.



Un-selected is correct

☐ There are two eigenvalues of 1.



Correct

The eigensystem is degenerate. Any linear combination of eigenvectors with the same eigenvalue is also an eigenvector.

Eigenvalues and eigenvectors

10/10 points (100%)

Quiz, 10 questions



The system has zero determinant.



Un-selected is correct



1 / 1
points

5.

By similarly applying damping to the link matrix from the previous question. What happens now?



The negative eigenvalues disappear.



Un-selected is correct



There becomes two eigenvalues of 1.



Un-selected is correct



Damping does not help this system.



Un-selected is correct



None of the other options.



Correct

There is now only one eigenvalue of 1, and PageRank will settle to it's eigenvector through repeating the power iteration method.



The system settles into a single loop.



Un-selected is correct

Eigenvalues and eigenvectors

10/10 points (100%)

Quiz, 10 questions


 1 / 1
points

6.

Given the matrix $A = \begin{bmatrix} 3/2 & -1 \\ -1/2 & 1/2 \end{bmatrix}$, calculate its characteristic polynomial.

☒ $\lambda^2 - 2\lambda + \frac{1}{4}$


Correct

Well done - this is indeed the characteristic polynomial of A .

☐ $\lambda^2 + 2\lambda - \frac{1}{4}$

☐ $\lambda^2 + 2\lambda + \frac{1}{4}$

☐ $\lambda^2 - 2\lambda - \frac{1}{4}$


 1 / 1
points

7.

By solving the characteristic polynomial above or otherwise, calculate the

eigenvalues of the matrix $A = \begin{bmatrix} 3/2 & -1 \\ -1/2 & 1/2 \end{bmatrix}$.

☒ $\lambda_1 = 1 - \frac{\sqrt{3}}{2}, \lambda_2 = 1 + \frac{\sqrt{3}}{2}$


Correct

Well done! These are the roots of the above characteristic polynomial, and hence these are the eigenvalues of A .

☐ $\lambda_1 = 1 - \frac{\sqrt{5}}{2}, \lambda_2 = 1 + \frac{\sqrt{5}}{2}$

☐ $\lambda_1 = -1 - \frac{\sqrt{5}}{2}, \lambda_2 = -1 + \frac{\sqrt{5}}{2}$



$$\lambda_1 = -1 - \frac{\sqrt{3}}{2}, \lambda_2 = -1 + \frac{\sqrt{3}}{2}$$

Eigenvalues and eigenvectors

10/10 points (100%)

Quiz, 10 questions



1 / 1
points

8.

Select the two eigenvectors of the matrix $A = \begin{bmatrix} 3/2 & -1 \\ -1/2 & 1/2 \end{bmatrix}$.



$$\mathbf{v}_1 = \begin{bmatrix} 1 - \sqrt{3} \\ 1 \end{bmatrix}, \mathbf{v}_2 = \begin{bmatrix} 1 + \sqrt{3} \\ 1 \end{bmatrix}$$



$$\mathbf{v}_1 = \begin{bmatrix} 1 - \sqrt{5} \\ 1 \end{bmatrix}, \mathbf{v}_2 = \begin{bmatrix} 1 + \sqrt{5} \\ 1 \end{bmatrix}$$



$$\mathbf{v}_1 = \begin{bmatrix} -1 - \sqrt{5} \\ 1 \end{bmatrix}, \mathbf{v}_2 = \begin{bmatrix} -1 + \sqrt{5} \\ 1 \end{bmatrix}$$



$$\mathbf{v}_1 = \begin{bmatrix} -1 - \sqrt{3} \\ 1 \end{bmatrix}, \mathbf{v}_2 = \begin{bmatrix} -1 + \sqrt{3} \\ 1 \end{bmatrix}$$



Correct

These are the eigenvectors for the matrix A . They have the eigenvalues λ_1 and λ_2 respectively.



1 / 1
points

9.

Form the matrix C whose left column is the vector \mathbf{v}_1 and whose right column is \mathbf{v}_2 from immediately above.

By calculating $D = C^{-1}AC$ or by using another method, find the diagonal matrix D .



$$\begin{bmatrix} -1 - \frac{\sqrt{3}}{2} & 0 \\ 0 & -1 + \frac{\sqrt{3}}{2} \end{bmatrix}$$



Eigenvalues and eigenvectors

10/10 points (100%)

Quiz, 10 questions



$$\begin{bmatrix} 1 + \frac{\sqrt{3}}{2} & 0 \\ 0 & 1 - \frac{\sqrt{3}}{2} \end{bmatrix}$$

**Correct**

Well done! Recall that when a matrix is transformed into its diagonal form, the entries along the diagonal are the eigenvalues of the matrix - this can save lots of calculation!



$$\begin{bmatrix} -1 - \frac{\sqrt{5}}{2} & 0 \\ 0 & -1 + \frac{\sqrt{5}}{2} \end{bmatrix}$$

1 / 1
points

10.

By using the diagonalisation above or otherwise, calculate A^2 .



$$\begin{bmatrix} -11/4 & 2 \\ 1 & -3/4 \end{bmatrix}$$



$$\begin{bmatrix} 11/4 & -1 \\ -2 & 3/4 \end{bmatrix}$$



$$\begin{bmatrix} -11/4 & 1 \\ 2 & -3/4 \end{bmatrix}$$



$$\begin{bmatrix} 11/4 & -2 \\ -1 & 3/4 \end{bmatrix}$$

**Correct**

Well done! In this particular case, calculating A^2 directly is probably easier - so always try to look for the method which solves the question with the least amount of pain possible!

Eigenvalues and eigenvectors

10/10 points (100%)Quiz, 10 questions

