

MOOC Econometrics


Lecture 3.1 on Model Specification: Motivation

Michel van der Wel

Erasmus University Rotterdam



Example: Stock market index



Introduction

- Question 1: Do we include all explanatory variables or only a few?
- Question 2: Should we transform the variables?
- Question 3: How to evaluate a model?

Test

Suppose that all explanatory variables in a dataset are relevant for the dependent variable. Should we include all?

Answer: Not necessarily. Lecture 3.2 will explain the why and how.




Explanatory variables

- Stock characteristics: Dividends, earnings, volatility, book value, issuing activity
- Interest-rate related: Treasury bill rates, long term yields, corporate bond returns
- Macroeconomic: Inflation, investment, consumption




Stock index and book-to-market ratio




Lecture 3.1, Slide 5 of 10, Erasmus School of Economics

Log stock index and book-to-market ratio



Lecture 3.1, Slide 6 of 10, Erasmus School of Economics

Change in log stock index, and book-to-market ratio



Lecture 3.1, Slide 7 of 10, Erasmus School of Economics

Regression output

Dependent variable: change in log(SP500 index)

Sample size: 86

	Coefficient b_j	Standard error $SE(b_j)$	t-Statistic t_j	p-value $H_0: \beta_j = 0$
Constant	0.177	0.050	3.543	0.001
Book-to-market	-0.213	0.079	-2.702	0.008
R-squared	0.080			
SE of regression	0.191			


Test

What is the interpretation of the negative sign of Book-to-market?

Answer: High book-to-market usually coincides with periods when market-value decreased.

Erasmus

Lecture 3.1, Slide 8 of 10, Erasmus School of Economics



Lecture 3.1, Slide 9 of 10, Erasmus School of Economics

- Train yourself by making the training exercise (see the website).
- After making this exercise, check your answers by studying the webcast solution (also available on the website).

Lecture 3.1, Slide 10 of 10, Erasmus School of Economics

MOOC Econometrics

Lecture 3.2 on Model Specification:
Specification

Michel van der Wel

Erasmus University Rotterdam



Consequences of omitting variables

DGP: $y = X_1\beta_1 + X_2\beta_2 + \varepsilon \rightarrow b_1 \text{ and } b_2$ Model: $y = X_1\beta_1 + \tilde{\varepsilon} \rightarrow b_R$

Test

Express $E(b_R)$ as function of β_1 and β_2 .

Answer:

$$\begin{aligned}
 E(b_R) &= E((X_1'X_1)^{-1}X_1'y), \\
 &= E((X_1'X_1)^{-1}X_1'(X_1\beta_1 + X_2\beta_2 + \varepsilon)), \\
 &= E((X_1'X_1)^{-1}X_1'X_1\beta_1 + (X_1'X_1)^{-1}X_1'X_2\beta_2 + (X_1'X_1)^{-1}X_1'\varepsilon)), \\
 &= \beta_1 + (X_1'X_1)^{-1}X_1'X_2\beta_2 + 0.
 \end{aligned}$$



Bias-efficiency trade-off

Setting:

$$y_i = x_i'\beta + \varepsilon_i, \quad i = 1, \dots, n,$$

or

$$y = X\beta + \varepsilon$$

in matrix form.

Which variables should we include in X ?

- Too few variables \rightarrow Bias.
- Too many variables \rightarrow Efficiency loss.
(Even if all variables really matter!)



Consequences of omitting variables

DGP: $y = X_1\beta_1 + X_2\beta_2 + \varepsilon \rightarrow b_1 \text{ and } b_2$ Model: $y = X_1\beta_1 + \tilde{\varepsilon} \rightarrow b_R$

It holds:

- $E(b_R) = \beta_1 + \underbrace{(X_1'X_1)^{-1}X_1'X_2}_P \beta_2 = \beta_1 + P\beta_2$
 \rightarrow Bias if $\beta_2 \neq 0$ (omitted variable bias).
- $Var(b_R) = Var(b_1) - PVar(b_2)P'$
 \rightarrow Variance of b_R is smaller than that of b_1 (even if $\beta_2 = 0$!).



Decision metrics

Possible decision metrics:

- Information criteria
- Out-of-sample prediction




Lecture 3.2, Slide 5 of 9, Erasmus School of Economics

Out-of-sample prediction

Commonly used out-of-sample prediction metrics:

- $RMSE = \left(\frac{1}{n_f} \sum_{i=1}^{n_f} (y_i - \hat{y}_i)^2 \right)^{1/2}$
- $MAE = \frac{1}{n_f} \sum_{i=1}^{n_f} |y_i - \hat{y}_i|$

with n_f the number of observations “saved” for out-of-sample evaluation and \hat{y}_i the i -th predicted value of the dependent variable.



Lecture 3.2, Slide 7 of 9, Erasmus School of Economics

Information criteria

Commonly used information criteria:

- Akaike: $AIC = \log(s^2) + \frac{2k}{n}$
- Bayes: $BIC = \log(s^2) + \frac{k \log n}{n}$

with s the standard error of the regression and k the number of variables.

Test

Which information criterion imposes the strongest penalty on the number of variables?

Answer: Penalty is $2/n$ for AIC and $\log(n)/n$ for BIC; BIC imposes stronger penalty if $\log(n) > 2$, $n \geq 8$.



Lecture 3.2, Slide 6 of 9, Erasmus School of Economics

Iterative selection methods

Commonly used methods to select explanatory variables:

- t -test and F -test
- Information criteria
- Out-of-sample predictions

Also iterative methods (based on tests) are commonly used:

- General-to-specific / backward elimination
- Specific-to-general / forward selection



Lecture 3.2, Slide 8 of 9, Erasmus School of Economics

- Train yourself by making the training exercise (see the website).
- After making this exercise, check your answers by studying the webcast solution (also available on the website).



MOOC Econometrics

Lecture 3.3 on Model Specification: Transformation

Michel van der Wel

Taking logarithms

Use for:

- Exponential growth.

Data transformation

Setting:

$$y_i = x_i' \beta + \varepsilon_i, \quad i = 1, \dots, n,$$

or

$$y = X\beta + \varepsilon$$

in matrix form.

What is the most appropriate form of the data (y and X)?

- All variables should be incorporated in a compatible manner.
- If this is not the case, data can be transformed.

Taking differences

Use for:

- Trending patterns
→ Statistical assumptions may not hold.

First difference:

$$\Delta y_i = y_i - y_{i-1}.$$

Test

What is the result if you take the difference of $y_i = i$?

Answer: For $y_i = i$ the difference is:

$$\Delta y_i = y_i - y_{i-1} = i - (i - 1) = 1.$$

Non-linear effects

$$y_i = x_i' \beta + \varepsilon_i = \beta_1 + \sum_{j=2}^k \beta_j x_{ji} + \varepsilon_i, \quad i = 1, \dots, n,$$

has linear set-up with fixed marginal effects ($dy_i/dx_{ji} = \beta_j$).

Extension with interaction and quadratic terms:

$$y_i = \beta_1 + \sum_{j=2}^k \beta_j x_{ji} + \sum_{j=2}^k \gamma_{jj} x_{ji}^2 + \sum_{j=2}^k \sum_{h=j+1}^k \gamma_{jh} x_{ji} x_{hi} + \varepsilon_i.$$

Advantages:

- Get non-linear functional form.
- May provide economically meaningful specification.

Erasmus

Lecture 3.3, Slide 5 of 9, Erasmus School of Economics

Dummy variables

Quarterly data

$$y_i = \alpha_i + \sum_{j=2}^k \beta_j x_{ji} + \varepsilon_i, \quad i = 1, \dots, n,$$

where α_i is the quarter-specific mean level.

Use dummy variables:

- D_{hi} for $h = 1, 2, 3, 4$, with $D_{hi} = 1$ if observation i is in quarter h (and $D_{hi} = 0$ otherwise).

- Then

$$y_i = \alpha_1 D_{1i} + \alpha_2 D_{2i} + \alpha_3 D_{3i} + \alpha_4 D_{4i} + \sum_{j=2}^k \beta_j x_{ji} + \varepsilon_i.$$

Erasmus

Lecture 3.3, Slide 7 of 9, Erasmus School of Economics

Example (from lecture 2.2)

- $\log(\text{Wage})_i = \beta_1 + \beta_2 \text{Female}_i + \beta_3 \text{Age}_i + \beta_4 \text{Educ}_i + \beta_5 \text{Parttime}_i + \varepsilon_i.$

- $\log(\text{Wage})_i = \beta_1 + \beta_2 \text{Female}_i + \beta_3 \text{Age}_i + \beta_4 \text{Educ}_i + \beta_5 \text{Parttime}_i + \gamma_1 \text{Female}_i \text{Educ}_i + \gamma_2 \text{Age}_i^2 + \varepsilon_i.$

Now:

- Wage differential may depend on education ($\beta_2 + \gamma_1 \text{Educ}_i$).
- Age has non-linear effect for wage ($\beta_3 + 2\gamma_2 \text{Age}_i$).

Erasmus

Lecture 3.3, Slide 6 of 9, Erasmus School of Economics

Dummy variables

$$y_i = \alpha_1 D_{1i} + \alpha_2 D_{2i} + \alpha_3 D_{3i} + \alpha_4 D_{4i} + \sum_{j=2}^k \beta_j x_{ji} + \varepsilon_i.$$

Test

Can we add a constant term to the above specification that has a dummy for each quarter?

Answer: Only if one of the dummy parameters is set to 0.

If we omit D_{1i} , so $\alpha_1 = 0$, we get

$$y_i = \alpha_1 + \gamma_2 D_{2i} + \gamma_3 D_{3i} + \gamma_4 D_{4i} + \sum_{j=2}^k \beta_j x_{ji} + \varepsilon_i,$$

where $\gamma_h = \alpha_h - \alpha_1$ for $h = 2, 3, 4$.

Erasmus

Lecture 3.3, Slide 8 of 9, Erasmus School of Economics

- Train yourself by making the training exercise (see the website).
- After making this exercise, check your answers by studying the webcast solution (also available on the website).



MOOC Econometrics

Lecture 3.4 on Model Specification:
Evaluation

Michel van der Wel

Erasmus University Rotterdam



RESET

Instead, add fitted y -values $\hat{y} = Xb = X(X'X)^{-1}X'y$ to the model:

$$y_i = x_i'\beta + \sum_{j=1}^p \gamma_j (\hat{y}_i)^{j+1} + \varepsilon_i,$$

and test for joint significance of γ 's. Under null of correct specification, $H_0: \gamma_j = 0$ for all j , test distribution approximately $F(p, n - k - p)$.

Test

For $p = 1$, compute the number of extra parameters in the alternative specification as compared to the total number of parameters in the RESET specification.

Answer: Above model with $p = 1$ has $k + 1$ parameters. Model with squares and cross-terms has $k + (k - 1) + \frac{1}{2}(k - 2)(k - 1)$ coefficients. For example, if $k = 6$, then this is 7 compared to 21.



RESET

Extend linear model

$$y_i = \beta_1 + \sum_{j=2}^k \beta_j x_{ji} + \varepsilon_i,$$

to non-linear model

$$y_i = \beta_1 + \sum_{j=2}^k \beta_j x_{ji} + \sum_{j=2}^k \gamma_{jj} x_{ji}^2 + \sum_{j=2}^k \sum_{h=j+1}^k \gamma_{jh} x_{ji} x_{hi} + \varepsilon_i.$$

Test for linearity by testing significance of γ coefficients.

Challenge: Nonlinear model contains many parameters.



Chow break test

In case of a possible break, split the sample and test for constancy of parameters.

$$\begin{aligned} y_1 &= X_1 \beta_1 + \varepsilon_1 & (n_1 \text{ observations}) \\ y_2 &= X_2 \beta_2 + \varepsilon_2 & (n_2 = n - n_1 \text{ observations}) \end{aligned}$$

Combine:

$$\begin{pmatrix} y_1 \\ y_2 \end{pmatrix} = \begin{pmatrix} X_1 & 0 \\ 0 & X_2 \end{pmatrix} \begin{pmatrix} \beta_1 \\ \beta_2 \end{pmatrix} + \begin{pmatrix} \varepsilon_1 \\ \varepsilon_2 \end{pmatrix}$$

Test $H_0: \beta_1 = \beta_2$ against this unrestricted set-up.



Chow break test

F-test for null hypothesis of no break:

$$F = \frac{(e'_R e_R - e'_U e_U)/k}{e'_U e_U/(n-2k)}.$$

Here:

- Have $e_U = \begin{pmatrix} e_1 \\ e_2 \end{pmatrix}$, with e_j the OLS residuals of each group.
- Thus $e'_U e_U = e'_1 e_1 + e'_2 e_2 \equiv S_1 + S_2$.
- Get $F = \frac{(S_0 - S_1 - S_2)/k}{(S_1 + S_2)/(n-2k)}$, with $S_0 = e'_R e_R$.




Lecture 3.4, Slide 5 of 8, Erasmus School of Economics

Test for normality of error terms

- Model misspecification may appear in the error terms.
- Normality of ε can be tested by distribution of residuals.
- Jarque-Bera test evaluates skewness S and kurtosis K :

$$JB = \left(\sqrt{\frac{n}{6}} S \right)^2 + \left(\sqrt{\frac{n}{24}} (K - 3) \right)^2,$$

which approximately has $\chi^2(2)$ distribution if $H_0: \varepsilon_i \sim NID(0, \sigma^2)$ holds true.



Lecture 3.4, Slide 7 of 8, Erasmus School of Economics

Chow forecast test

A variation on the Chow break test is based on

$$y_i = x'_i \beta + \sum_{j=n_1+1}^{n_1+n_2} \gamma_j D_{ji} + \varepsilon_i,$$

test $H_0: \gamma_j = 0$ for all j .

Test

What is the number of parameters in the above specification?

Answer: The model contains the usual k variables and n_2 dummy-variables (one for each observation in group 2), so in total $k + n_2$ parameters.

- Perfect fit in second sample, thus $e_2 = 0$.
- Thus $F = \frac{(S_0 - S_1)/n_2}{S_1/(n_1 - k)}$.



Lecture 3.4, Slide 6 of 8, Erasmus School of Economics

TRAINING EXERCISE 3.4

- Train yourself by making the training exercise (see the website).
- After making this exercise, check your answers by studying the webcast solution (also available on the website).



Lecture 3.4, Slide 8 of 8, Erasmus School of Economics

MOOC Econometrics


Lecture 3.5 on Model Specification: Application

Michel van der Wel

Erasmus University Rotterdam

Erasmus

Transformation



Lecture 3.5, Slide 3 of 9, Erasmus School of Economics

Setting

Application:

- Model/forecast S&P500 stock index
 - ▶ Should we transform this index series?
- Large set of explanatory variables
 - ▶ Which to select?
- Choice of model
 - ▶ How to evaluate candidate models and how to compare them?
 - ▶ Is relationship stable over time?

Erasmus

Lecture 3.5, Slide 2 of 9, Erasmus School of Economics

Variable selection

Dependent variable: Log of equity premium; Sample size: 87

	Coefficients (p-values) by specification				
	(A)	(B)	(C)	(D)	(E)
Constant	0.166 (0.001)	0.062 (0.015)	-0.266 (0.076)	-0.027 (0.848)	0.065 (0.015)
Book-to-market	-0.185 (0.019)				
Issued Stock		-0.147 (0.850)			
Dividend/Price			-0.097 (0.029)		
Earnings/price				-0.032 (0.532)	
Inflation					-0.166 (0.746)
R-squared	0.063	0.000	0.055	0.005	0.001

Erasmus

Lecture 3.5, Slide 4 of 9, Erasmus School of Economics

General to specific

Dependent variable: Log of equity premium; Sample size: 87					
	Coefficients (p-values) by specification				
	(1)	(2)	(3)	(4)	(5)
Constant	0.234 (0.544)	0.205 (0.554)	0.215 (0.537)	0.489 (0.038)	0.166 (0.001)
Book-to-market	-0.176 (0.257)	-0.166 (0.249)	-0.191 (0.178)	-0.290 (0.008)	-0.185 (0.019)
Issued Stock	-0.146 (0.859)				
Dividend/Price	-0.120 (0.226)	-0.126 (0.174)	-0.090 (0.286)		
Earnings/price	0.167 (0.052)	0.167 (0.051)	0.127 (0.088)	0.097 (0.159)	
Inflation	-0.567 (0.337)	-0.564 (0.336)			
R-squared	0.108	0.108	0.098	0.085	0.063

Erasmus

Lecture 3.5, Slide 5 of 9, Erasmus School of Economics

Model comparison

	Full model	Book-to-market
R^2	0.108	0.063
AIC	-0.444	-0.486
BIC	-0.273	-0.430

Erasmus

Lecture 3.5, Slide 7 of 9, Erasmus School of Economics

Stability

$$\log(\text{EqPr})_i = \beta_1 + \beta_2 \text{BTM}_i + \beta_3 \text{BTM}_i \times D_i^{\text{War}} + \beta_4 \text{BTM}_i \times D_i^{\text{Oil}} + \varepsilon_i,$$

Dependent variable: Log of equity premium; Sample size: 87		
	Coefficients	p-values
Constant	0.160	0.002
Book-to-market	-0.175	0.036
Book-to-market \times War-dummy	0.078	0.440
Book-to-market \times Oil-dummy	-0.133	0.287
R-squared	0.085	

Test

What is the coefficient on Book-to-market during the war years?

Answer: $-0.175 + 0.078 = -0.097$.

Erasmus

Lecture 3.5, Slide 6 of 9, Erasmus School of Economics

Model evaluation

Book-to-market		
	Test statistic	p-value
RESET ($p = 1$)	3.446	0.067
Chow Break	2.269	0.110
Chow Forecast	0.765	0.794
Jarque-Bera	7.155	0.028

Note: As break-point 1980 is chosen.

Test

Will the p-values increase if the full model is considered?

Answer: Not possible to say.

Erasmus

Lecture 3.5, Slide 8 of 9, Erasmus School of Economics

- Train yourself by making the training exercise (see the website).
- After making this exercise, check your answers by studying the webcast solution (also available on the website).

