

HOME DOWNLOAD SALES EBOOK SITE MAP

# Kruskal-Wallis Test

A collection of data samples are **independent** if they come from unrelated populations and the samples do not affect each other. Using the **Kruskal-Wallis Test**, we can decide whether the population distributions are identical *without* assuming them to follow the **normal distribution**.

### **Example**

In the built-in data set named **airquality**, the daily air quality measurements in New York, May to September 1973, are recorded. The ozone density are presented in the data frame column Ozone.

```
> head(airquality)
  Ozone Solar.R Wind Temp Month Day
1    41    190   7.4   67   5   1
2    36    118   8.0   72   5   2
.....
```

#### **Problem**

Without assuming the data to have normal distribution, test at .05 significance level if the monthly ozone density in New York has identical data distributions from May to September 1973

#### **Solution**

The null hypothesis is that the monthly ozone density are identical populations. To test the hypothesis, we apply the kruskal test function to compare the independent monthly data. The p-value turns out to be nearly zero (6.901e-06). Hence we reject the null hypothesis.

```
> kruskal.test(Ozone ~ Month, data = airquality)

Kruskal-Wallis rank sum test

data: Ozone by Month
Kruskal-Wallis chi-squared = 29.267, df = 4, p-value = 6.901e-06
```

#### **Answer**


At .05 significance level, we conclude that the monthly ozone density in New York from May to September 1973 are *nonidentical* populations.

< Mann-Whitney-Wilcoxon Test	up Kruskal-Wallis test	Simple Linear Regression >		
Tags: Elementary Statistics with R airquality		non-parametric	head	kruskal.test

### Search this site:

Search

## R Tutorial eBook



# R Tutorials

R Tutorials
R Introduction
Elementary Statistics with R
Qualitative Data
Quantitative Data
Numerical Measures
Probability Distributions
Interval Estimation
Hypothesis Testing
Type II Error
Inference About Two
Populations
Goodness of Fit
Analysis of Variance
Non-parametric Methods
Sign Test
Wilcoxon Signed-Rank Test
Mann-Whitney-Wilcoxon Test
Kruskal-Wallis Test
Simple Linear Regression
Multiple Linear Regression
Logistic Regression

GPU Computing with R

## Recent Articles

- Installing CUDA Toolkit 7.5 on Fedora 21 Linux
   September 10, 2015
- Installing CUDA Toolkit 7.5 on Ubuntu 14.04 Linux
   September 10, 2015
- Hierarchical Linear Model
   July 22, 2013
- Bayesian Classification with Gaussian Process
   January 6, 2013

Copyright © 2009 - 2016 Chi Yau All Rights Reserved Theme design by styleshout Fractal graphics by zyzstar Adaptation by Chi Yau