What properties should a probability measure have?

Sum of face values of a pair of dice: 50 throws

Sum of face values of a pair of dice: 50 throws



Sum of face values of a pair of dice: 50 throws



Sample space: $\Omega = \{2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12\}$

Win: $A = \{7, 11\}$

Lose: $B = \{2, 3, 12\}$

Win or lose: $A \cup B = \{2, 3, 7, 11, 12\}$

Sum of face values of a pair of dice: 50 throws



Frequency of occurrence: $\nu(\cdot)$

Sample space:
$$\Omega = \{2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12\}$$

Win:
$$A = \{7, 11\}$$

Lose:
$$B = \{2, 3, 12\}$$

Win or lose:
$$A \cup B = \{2, 3, 7, 11, 12\}$$

$$\nu(\Omega) = \frac{50}{50} = 1$$

$$v(A) = \frac{9}{50}$$

$$\nu(B) = \frac{7}{50}$$

$$\nu(A \cup B) = \frac{16}{50} = \frac{9}{50} + \frac{7}{50} = \nu(A) + \nu(B)$$

Sum of face values of a pair of dice: 50 throws



Frequency of occurrence: $\nu(\cdot)$

Sample space: $\Omega = \{2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12\}$

Win:
$$A = \{7, 11\}$$

Lose: $B = \{2, 3, 12\}$

Win or lose: $A \cup B = \{2, 3, 7, 11, 12\}$

$$\nu(\Omega) = \frac{50}{50} = 1$$

$$v(A) = \frac{9}{50}$$

$$\nu(B) = \frac{7}{50}$$

$$\nu(A \cup B) = \frac{16}{50} = \frac{9}{50} + \frac{7}{50} = \nu(A) + \nu(B)$$

Normalisation: The certain event

Positivity

Additivity: A and B are disjoint, $A \cap B = \emptyset$