

Lesson 5.3-5.4

[Back to Week 2](#)

8/8 points earned
(100%)

Quiz passed!



Be Recognized for Your Achievements. "Course Certificates give you the recognition you need to get the job, the material gives you the skills to do the job. It makes you look more valuable because you are more valuable." - Peter B., USA, Software Developer

Showcase Your Accomplishment! Earn Your Course Certificate! ₹1,953 ➔



1 / 1
points

1.

We use the continuous version of Bayes' theorem if:



θ is continuous

Correct Response

If θ is continuous, we use a probability density for the prior.

- ☐ Y is continuous
- ☐ $f(y | \theta)$ is continuous
- ☐ All of the above
- ☐ None of the above



1 / 1
points

2.

Consider the coin-flipping example from the lesson. Recall that the likelihood for this experiment was Bernoulli with unknown probability of heads, i.e., $f(y | \theta) = \theta^y (1 - \theta)^{1-y} I_{\{0 \leq \theta \leq 1\}}$, and we started with a uniform prior on the interval $[0, 1]$.

After the first flip resulted in heads ($Y_1 = 1$), the posterior for θ became $f(\theta | Y_1 = 1) = 2\theta I_{\{0 \leq \theta \leq 1\}}$.

Now use this posterior as your prior for θ before the next (second) flip. Which of the following represents the posterior PDF for θ after the second flip also results in heads ($Y_2 = 1$)?

- ☐ $f(\theta | Y_2 = 1) = \frac{(1-\theta) \cdot 2\theta}{\int_0^1 (1-\theta) \cdot 2\theta d\theta} I_{\{0 \leq \theta \leq 1\}}$
- ☐ $f(\theta | Y_2 = 1) = \frac{\theta(1-\theta) \cdot 2\theta}{\int_0^1 \theta(1-\theta) \cdot 2\theta d\theta} I_{\{0 \leq \theta \leq 1\}}$
- ☒ $f(\theta | Y_2 = 1) = \frac{\theta \cdot 2\theta}{\int_0^1 \theta \cdot 2\theta d\theta} I_{\{0 \leq \theta \leq 1\}}$

Correct Response

This simplifies to the posterior PDF $f(\theta | Y_2 = 1) = 3\theta^2 I_{\{0 \leq \theta \leq 1\}}$.

Incidentally, if we assume that the two coin flips are independent, we would have arrived at the same posterior if we had again started with a uniform prior and performed a single update using $Y_1 = 1$ and $Y_2 = 1$.



1 / 1
points

3.

Consider again the coin-flipping example from the lesson. Recall that we used a $\text{Uniform}(0,1)$ prior for θ . Which of the following is a correct interpretation of $P(0.3 < \theta < 0.9) = 0.6$?



(0.3, 0.9) is a 60% credible interval for θ before observing any data.



Correct Response

The probability statement came from our prior, so the prior probability that θ is in this interval is 0.6.



(0.3, 0.9) is a 60% credible interval for θ after observing $Y = 1$.



(0.3, 0.9) is a 60% confidence interval for θ .



The posterior probability that $\theta \in (0.3, 0.9)$ is 0.6.



1 / 1
points

4.

Consider again the coin-flipping example from the lesson. Recall that the posterior PDF for θ , after observing $Y = 1$, was $f(\theta | Y = 1) = 2\theta I_{\{0 \leq \theta \leq 1\}}$. Which of the following is a correct interpretation of $P(0.3 < \theta < 0.9 | Y = 1) = \int_{0.3}^{0.9} 2\theta d\theta = 0.72$?

- ☐ (0.3, 0.9) is a 72% credible interval for θ before observing any data.
- ☒ (0.3, 0.9) is a 72% credible interval for θ after observing $Y = 1$.

Correct Response

The probability statement came from the posterior, so the posterior probability that θ is in this interval is 0.72.

- ☐ (0.3, 0.9) is a 72% confidence interval for θ .
- ☐ The prior probability that $\theta \in (0.3, 0.9)$ is 0.72.



1 / 1
points

5.

Which two quantiles are required to capture the middle 90% of a distribution (thus producing a 90% equal-tailed interval)?

- ☐ .025 and .975
- ☐ 0 and .9
- ☐ .10 and .90
- ☒ .05 and .95

Correct Response

90% of the probability mass is contained between the .05 and .95 quantiles (or equivalently, the 5th and 95th percentiles). 5% of the probability lies on either side of this interval.



1 / 1
points

6.

Suppose you collect measurements to perform inference about a population mean θ . Your posterior distribution after observing data is $\theta \mid \mathbf{y} \sim N(0, 1)$.

Report the upper end of a 95% equal-tailed interval for θ . Round your answer to two decimal places.

1.96

Correct Response

The 95% equal-tailed interval for a standard normal distribution is (-1.96, 1.96).

Because the normal distribution is symmetric and unimodal (has only one peak), the equal-tailed interval is also the highest posterior density (HPD) interval.

In R:

```
1 qnorm(p=0.975, mean=0, sd=1)
```

In Excel:

```
1 = NORM.INV(0.975, 0, 1)  
2
```

where probability=0.975, mean=0, standard_dev=1.



1 / 1
points

7.

What does "HPD interval" stand for?

- ☐ Highest precision density interval
- ☐ Highest point distance interval
- ☒ Highest posterior density interval

Correct Response

- ☐ Highest partial density interval
-




1 / 1
points


8.

Each of the following graphs depicts a 50% credible interval from a posterior distribution. Which of the intervals represents the HPD interval?


- ☐ 50% interval: $\theta \in (0.500, 1.000)$



☐ 50% interval: $\theta \in (0.196, 0.567)$



☒ 50% interval: $\theta \in (0.326, 0.674)$



Correct Response

This is the 50% credible interval with the highest posterior density values. It is the shortest possible interval containing 50% of the probability under this posterior distribution.

☐ 50% interval: $\theta \in (0.400, 0.756)$

