Expectation and variance in continuous spaces

The continuous sample space

The continuous sample space

Random variable $X \sim p(x)$ density

The continuous sample space

Random variable $X \sim p(x)$ density

$$p(x) \geq 0, \quad \int_{-\infty}^{\infty} p(x)\, dx = 1$$

The continuous sample space

Random variable $X \sim p(x)$ density

$$p(x) \geq 0, \quad \int_{-\infty}^{\infty} p(x)\, dx = 1$$

$$\mu = \int_{-\infty}^{\infty} x \cdot p(x)\, dx$$

The continuous sample space

Random variable $X \sim p(x)$ density

$$p(x) \geq 0, \quad \int_{-\infty}^{\infty} p(x)\, dx = 1$$

The expectation of X (if the integral converges absolutely):

$$\mu = \int_{-\infty}^{\infty} x \cdot p(x)\, dx \;=: \mathsf{E}(X)$$

The continuous sample space

Random variable $X \sim p(x)$ density

$$p(x) \geq 0, \quad \int_{-\infty}^{\infty} p(x)\, dx = 1$$

The expectation of X (if the integral converges absolutely):

$$\mu = \int_{-\infty}^{\infty} x \cdot p(x)\, dx \;=: \mathsf{E}(X)$$

$$\sigma^2 = \int_{-\infty}^{\infty} (x-\mu)^2 \cdot p(x)\, dx$$

The continuous sample space

Random variable $X \sim p(x)$ density

$$p(x) \geq 0, \quad \int_{-\infty}^{\infty} p(x)\, dx = 1$$

The expectation of X (if the integral converges absolutely):

$$\mu = \int_{-\infty}^{\infty} x \cdot p(x)\, dx \ =: \mathsf{E}(X)$$

The variance of X:

$$\sigma^2 = \int_{-\infty}^{\infty} (x - \mu)^2 \cdot p(x)\, dx \ =: \mathrm{Var}(X)$$

Coordinate transformations: one dimension

Random variable $X \sim p(x)$ density

Coordinate transformations: one dimension

Random variable $X \sim p(x)$ density

Transformation: $Y = f(X)$

Coordinate transformations: one dimension

Random variable $X \sim p(x)$ density

Transformation: $Y = f(X)$

$$m = \int_{-\infty}^{\infty} f(x) \cdot p(x)\, dx$$

Coordinate transformations: one dimension

Random variable $X \sim p(x)$ density

Transformation: $Y = f(X)$

$$m = \int_{-\infty}^{\infty} f(x) \cdot p(x)\, dx =: \mathbb{E}\big(f(X)\big)$$

Coordinate transformations: one dimension

Random variable $X \sim p(x)$ density

Transformation: $Y = f(X)$

change of variable formula

$$m = \int_{-\infty}^{\infty} f(x) \cdot p(x)\, dx =: E\big(f(X)\big)$$

Coordinate transformations: one dimension

Random variable $X \sim p(x)$ density

Transformation: $Y = f(X)$

change of variable formula

$$m = \int_{-\infty}^{\infty} f(x) \cdot p(x)\, dx =: \mathrm{E}\big(f(X)\big)$$

$$s^2 = \int_{-\infty}^{\infty} \big(f(x) - m\big)^2 \cdot p(x)\, dx$$

Coordinate transformations: one dimension

Random variable $X \sim p(x)$ density

Transformation: $Y = f(X)$

change of variable formula

$$m = \int_{-\infty}^{\infty} f(x) \cdot p(x)\, dx \;=: \mathsf{E}\big(f(X)\big)$$

$$s^2 = \int_{-\infty}^{\infty} \big(f(x) - m\big)^2 \cdot p(x)\, dx \;=: \mathrm{Var}\big(f(X)\big)$$

Coordinate transformations: two and more dimensions

Coordinate transformations: two and more dimensions

Random vector $(X_1, \ldots, X_n) \sim p(x_1, \ldots, x_n)$ density

Coordinate transformations: two and more dimensions

Random vector $(X_1, \ldots, X_n) \sim p(x_1, \ldots, x_n)$ density

Transformation: $Y = f(X_1, \ldots, X_n)$

Coordinate transformations: two and more dimensions

Random vector $(X_1, \dots, X_n) \sim p(x_1, \dots, x_n)$ density

Transformation: $Y = f(X_1, \dots, X_n)$

$$m = \int_{-\infty}^{\infty} \cdots \int f(x_1, \dots, x_n) \cdot p(x_1, \dots, x_n) \, dx_n \cdots dx_1$$

Coordinate transformations: two and more dimensions

Random vector $(X_1, \ldots, X_n) \sim p(x_1, \ldots, x_n)$ density

Transformation: $Y = f(X_1, \ldots, X_n)$

$$m = \int_{-\infty}^{\infty} \cdots \int f(x_1, \ldots, x_n) \cdot p(x_1, \ldots, x_n)\, dx_n \cdots dx_1 \; =: \mathsf{E}\big(f(X_1, \ldots, X_n)\big)$$

Coordinate transformations: two and more dimensions

Random vector $(X_1, \ldots, X_n) \sim p(x_1, \ldots, x_n)$ density

Transformation: $Y = f(X_1, \ldots, X_n)$

$$m = \int_{-\infty}^{\infty} \cdots \int f(x_1, \ldots, x_n) \cdot p(x_1, \ldots, x_n)\, dx_n \cdots dx_1 \;=: \mathsf{E}\big(f(X_1, \ldots, X_n)\big)$$

$$s^2 = \int_{-\infty}^{\infty} \cdots \int \big(f(x_1, \ldots, x_n) - m\big)^2 \cdot p(x_1, \ldots, x_n)\, dx_n \cdots dx_1$$

Coordinate transformations: two and more dimensions

Random vector $(X_1, \dots, X_n) \sim p(x_1, \dots, x_n)$ density

Transformation: $Y = f(X_1, \dots, X_n)$

$$m = \int_{-\infty}^{\infty} \cdots \int f(x_1, \dots, x_n) \cdot p(x_1, \dots, x_n)\, dx_n \cdots dx_1 \;=: \mathsf{E}\big(f(X_1, \dots, X_n)\big)$$

$$s^2 = \int_{-\infty}^{\infty} \cdots \int \big(f(x_1, \dots, x_n) - m\big)^2 \cdot p(x_1, \dots, x_n)\, dx_n \cdots dx_1 =: \mathrm{Var}\big(f(X_1, \dots, X_n)\big)$$